

**CAPRON MAX- dextromethorphan hbr and pyrilamine maleate liquid**  
**Capital Pharmaceutical, LLC**

-----  
**Capron DM Liquid**

***Active Ingredients***

(in each 5 mL teaspoonful)

Dextromethorphan HBr 15 mg

Pyrilamine Maleate 15 mg

***Purpose***

Dextromethorphan HBr 15 mg ..... Antitussive

Pyrilamine Maleate 15 mg ..... Antihistamine

***Uses***

temporarily relieves these symptoms due to hay fever (allergic rhinitis) or other upper respiratory allergies:

- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes
- cough due to minor throat and bronchial irritation associated with a cold
- alleviates cough to help you sleep
- nonnarcotic cough suppressant for the relief of cough

***Warnings***

**Do not exceed recommended dosage.**

- a persistent cough may be a sign of a serious condition. If cough persist for more than 1 week, tends to recur, or is accompanied by a fever, rash, or persistent headache, consult a doctor.

**Do not use this product**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product
- for persistent or chronic cough such as occurs with smoking, asthma, or emphysema, or if cough is accompanied by excessive phlegm (mucus) unless directed by a doctor

**Ask a doctor before use if you have**

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- difficulty in urination due to enlargement of the prostate gland

**Ask a doctor before use if you are taking  
sedatives or tranquilizers.**

**When using this product**

- excitability may occur, especially in children
- may cause marked drowsiness
- sedatives and tranquilizers may increase drowsiness effect
- avoid alcoholic beverages
- use caution when driving a motor vehicle or operating machinery

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of accidental overdose, seek professional help or contact a Poison Control Center immediately.

***Directions***

**Do not exceed recommended dosage.**

Adults and children 12 years of age and over:	2 teaspoonfuls every 6-8 hours, not to exceed 8 teaspoonfuls in 24 hours, or as directed by a doctor
Children 6 to under 12 years of age:	1 teaspoonfuls every 6-8 hours, not to exceed 4 teaspoonfuls in 24 hours, or as directed by a doctor
Children under 6 years of age	Consult a doctor

***Inactive ingredients***

Citric acid, flavor, methylparaben, monoammonium glycyrrhizinate, potassium citrate, potassium sorbate, propylene glycol, propylparaben, purified water, sorbitol, sucralose

***Questions or Comments?***

Call 614.638.4622

**PRINCIPAL DISPLAY PANEL**

Serialization Area

NDC 29978-123-16

Capron MAX

Liquid

Antitussive • Antihistamine

Sugar Free • Alcohol Free

Dye Free • Gluten Free

Each teaspoonful (5 mL) for oral administration contains:


Dextromethorphan HBr..... 15 mg

Pyrilamine Maleate ..... 15 mg

Orange Pineapple Flavor

Tamper evident by foil seal under cap. Do not use if foil seal is broken or missing.

Supplied in a tight container with a child-resistant cap.



Manufactured for:  
Capital Pharmaceutical, LLC  
Powell, OH 43065  
1 (614) 638-4622

16 oz. (473 mL)

Drug Facts

Active Ingredients (in each 5 mL teaspoonful)

Dextromethorphan HBr 15 mg ..... Antitussive

Pyrilamine Maleate 15 mg ..... Antihistamine

Purpose

Uses temporarily relieves these symptoms due to hay fever (allergic rhinitis) or other upper respiratory allergies: runny nose sneezing itching of the nose or throat itchy, watery eyes cough due to minor throat and bronchial irritation associated with a cold alleviates cough to help you sleep nonnarcotic cough suppressant for the relief of cough

Warnings Do not exceed recommended dosage. a persistent cough may be a sign of a serious condition. If cough persist for more than 1 week, tends to recur, or is accompanied by a fever, rash, or persistent headache, consult a doctor.

Do not use this product if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product for persistent or chronic cough such as occurs with smoking, asthma, or emphysema, or if cough is accompanied by excessive phlegm (mucus) unless directed by a doctor

Ask a doctor before use if you have a breathing problem such as emphysema or chronic bronchitis glaucoma difficulty in urination due to enlargement of the prostate gland

Ask a doctor before use if you are taking sedatives or tranquilizers.

Drug Facts (continued)

When using this product excitability may occur, especially in children may cause marked drowsiness sedatives and tranquilizers may increase drowsiness effect avoid alcoholic beverages use caution when driving a motor vehicle or operating machinery

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of accidental overdose, seek professional help or contact a Poison Control Center immediately.

Directions Do not exceed recommended dosage.


Adults and children 12 years of age and over: 2 teaspoonfuls every 6-8 hours, not to exceed 8 teaspoonfuls in 24 hours, or as directed by a doctor

Children 6 to under 12 years of age: 1 teaspoonful every 6-8 hours, not to exceed 4 teaspoonfuls in 24 hours, or as directed by a doctor

Children under 6 years of age: consult a doctor

Inactive Ingredients Citric acid, flavor, methylparaben, monoammonium glycyrrhizate, potassium citrate, potassium sorbate, propylene glycol, propylparaben, purified water, sorbitol, sucralose

Questions or Comments? Call 614.638.4622



N 3 29978 12316 1

Rev. 01/25

## CAPRON MAX

dextromethorphan hbr and pyrilamine maleate liquid

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:29978-123
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg in 5 mL
PYRILAMINE MALEATE (UNII: R35D29L3ZA) (PYRILAMINE - UNII:HPE317O9TL)	PYRILAMINE MALEATE	15 mg in 5 mL

### Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
AMMONIUM GLYCYRRHIZATE (UNII: 3VRD35U26C)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
POTASSIUM CITRATE (UNII: EE90ONI6FF)	
POTASSIUM SORBATE (UNII: 1VPU26JZZ4)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

<b>WATER</b> (UNII: 059QF0KO0R)				
<b>SORBITOL</b> (UNII: 506T60A25R)				
<b>SUCRALOSE</b> (UNII: 96K6UQ3Z D4)				
<b>Product Characteristics</b>				
<b>Color</b>				<b>Score</b>
<b>Shape</b>				<b>Size</b>
<b>Flavor</b>	ORANGE (Orange Pineapple) , PINEAPPLE (Orange Pineapple)			<b>Imprint Code</b>
<b>Contains</b>				
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
<b>1</b>	NDC:29978-123-16	1 in 1 BOX	06/11/2025	
<b>1</b>		473 mL in 1 BOTTLE; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
<b>Marketing Category</b>		<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug		M012	06/11/2025	

**Labeler -** Capital Pharmaceutical, LLC (831545541)