

NYSTATIN - nystatin cream
Dispensing Solutions, Inc.

NYSTATIN CREAM, USP


Rx only

FOR TOPICAL USE ONLY • NOT FOR OPHTHALMIC USE

DESCRIPTION

Nystatin is a polyene antifungal antibiotic obtained from *Streptomyces noursei*.

Structural formula:



$C_{47}H_{75}NO_{17}$

MW 926.13

Nystatin cream is for dermatologic use.

Nystatin cream for topical use, contains 100,000 USP nystatin units per gram. Inactive ingredients: aluminum hydroxide compressed wet gel, cetearyl alcohol (and) cetareth 20, glyceryl monostearate, polyoxyl 40 stearate, propylene glycol, purified water, simethicone, sorbic acid, sorbitol solution, titanium dioxide, and white petrolatum.

CLINICAL PHARMACOLOGY

Pharmacokinetics

Nystatin is not absorbed from intact skin or mucous membrane.

Microbiology

Nystatin is an antibiotic which is both fungistatic and fungicidal *in vitro* against a wide variety of yeasts and yeast-like fungi, including *Candida albicans*, *C. parapsilosis*, *C. tropicalis*, *C. guilliermondi*, *C. pseudotropicalis*, *C. krusei*, *Torulopsis glabrata*, *Trichophyton rubrum*, *T. mentagrophytes*.

Nystatin acts by binding to sterols in the cell membrane of susceptible species resulting in a change in membrane permeability and the subsequent leakage of intracellular components. On repeated subculturing with increasing levels of nystatin, *Candida albicans* does not develop resistance to nystatin. Generally, resistance to nystatin does not develop during therapy. However, other species of *Candida* (*C. tropicalis*, *C. guilliermondi*, *C. krusei*, and *C. stellatoidea*) become quite resistant on treatment with nystatin and simultaneously become cross resistant to amphotericin as well. This resistance is lost when the antibiotic is removed.

Nystatin exhibits no appreciable activity against bacteria, protozoa, or viruses.

INDICATIONS AND USAGE

Nystatin cream is indicated in the treatment of cutaneous or mucocutaneous mycotic infections caused by *Candida albicans* and other susceptible *Candida* species.

This cream is not indicated for systemic, oral, intravaginal or ophthalmic use.

CONTRAINDICATIONS

Nystatin cream is contraindicated in patients with a history of hypersensitivity to any of its components.

PRECAUTIONS

General

Nystatin cream should not be used for the treatment of systemic, oral, intravaginal or ophthalmic infections.

If irritation or sensitization develops, treatment should be discontinued and appropriate measures taken as indicated. It is recommended that KOH smears, cultures, or other diagnostic methods be used to confirm the diagnosis of cutaneous or mucocutaneous candidiasis and to rule out infection caused by other pathogens.

INFORMATION FOR THE PATIENT

Patients using this medication should receive the following information and instructions:

1. The patient should be instructed to use this medication as directed (including the replacement of missed doses). This medication is not for any disorder other than that for which it is prescribed.
2. Even if symptomatic relief occurs within the first few days of treatment, the patient should be advised not to interrupt or discontinue therapy until the prescribed course of treatment is completed.
3. If symptoms of irritation develop, the patient should be advised to notify the physician promptly.

Laboratory Tests

If there is a lack of therapeutic response, KOH smears, cultures, or other diagnostic methods should be repeated.

Carcinogenesis, Mutagenesis, Impairment of Fertility

No long-term animal studies have been performed to evaluate the carcinogenic potential of nystatin. No studies have been performed to determine the mutagenicity of nystatin or the effects on male or female fertility.

Pregnancy:

Teratogenic Effects

Category C.

Animal reproduction studies have not been conducted with any nystatin cream. It also is not known whether this cream can cause fetal harm when used by a pregnant woman or can affect reproductive capacity. Nystatin cream should be prescribed for a pregnant woman only if the potential benefit to the mother outweighs the potential risk to the fetus.

Nursing Mothers

It is not known whether nystatin is excreted in human milk. Caution should be exercised when nystatin is prescribed for a nursing woman.

Pediatric Use

Safety and effectiveness have been established in the pediatric population from birth to 16 years.
(See **DOSAGE AND ADMINISTRATION**.)

Geriatric Use

Clinical studies with nystatin cream did not include sufficient numbers of subjects aged 65 years and older to determine whether they respond differently than younger subjects. Other reported clinical experience has not identified differences in responses between elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out.

ADVERSE REACTIONS

The frequency of adverse events reported in patients using nystatin cream is less than 0.1%. The more common events that were reported include allergic reactions, burning, itching, rash, eczema, and pain on application.

(See **PRECAUTIONS: General**.)

DOSAGE AND ADMINISTRATION

Adults and Pediatric Patients (Neonates and Older):

Apply liberally to affected areas twice daily or as indicated until healing is complete.

HOW SUPPLIED

Nystatin cream: 100,000 units nystatin per gram in an aqueous, vanishing cream base, in 15 g, 30 g, and 60 g tubes.

STORAGE

Store at room temperature, avoid freezing.

Manufactured for:

QUALITEST PHARMACEUTICALS

Huntsville, AL 35811

8181604

R5/08-R2

PRINCIPAL DISPLAY PANEL

BULK SOURCE DATA
MFD. FOR: QUALITEST PHARMACEUTICALS
HUNTSVILLE, AL 35811
PRODUCT ID:
TAN / BLACK / BROWN / WHITE
BOX PRINTED NDC 0603-7818-78
BULK SOURCE NDC: 00603-7818-78
MFR. LOT: XXXXXX
PEDIGREE #: 815503
DISPENSE IN THIS
TIGHT/LIGHT RESISTANT CONTAINER



APPLY EXTERNALLY TO AFFECTED AREA(S) EVERY _____ HOURS _____ TIMES A DAY. NOT FOR OPHTHALMIC USE. FOR DERMATOLOGIC USE ONLY. AVOID FREEZING. AVOID EXPOSURE TO EXCESSIVE HEAT.

DispenseQuick™
Making Medicine Easy

NYSTATIN 100,000 USP units/gram
30g CREAM
NDC 68258-3046-03
PRODUCT # 3N0126

Rev. Date: 04/11

EACH GRAM CONTAINS:
100,000 USP NYSTATIN UNITS IN
AN AQUEOUS VANISHING CREAM BASE

LOT# SAMPLE EXP: 00-00 Rx # 24193121
RX ONLY

WARNING: KEEP OUT OF CHILDREN'S REACH
STORE AT ROOM TEMPERATURE. SEE USP.

3N0126 NDC 68258-3046-03
NYSTATIN 100,000 USP units/gram
30g CREAM
LOT # SAMPLE EXP: 00-00
MN 00603-7818-78 RX# 24193121

3N0126 NDC 68258-3046-03
NYSTATIN 100,000 USP units/gram
30g CREAM
LOT # SAMPLE EXP: 00-00
MN 00603-7818-78 RX# 24193121

3N0126 NDC 68258-3046-03
NYSTATIN 100,000 USP units/gram
30g CREAM
LOT # SAMPLE EXP: 00-00
MN 00603-7818-78 RX# 24193121



Packaged Exclusively By:
DISPENSING SOLUTIONS^{INC}
Santa Ana, CA 92704

NDC 68258-3046-03

NYSTATIN				
nystatin cream				
Product Information				
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:68258-3046(NDC:0603-7818)	
Route of Administration	TOPICAL			
Active Ingredient/Active Moiety				
	Ingredient Name	Basis of Strength	Strength	
	NYSTATIN (UNII: BDF1O1C72E) (NYSTATIN - UNII:BDF1O1C72E)	NYSTATIN	100000 [USP.U] in 1g	
Inactive Ingredients				
	Ingredient Name	Strength		
	ALUMINUM HYDRO XIDE (UNII: 5QB0T2IUN0)			
	CETOSTEARYL ALCOHOL (UNII: 2DMT128M1S)			
	GLYCERYL MONOSTEARATE (UNII: 230OU9XXE4)			
	POLYOXYL 40 STEARATE (UNII: 13A4J4NH9I)			
	PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
	WATER (UNII: 059QF0KO0R)			
	SORBIC ACID (UNII: X045WJ989B)			
	SORBITOL (UNII: 506T60A25R)			
	TITANIUM DIOXIDE (UNII: 15FIX9V2JP)			
	PETROLATUM (UNII: 4T6H12BN9U)			
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date

1	NDC:68258-3046-3	30 g in 1 TUBE		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA065315	05/31/2006		

Labeler - Dispensing Solutions, Inc. (066070785)

Establishment			
Name	Address	ID/FEI	Business Operations
Dispensing Solutions, Inc.		066070785	relabel, repack

Revised: 10/2011

Dispensing Solutions, Inc.