

LICEOUT- piperonyl butoxide and pyrethrum extract shampoo
Bob Barker Company Inc.

LiceOut™

Active Ingredients:

*Piperonyl Butoxide,
Technical.....3.0% Pyrethrum
Extract.....0.3%

*Equivalent to min. 2.4% (butylcarbityl) (6-propylpiperonyl) ether and 0.6% related compounds.

Active Ingredients	Purpose
Piperonyl Butoxide, Technical 3.0%	Lice Treatment
Pyrethrum Extract 0.3%	Lice Treatment
*Equivalent to min. 2.4% (butylcarbityl) (6-propylpiperonyl) ether and 0.6% related compounds.	

Uses: Treats head, pubic (crab) and body lice.

Warnings: For external use only.

Do not use • near eyes • inside nose, mouth, or vagina • on lice in eyebrows or eyelashes. See a doctor if lice are present in these areas.

Ask a doctor before use if you are • allergic to ragweed. May cause breathing difficulty or an asthmatic attack. When using this product • keep eyes tightly closed and protect eyes with a washcloth or towel • if product gets in eyes, flush with water right away • scalp itching or redness may occur.

Stop use and ask a doctor if • breathing difficulty occurs • eye irritation occurs • skin or scalp irritation continues or infection occurs. Do not freeze. Do not reuse original container.

Children under 2 years ask a doctor

Do not use

- near eyes
- inside nose, mouth, or vagina
- on lice in eyebrows or eyelashes. See a doctor if lice are present in these areas.

Ask a doctor before use if you are

- allergic to ragweed.
- if product gets in eyes, flush with water right away
- scalp itch-ing or redness may occur.

Stop use and ask a doctor if

- breathing difficulty occurs
- eye irritation occurs
- skin or scalp irritation continues or infection occurs

Children under 2 yearsask a doctor.

Caution

Keep Out of the Reach of Children

Directions

Shake Well Before Using

Adults and children 2 years and over.

Treat

- Apply thoroughly to hair or other affected area. For head lice, first apply behind ears and to back of neck.
- Allow product to remain for 10 minutes, but no longer
- use warm water to form a lather, shampoo, then thoroughly rinse
- For head lice, towel dry hair and comb out tangles.
- Check daily for any lice/nits.
- A second treatment must be done in 7 to 10 days to kill any newly hatched lice.
- If infestation continues, see a doctor for other treatments.

Other Information

Other Information

See insert in carton for additional information.

LICE OUT[®]

Non-foaming Shampoo Liquid Lice Treatment for Human Use

Kills

Head Lice (*Pediculus Humanus capitis* DeG)

Body Lice (*Pediculus Humanus Humanus* L.)

Pubic (Crab) Lice (*Phthirus pubis* L.)

Caution

Keep Out of the Reach of Children

Contents: 1 FL. OZ. (29.6 mL)

Active Ingredients:

*Piperonyl Butoxide, Technical.....	3.0%
Pyrethrum Extract.....	0.3%

Total Ingredients:

Water, Octylphenoxypolyethoxyethanol, Piperonyl Butoxide, Benzyl Alcohol, C 13-14 Isoparaffin, Pyrethrum Extract. *Equivalent to min. 2.4% (butylcarbityl) (6-propylpiperonyl) ether and 0.6% related compounds.

Uses: Treats head, pubic (crab) and body lice.

Warnings:

For external use only. Do not use • near eyes • inside nose, mouth, or vagina • on lice in eyebrows or eyelashes. See a doctor if lice are present in these areas. Ask a doctor before use if you are • allergic to ragweed. May cause breathing difficulty or an asthmatic attack. When using this product • keep eyes tightly closed and protect eyes with a washcloth or towel • if product gets in eyes, flush with water right away • scalp itching or redness may occur. Stop use and ask a doctor if • breathing difficulty occurs • eye irritation occurs • skin or scalp irritation continues or infection occurs. Do not freeze. Do not reuse original container.

Directions

Important: Read Warnings Before Use

Adults and children 2 years and over. Inspect • check each household member with a magnifying glass in bright light for lice/nits (eggs) • look for tiny nits near scalp, beginning at back of neck and behind ears • examine small sections of hair at a time • unlike dandruff which moves when touched, nits stick to the hair • if either lice or nits are found, treat with this product.

– SHAKE WELL BEFORE USING –

Treat • Apply thoroughly to hair or other affected area. For head lice, first apply behind ears and to back of neck. • Allow product to remain for 10 minutes, but no longer • use warm water to form a lather, shampoo, then thoroughly rinse • For head lice, towel dry hair and comb out tangles. Remove lice and their eggs (nits) • Use a fine-tooth or special lice/nit comb. Remove any remaining nits by hand (using a throw-away glove). Hair should remain slightly damp while removing nits • if hair dries during combing, dampen slightly with water • for head lice, part hair into sections. Do one section at a time starting on top of head. Longer hair may take 1 to 2 hours. • Lift a 1- to 2-inch wide strand of hair. Place comb as close to scalp as possible and comb with a firm, even motion away from scalp. • Pin back each strand of hair after combing • clean comb often. Wipe nits away with tissue and discard in a plastic bag. Seal bag and discard to prevent lice from coming back. • After combing, thoroughly recheck for lice/nits. Repeat combing if necessary. • Check daily for any lice/nits that you missed. • A second treatment must be done in 7 to 10 days to kill any newly hatched lice. • If infestation continues, see a doctor for other treatments.

Children under 2 years ask a doctor.

Other Information:

Head lice • lay small white eggs (nits) on hair shaft close to scalp. • Nits are most easily found on back of neck or behind ears. • Disinfect hats, hair ribbons, scarves, coats, towels and bed linens by machine washing in hot water [above 54° C (130° F)], then using hottest dryer cycle for at least 20 minutes. • Items that cannot be washed (bedspreads, blankets, pillows, stuffed toys, etc.) should be dry-cleaned or sealed in a plastic bag for 4 weeks, then removed outdoors and shaken out very hard before using again. • Items that cannot be washed, dry-cleaned, or stored may be sprayed with a product designed for this purpose. • Soak all combs and brushes in hot water [above 54° C (130° F)] for at least 10 minutes. • Vacuum all carpets, mattresses, upholstered furniture, and car seats that may have been used by affected people. **Pubic (crab) lice** • may be transmitted by sexual contact. Sexual partners should be treated simultaneously to avoid reinfection. • Lice are very small and look like brown or grey dots on skin • usually cause intense itching and lay small white eggs (nits) on the hair shaft generally close to the skin surface • may be present on the short hairs of groin, thighs, trunk, and underarms, and occasionally on the beard and mustache. • Disinfect underwear by machine washing in hot water [above 54° C (130° F)], then using hottest dryer cycle for at least 20 minutes. **Body lice** • body lice and their eggs (nits) are generally found in the seams of clothing, particularly in waistline and armpit area. • Body lice feed on skin then return to clothing to lay their eggs. • Disinfect clothing by machine washing in hot water [above 54° C (130° F)], then using hottest dryer cycle for at least 20 minutes. • Do not seal clothing in a plastic bag because nits can remain dormant for up to 30 days.

Distributed By: Bob Barker Company, Inc.
7925 Purfoy Road Fuquay-Varina, NC 27526

Inactive Ingredients

Water, Octylphenoxypolyethoxyethanol, Piperonyl Butoxide, Benzyl Alcohol, C 13-14 Isoparaffin, Pyrethrum Extract.

Distributed By: Bob Barker Company, Inc.
7925 Purfoy Road, Fuquay-Varina, NC 27526

PRINCIPAL DISPLAY PANEL - 150 Sachet Case LICEOUT

LICEOUT

Non-foaming Shampoo
Liquid Lice Treatment for Human Use

Kills

Head Lice (*Pediculus Humanus capitis* DeG)

Body Lice (*Pediculus Humanus Humanus* L.)

Pubic (Crab) Lice (*Phthirus pubis* L.)

Caution

Keep Out of the Reach of Children

Contents: 1 FL. OZ. (29.6 mL)

LICEOUT[®]

Drug Facts (continued)

Stop use and ask a doctor if • breathing difficulty occurs • eye irritation occurs • skin or scalp irritation continues or infection occurs.

Children under 2 years ask a doctor.

Directions

Shake Well Before Using

Adults and children 2 years and over.

Treat • Apply thoroughly to hair or other affected area. For head lice, first apply behind ears and to back of neck. • Allow product to remain for 10 minutes, but no longer • use warm water to form a lather, shampoo, then thoroughly rinse • For head lice, towel dry hair and comb out tangles. • Remove lice and their eggs (nits). • Check daily for any lice/nits. • A second treatment must be done in 7 to 10 days to kill any newly hatched lice. • If infestation continues, see a doctor for other treatments.

Other Information

See insert in carton for additional information.

Inactive Ingredients

Water, Octylphenoxypolethoxyethanol, Piperonyl Butoxide, Benzyl Alcohol, C 13-14 Isoparaffin, Pyrethrum Extract.

Distributed By: Bob Barker Company, Inc.
7925 Purfoy Road, Fuquay-Varina, NC 27526

LICEOUT[®]

Non-foaming Shampoo
Liquid Lice Treatment for Human Use

Kills

Head Lice (*Pediculus Humanus capitis* DeG)

Body Lice (*Pediculus Humanus Humanus* L.)

Pubic (Crab) Lice (*Phthirus pubis* L.)

Caution

Keep Out of the Reach of Children

Contents: 1 FL. OZ. (29.6 mL)

Drug Facts

Active Ingredients

Purpose

Piperonyl Butoxide, Technical 3.0%...Lice Treatment
Pyrethrum Extract 0.3%.....Lice Treatment
*Equivalent to min. 2.4% (butylcarbityl) (6-propylpiperonyl) ether and 0.6% related compounds.

Uses Treats head, pubic (crab) and body lice

Warnings

For external use only.

Do not use • near eyes • inside nose, mouth, or vagina • on lice in eyebrows or eyelashes. See a doctor if lice are present in these areas.

Ask a doctor before use if you are • allergic to ragweed. • if product gets in eyes, flush with water right away • scalp itching or redness may occur.



LICEOUT

Non-foaming Shampoo
Liquid Lice Treatment for Human Use

Kills
Head Lice (Pediculus Humanus capitis DeG)
Body Lice (Pediculus Humanus Humanus L.)
Pubic (Crab) Lice (Phthirus pubis L.)

Caution
Keep Out of the Reach of Children

Contents: 128 FL. OZ. (3785.4 mL)



Non-foaming Shampoo
Liquid Lice Treatment for Human Use
Kills
Head Lice (Pediculus Humanus capitis DeG)
Body Lice (Pediculus Humanus Humanus L.)
Pubic (Crab) Lice (Phthirus pubis L.)
Caution
Keep Out of the Reach of Children
Contents: 128 FL. OZ. (3785.4 mL)

Active Ingredients:
Piperonyl Butoxide, Technical 3.0%
Pyrethrum Extract 0.3%

Total Ingredients:
Water, Octylphenoxypolyethoxyethanol, Piperonyl Butoxide, Benzyl Alcohol, C 13-14 Isoparaffin, Pyrethrum Extract, *Equivalent to min. 2.4% (butylcarbityl) (6-propylpiperonyl) ether and 0.6% related compounds.

Uses: Treats head, pubic (crab) and body lice.

Warnings:
For external use only. Do not use • near eyes • inside nose, mouth, or vagina • on lice in eyebrows or eyelashes. See a doctor if lice are present in these areas. Ask a doctor before use if you are allergic to ragweed. May cause breathing difficulty or an asthmatic attack. When using this product • keep eyes tightly closed and protect eyes with a washcloth or towel • if product gets in eyes, flush with water right away • scalp itching or redness may occur. Stop use and ask a doctor if • breathing difficulty occurs • eye irritation occurs • skin or scalp irritation continues or infection occurs. Do not freeze. Do not reuse original container.

Directions
Important: Read Warnings Before Use
Adults and children 2 years and over. Inspect • check each household member with a magnifying glass in bright light for lice(nits (eggs)) • look for tiny nits near scalp, beginning at back of neck and behind ears • examine small sections of hair at a time • unlike dandruff which moves when touched, nits stick to the hair • if either lice or nits are found, treat with this product.
– **SHAKE WELL BEFORE USING** –
Treat • Apply thoroughly to hair or other affected area. For head lice, first apply behind ears and to back of neck • Allow product to remain for 10 minutes, but no longer • use warm water to form a lather, shampoo, then thoroughly rinse • For head lice, towel dry hair and comb out tangles. Remove lice and their eggs (nits) • Use a fine-tooth or special lice(nit) comb. Remove any remaining nits by hand using a throw-away glove. Hair should remain slightly damp while removing nits • if hair dries during combing, dampen slightly with water • for head lice, part hair into sections. Do one section at a time starting on top of head. Longer hair may take 1 to 2 hours.
• Lift a 1- to 2-inch wide strand of hair. Place comb as close to scalp as possible and comb with a firm, even motion away from scalp • Pin back each strand of hair after combing • clean comb often. Wipe nits away with tissue and discard in a plastic bag. Seal bag and discard to prevent lice from coming back • After combing, thoroughly recheck for lice(nits). Repeat combing if necessary.
• Check daily for any lice(nits) that you missed • A second treatment must be done in 7 to 10 days to kill any newly hatched lice • If infestation continues, see a doctor for other treatments.
Children under 2 years ask a doctor.
Other Information:
Head lice • lay small white eggs (nits) on hair shaft close to scalp • Nits are most easily found on back of neck or behind ears • Disinfest hats, hair ribbons, scarves, coats, towels and bed linens by machine washing in hot water [above 54° C (130° F)], then using hottest dryer cycle for at least 20 minutes • Items that cannot be washed (bedspreads, blankets, pillows, stuffed toys, etc.) should be dry-cleaned or sealed in a plastic bag for 4 weeks, then removed outdoors and shaken out very hard before using again • Items that cannot be washed, dry-cleaned, or stored may be sprayed with a product designed for this purpose • Soak all combs and brushes in hot water [above 54° C (130° F)] for at least 10 minutes • Vacuum all carpets, mattresses, upholstered furniture, and car seats that may have been used by affected people. **Pubic (crab) lice** • may be transmitted by sexual contact. Sexual partners should be treated simultaneously to avoid reinfestation • Lice are very small and look like brown or grey dots on skin • usually cause intense itching and lay small white eggs (nits) on the hair shaft generally close to the skin surface • may be present on the short hairs of groin, thighs, trunk, and underarms, and occasionally on the beard and mustache • Disinfest underwear by machine washing in hot water [above 54° C (130° F)], then using hottest dryer cycle for at least 20 minutes. **Body lice** • body lice and their eggs (nits) are generally found in the seams of clothing, particularly in waistline and armpit area • Body lice feed on skin then return to clothing to lay their eggs • Disinfest clothing by machine washing in hot water [above 54° C (130° F)], then using hottest dryer cycle for at least 20 minutes • Do not seal clothing in a plastic bag because nits can remain dormant for up to 30 days.



Drug Facts	
Active Ingredients Piperonyl Butoxide, Technical 3.0% Pyrethrum Extract 0.3% *Equivalent to min. 2.4% (butylcarbityl) (6-propylpiperonyl) ether and 0.6% related compounds.	Purpose Lice Treatment Lice Treatment Lice Treatment
Uses Treats head, pubic (crab) and body lice	
Warnings For external use only. Do not use • near eyes • inside nose, mouth, or vagina • on lice in eyebrows or eyelashes. See a doctor if lice are present in these areas. Ask a doctor before use if you are • allergic to ragweed • if product gets in eyes, flush with water right away • scalp itching or redness may occur. Stop use and ask a doctor if • breathing difficulty occurs • eye irritation occurs • skin or scalp irritation continues or infection occurs. Children under 2 years ask a doctor.	
Directions Shake Well Before Using Adults and children 2 years and over. Treat • Apply thoroughly to hair or other affected area. For head lice, first apply behind ears and to back of neck • Allow product to remain for 10 minutes, but no longer • use warm water to form a lather, shampoo, then thoroughly rinse • For head lice, towel dry hair and comb out tangles. • Check daily for any lice(nits). • A second treatment must be done in 7 to 10 days to kill any newly hatched lice • If infestation continues, see a doctor for other treatments.	
Inactive Ingredients Water, Octylphenoxypolyethoxyethanol, Piperonyl Butoxide, Benzyl Alcohol, C 13-14 Isoparaffin, Pyrethrum Extract. Distributed By: Bob Barker Company, Inc. 7925 Purfoy Road, Fuquay-Varina, NC 27526	

LICEOUT
piperonyl butoxide and pyrethrum extract shampoo

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53247-124
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety		
Ingredient Name		Strength
PIPERONYL BUTOXIDE (UNII: LWK91TU9AH) (PIPERONYL BUTOXIDE - UNII:LWK91TU9AH)		3 g in 100 mL
PYRETHRUM EXTRACT (UNII: ZUM06L90GV) (PYRETHRUM EXTRACT - UNII:ZUM06L90GV)		0.3 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
C13-14 ISOPARAFFIN (UNII: E4F12ROE70)	
OCTOXYNOL-9 (UNII: 7JPC6Y25QS)	
WATER (UNII: 059QF0KO0R)	
BENZYL ALCOHOL (UNII: LKG8494WBH)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53247-124-01	150 in 1 CASE	01/02/2022	09/05/2025
1		29.6 mL in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:53247-124-02	3785 mL in 1 JUG; Type 0: Not a Combination Product	01/02/2022	04/11/2026

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	01/02/2022	04/11/2026

Labeler - Bob Barker Company Inc. (058525536)

Revised: 11/2024

Bob Barker Company Inc.