

NYSTATIN- nystatin suspension
ATLANTIC BIOLOGICALS CORP.


Nystatin Oral Suspension USP

[100,000 units per mL]

Rx Only

DESCRIPTION

Nystatin is an antimycotic polyene antibiotic obtained from *Streptomyces noursei*. Structural formula:



C₄₇H₇₅NO₁₇ MW 926.13

Nystatin Oral Suspension USP, for oral administration, contains 100,000 USP Nystatin Units per mL. Inactive ingredients: alcohol ($\leq 1\%$ v/v), artificial peppermint flavor, cherry flavor, citric acid, D&C Yellow No. 10, FD&C Red No. 40, glycerin, magnesium aluminum silicate, methylparaben, potassium phosphate dibasic, propylene glycol, propylparaben, purified water and sucrose.

CLINICAL PHARMACOLOGY

Pharmacokinetics

Gastrointestinal absorption of nystatin is insignificant. Most orally administered nystatin is passed unchanged in the stool. In patients with renal insufficiency receiving oral therapy with conventional dosage forms, significant plasma concentrations of nystatin may occasionally occur.

Microbiology

Nystatin is both fungistatic and fungicidal *in vitro* against a wide variety of yeasts and yeast-like fungi. *Candida albicans* demonstrates no significant resistance to nystatin *in vitro* on repeated subculture in increasing levels of nystatin; other *Candida* species become quite resistant. Generally, resistance does not develop *in vivo*. Nystatin acts by binding to sterols in the cell membrane of susceptible *Candida* species with a resultant change in membrane permeability allowing leakage of intracellular components. Nystatin exhibits no appreciable activity against bacteria, protozoa, or viruses.

INDICATIONS AND USAGE

Nystatin oral suspension is indicated for the treatment of candidiasis in the oral cavity.

CONTRAINDICATIONS

The preparation is contraindicated in patients with a history of hypersensitivity to any of its components.

PRECAUTIONS

General

This medication is not to be used for the treatment of systemic mycoses. Discontinue treatment if sensitization or irritation is reported during use.

Carcinogenesis, Mutagenesis, Impairment of Fertility

No long-term animal studies have been performed to evaluate carcinogenic potential. There also have been no studies to determine mutagenicity or whether this medication affects fertility in males or females.

Pregnancy

Teratogenic Effects Category C

Animal reproduction studies have not been conducted with nystatin oral suspension. It is also not known whether nystatin oral suspension can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Nystatin oral suspension should be given to a pregnant woman only if clearly needed.

Nursing Mothers

It is not known whether nystatin is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when nystatin is administered to a nursing woman.

Pediatric Use

See DOSAGE AND ADMINISTRATION.

ADVERSE REACTIONS

Nystatin is well tolerated even with prolonged therapy. Oral irritation and sensitization have been reported. (See PRECAUTIONS: General).

Gastrointestinal: Diarrhea (including one case of bloody diarrhea), nausea, vomiting, gastrointestinal upset/disturbances.

Dermatologic: Rash, including urticaria has been reported rarely. Stevens-Johnson syndrome has been reported very rarely.

Other: Tachycardia, bronchospasm, facial swelling, and non-specific myalgia have also been rarely reported.

OVERDOSAGE

Oral doses of nystatin in excess of five million units daily have caused nausea and gastrointestinal upset. There have been no reports of serious toxic effects of superinfections (See CLINICAL PHARMACOLOGY, Pharmacokinetics).

DOSAGE AND ADMINISTRATION

INFANTS: 2 mL (200,000 units) four times daily (in infants and young children, use dropper to place one-half of dose in each side of mouth and avoid feeding for 5 to 10 minutes).

NOTE: Limited clinical studies in premature and low birth weight infants indicate that 1 mL four times daily is effective.

CHILDREN AND ADULTS: 4 to 6 mL (400,000 to 600,000 units) four times daily (one-half of dose in each side of mouth). The preparation should be retained in the mouth as long as possible before swallowing.

Continue treatment for at least 48 hours after perioral symptoms have disappeared and cultures demonstrate eradication of *Candida albicans*.

HOW SUPPLIED

Nystatin Oral Suspension USP, 100,000 USP Nystatin Units per mL, is available in a cherry, peppermint flavored, light creamy yellow, ready-to-use suspension, supplied in the following oral dosage forms:

NDC 17856-0811-01 NYSTATIN 100,000 UNITS/ML 0.5ML ENFIT SYRINGE 120 ct UD

NDC 17856-0811-02 NYSTATIN 100,000 UNITS/ML 1ML ENFIT SYRINGE 120 ct UD

NDC 17856-0811-03 NYSTATIN 100,000 UNITS/ML 2ML ENFIT SYRINGE 120 ct UD

NDC 17856-0811-04 NYSTATIN 100,000 UNITS/ML 5ML ENFIT SYRINGE 48 ct UD

NDC 17856-0811-05 NYSTATIN 100,000 UNITS/ML 5ML CUP 72 ct UD

Storage

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature]. Avoid freezing.

DISTRIBUTED BY:

ATLANTIC BIOLOGICALS CORP

20101 N.E 16TH PLACE

MIAMI, FL 33179

PRINCIPAL DISPLAY PANEL

NDC 17856-0811

NYSTATIN

ORAL

SUSPENSION, USP

[100,000 units per mL]

Alcohol ≤ 1% v/v SHAKE WELL

Rx ONLY

FOR INSTITUTIONAL USE ONLY

SEE INSERT

17856-0811-01



Rev.01/19

**NYSTATIN ORAL SUSPENSION
50,000 UNITS/0.5 ML ENFIT SYR**

See package insert for indications and dosage schedule

Store at 20° to 25°C (68° to 77°F).[See USP Controlled Room Temperature]. Protect from light. **AVOID FREEZING**
**** Keep this and all Medication out of the reach of children

17856-0811-01
NYSTATIN

Dosage: 0.5 ML

Qty: 120 ENFIT Syringes



GTIN: 1785608111
S/N: 00000000000001
Exp: 10/08/19
Lot: 000000

RX



17856081101

Packaged by: Unit Dose Solutions
Morrisville, NC 27560

Distributed by: AtlanticBiologicals
Corp, Miami FL 33179

Call to Reorder: 800.509.7592

17856-0811-02



**NYSTATIN ORAL SUSPENSION
100,000 UNITS/ML ENFIT SYR**

Rev.01/19

See package insert for indications and dosage schedule

Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature]. Protect from light. AVOID FREEZING
**** Keep this and all Medication out of the reach of children

17856-0811-02
NYSTATIN

Dosage: 1 ML

Qty: 120 ENFIT Syringes



GTIN: 1785608112
S/N: 00000000000002
Exp: 10/08/19
Lot: 000000

RX



17856081102

Packaged by: Unit Dose Solutions
Morrisville, NC 27560

Distributed by: AtlanticBiologicals
Corp, Miami Fl 33179

Call to Reorder: 800.509.7592

17856-0811-03



**NYSTATIN ORAL SUSPENSION
200,000 UNITS/2 ML ENFIT SYR**

Rev.01/19

See package insert for indications and dosage schedule

Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature]. Protect from light. **AVOID FREEZING**
**** Keep this and all Medication out of the reach of children

17856-0811-03
NYSTATIN

Dosage: 2 ML

Qty: 120 ENFIT Syringes



GTIN: 1785608113
S/N: 00000000000003
Exp: 10/08/19
Lot: 000000



17856081103

Packaged by: Unit Dose Solutions
Morrisville, NC 27560

Distributed by: AtlanticBiologicals
Corp, Miami Fl 33179

Call to Reorder: 800.509.7592

17856-0811-04



Rev.01/19

**NYSTATIN ORAL SUSPENSION
500,000 UNITS/5 ML ENFIT SYR**

See package insert for indications and dosage schedule

Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature]. Protect from light. **AVOID FREEZING**
*** Keep this and all Medication out of the reach of children

17856-0811-04
NYSTATIN

Dosage: 5 ML

Qty: 48 ENFIT Syringes



GTIN: 1785608114
S/N: 00000000000004
Exp: 10/08/19
Lot: 000000



17856081104

Packaged by: Unit Dose Solutions
Morrisville, NC 27560
Distributed by: AtlanticBiologicals
Corp, Miami FL 33179

Call to Reorder: 800.509.7592

17856-0811-05



Rev.01/19

NYSTATIN ORAL SUSPENSION
500,000 UNITS/5 ML

See package insert for indications and dosage schedule

Store at 20° to 25°C (68° to 77°F).[See USP Controlled Room Temperature]. Protect from light. AVOID FREEZING
**** Keep this and all Medication out of the reach of children

17856-0811-05
NYSTATIN

Dosage: 5 ML

Qty: 72 Cups



GTIN: 1785608115
S/N: 00000000000005
Exp: 10/08/19
Lot: 000000



17856081105

Packaged by: Unit Dose Solutions
Morrisville, NC 27560

Distributed by: AtlanticBiologicals
Corp, Miami FL 33179

Call to Reorder: 800.509.7592

NYSTATIN

nystatin suspension

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:17856-0811(NDC:0121-0810)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NYSTATIN (UNII: BDF1O1C72E) (NYSTATIN - UNII:BDF1O1C72E)	NYSTATIN	100000 [USP'U] in 1 mL

Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	

CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
METHYL PARABEN (UNII: A2I8C7HI9T)	
POTASSIUM PHOSPHATE, DIBASIC (UNII: CI71S98N1Z)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYL PARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SUCROSE (UNII: C151H8M554)	
MAGNESIUM ALUMINUM SILICATE (UNII: 6M3P64V0NC)	

Product Characteristics

Color	yellow (Light - Creamy)	Score	
Shape		Size	
Flavor	CHERRY (w/Peppermint)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:17856-0811-1	120 in 1 BOX, UNIT-DOSE	01/26/2021	
1		0.5 mL in 1 SYRINGE; Type 0: Not a Combination Product		
2	NDC:17856-0811-2	120 in 1 BOX, UNIT-DOSE	01/26/2021	
2		1 mL in 1 SYRINGE; Type 0: Not a Combination Product		
3	NDC:17856-0811-3	120 in 1 BOX, UNIT-DOSE	01/26/2021	
3		2 mL in 1 SYRINGE; Type 0: Not a Combination Product		
4	NDC:17856-0811-4	48 in 1 BOX, UNIT-DOSE	01/26/2021	
4		5 mL in 1 SYRINGE; Type 0: Not a Combination Product		
5	NDC:17856-0811-5	72 in 1 BOX, UNIT-DOSE	01/26/2021	
5		5 mL in 1 CUP, UNIT-DOSE; Type 0: Not a Combination Product		
6	NDC:17856-0811-6	120 in 1 BOX, UNIT-DOSE	01/26/2021	
6		1 mL in 1 SYRINGE; Type 0: Not a Combination Product		
7	NDC:17856-0811-7	120 in 1 BOX, UNIT-DOSE	01/26/2021	
7		2 mL in 1 SYRINGE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA203621	05/07/2019	

Labeler - ATLANTIC BIOLOGICALS CORP. (047437707)

Establishment

Name	Address	ID/FEI	Business Operations
ATLANTIC BIOLOGICALS CORP.		047437707	relabel(17856-0811) , repack(17856-0811)

Revised: 1/2021

ATLANTIC BIOLOGICALS CORP.