

\_\_\_\_\_

Nuvicare NightTime Cold and Flu Softgel (Acetaminophen 325mg, Dextromethorphan HBr 15mg, Doxylamine succinate 6.25 mg)

### **Drug Facts**

## Active ingredients (in each softgel)

Acetaminophen 325mg

Dextromethorphan HBr 15mg

Doxylamine succinate 6.25 mg

### **Purpose**

Acetaminophen 325mg	Pain reliever/fever reducer
Dextromethorphan HBr 15mg	Cough suppressant
Doxylamine succinate 6.25 mg.	Antihistamine

#### Uses:

temporarily relieves these common cold/flu symptoms:

- minor aches & pains
- headache
- sore throat
- fever
- Runny nose and sneezing
- cough due to minor throat & bronchial irritation

### **Warning**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

 $\blacksquare$  more than 8 softgels in 24 hours, which is the maximum daily amount for this product  $\blacksquare$  with other drugs containing acetaminophen  $\blacksquare$  3 or more alcoholic drinks everyday while using this product.

**Allergy Alert:** acetamenophen may cause severe skin recations. Symptoms may include:

■ skin reddening ■ blisters ■ rash. if a skin reactions occurs, stop use and seek

medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### Do not use:

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

## Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that last or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

# Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedative or tranquilizers.

# When using this product

- do not exceed recommeded dosage.
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedative and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

# Stop use and ask a doctor if

- pain or cough get worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

Keep out of reach of children.

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. (1-800-222-1222)

#### **Directions**

- take only as directed- see OVERDOSE WARNING
- do not exceed 8 softgels per 24 hrs.
- take softgels with water

Adults and children age 12 yrs and over	2 softgels every 6 hrs.
Children 4 to under 12 yrs	ask a doctor
Children under 4 yrs	do not use

#### Other information

- store between 15-30°C (59 to 86 °F)
- avoid excessive heat

# **Inactive ingredients**

Polyethylene Glycol400, Propylene Glycol, Povidone (PVP K-30), Purified water, Gelatin, Glycerin, Sorbitol Sorbitan solution, FD & C Blue No. 1, D & C Yellow No. 10

## **Questions or Comments?**

Call 1-718-337-8733 or email: support@nuvicare.com

Distributed by: Nuvicare LLC, Brooklyn, NY 11211, USA



### **NUVICARE NIGHTTIME COLD AND FLU SOFTGEL**

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate capsule, liquid filled

### **Product Information**

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:84324-019
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg	
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg	
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg	

Inactive Ingredients			
Ingredient Name	Strength		
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
POVIDONE (UNII: FZ989GH94E)			
Gelatin (UNII: 2G86QN327L)			
Sorbitol (UNII: 506T60A25R)			
GLYCERIN (UNII: PDC6A3C0OX)			
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)			
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)			

Product Characteristics			
Color	green	Score	no score
Shape	capsule (Oblong)	Size	19mm
Flavor		Imprint Code	SD15
Contains			

Ш	Packaging			
	# Item Code	Package Description	Marketing Start Date	Marketing End Date
	NDC:84324- 019-01	1 in 1 CARTON	05/20/2025	
	1	10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/20/2025	

Revised: 5/2025 NUVICARE LLC