

ELAIMEI TATTOO NUMBING CREAM- lidocaine 5% cream
Shenzhen Shandian Jingling Technology Co., Ltd.

83804-005 ELAIMEI Tattoo Numbing Cream

Lidocaine 5%

Topical Anesthetic

Uses: For temporary relief of local discomfort, itching, pain, soreness or burning in the perianal area associated with anorectal disorders

Do not use this product if pregnant or breastfeeding, ask a health professional before use

Do not exceed the recommended daily dosage unless directed by a doctor.

Do not put into the rectum by using fingers or any medical device or applicator.

Children under 12 years of age, consult a doctor.

CORRECT USE STEPS BEFORE TATTOOING

1. Clean hands and treatment area with mild soap and warmwater (to open pores to help absorb numbing cream), then gently dry.
2. Apply a thick layer of about 2mm to the treated area. (Make sure the tattoo area and surroundings are white cream)
3. Wrap tightly with plastic wrap, wait 60 minutes (longer time better effect). It is recommended to massage or tap lightly every 10 minutes to help the absorption of the numbing cream.
4. Wipe away residue, wait another 20-30 minutes for the numbness to peak. And begin procedure.

Stop use and ask a doctor if Condition worsens, or does not improve within 7 days. Certain persons can develop allergic reactions to ingredients in this product. If the symptom being treated does not subside or if redness, irritation, swelling, pain, or other symptoms develop or increase, discontinue use.

KEEP OUT OF REACH OF CHILDREN SECTION

Adults and Children 12 years and older: When practical, cleanse the affected area with mild soap and warm water and rinse thoroughly, Gently dry by patting or blotting with toilet tissue or a soft cloth before application of this product. Apply externally to the affected area up to 6 times daily.

Aloe barbadensis leaf juice (Aloe vera)

Water

Lecithin (Soybean)

Propylene Glycol

Tocopheryl Acetate

Carbomer

Menthol

Triethanolamine



ELAIMEI®

Tattoo Numbing Cream

Topical Pain Treatment for Tattoos



Lidocaine 5%
with Vitamin E

6 hours

Maximum Strength Pain Relieves

50ml/ 1.76 fl.oz
For external use only

Drug Facts

Active ingredient

Lidocaine 5%.....Topical Anesthetic

Purpose

Inactive Ingredients:

Aloe barbadensis leaf juice (Aloe vera), Water, Lecithin (Soybean), Propylene Glycol, Tocopheryl Acetate, Carbomer, Menthol, Triethanolamine.

Uses: For temporary relief of local discomfort, itching, pain, soreness, or burning in the perianal area associated with anorectal disorders.

Warnings (For external use only)

Do not use this product if pregnant or breastfeeding, ask a health professional before use.

Drug Facts (Continued)

When using this product

- Do not exceed the recommended daily dosage unless directed by a doctor.
- Do not put into the rectum by using fingers or any medical device or applicator.
- Children under 12 years of age; consult a doctor.

Keep out of reach of children

Keep out of reach of children

In case of accidental ingestion, seek medical attention immediately.

CORRECT USE STEPS BEFORE TATTOOING:

1. Clean hands and treatment area with mild soap and warm water (to open pores to help absorb numbing cream), then gently dry.
2. Apply a thick layer of about 2mm to the treated area. (Make sure the tattoo area and surroundings are white cream)
3. Wrap tightly with plastic wrap, wait 60 minutes (longer time better effect). It is recommended to massage or tap lightly every 10 minutes to help the absorption of the numbing cream.
4. Wipe away residue, wait another 20-30 minutes for the numbness to peak. And begin procedure.

Notice:

Due to the difference of temperature and skin type, the duration of numbness effect will be different.

Recommended to test in a small area first to ensure the best numbness effect. (the longer wrap time, better effect)

Apex CE Limited (RP)

North Point House, North Point Business Park, Co. Cork,

T23 AT2P Ireland

Stop use and ask a doctor if Condition worsens, or does not improve within 7 days. Certain persons can develop allergic reactions to ingredients in this product. If the symptom being treated does not subside or if redness, irritation, swelling, pain, or other symptoms develop or increase, discontinue use.

Directions

- Adults and Children 12 years and older: When practical, cleanse the affected area with mild soap and warm water and rinse thoroughly. Gently dry by patting or blotting with toilet tissue or a soft cloth, before application of this product. Apply externally to the affected area up to 6 times daily.



ELAIMEI TATTOO NUMBING CREAM

lidocaine 5% cream

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:83804-005	
Route of Administration	TOPICAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)		LIDOCAINE	5 g in 100 mL	
Inactive Ingredients				
Ingredient Name			Strength	
WATER (UNII: 059QF0KO0R)				
.ALPHA.-TOCOPHEROL ACETATE (UNII: 9E8X80D2L0)				
ALOE BARBADENSIS LEAF (UNII: ZY81Z83H0X)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
TRIETHANOLAMINE (UNII: 9O3K93S3TK)				
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)				
CARBOMER (UNII: 0A5MM307FC)				
MENTHOL (UNII: L7T10EIP3A)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83804-005-01	1 in 1 BOX	06/16/2025	
1		50 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	06/16/2025	

Labeler - Shenzhen Shandian Jingling Technology Co., Ltd. (418636058)

Establishment

Name	Address	ID/FEI	Business Operations
Shenzhen Shandian Jingling Technology Co., Ltd.		418636058	manufacture(83804-005)

Revised: 6/2025

Shenzhen Shandian Jingling Technology Co., Ltd.