

**IBUPROFEN- ibuprofen tablet, film coated
DOLGENCORP, LLC**

Rite Aid 44-329

Active ingredient (in each brown tablet)

Ibuprofen USP, 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - muscular aches
 - backache
 - toothache
 - menstrual cramps
 - headache
 - the common cold
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- facial swelling
- blisters
- hives
- shock
- skin reddening
- asthma (wheezing)
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- take more or for a longer time than directed
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - slurred speech
 - leg swelling
 - trouble breathing
 - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen at 20

weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 tablet, 2 tablets may be used
 - do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store between 20°-25°C (68°-77°F)
- avoid excessive heat 40°C (104°F)
- see end flap for expiration date and lot number

Inactive ingredients

carnauba wax, colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?

1-888-309-9030

Principal Display Panel

DG™ | health

**Compare to
active ingredient
of Advil® Tablets****

Ibuprofen

Tablets USP, 200 mg

Pain Reliever/
Fever Reducer

(NSAID)

100 + 100 =
200 Coated Tablets

Actual Tablet Size

200
mg each

**TAMPER EVIDENT: DO NOT USE IF
IMPRINTED SAFETY SEAL UNDER
CAP IS BROKEN OR MISSING**

**This product is not manufactured or
distributed by PF Consumer Healthcare 1 LLC,
owner of the registered trademark Advil®
Tablets. 50844 ORG122129112

DISTRIBUTED BY OLD EAST MAIN CO.
100 MISSION RIDGE
GOODLETTSVILLE, TN 37072

Drug Facts (continued)

Warnings
 Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 facial swelling ■ hives ■ shock
 If an allergic reaction occurs, stop use and seek medical help right away.
 Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ take more or for a longer time than directed ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have had stomach ulcers or bleeding problems ■ have 3 or more alcoholic drinks every day while using this product
 Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure,

Uses
 ■ temporarily relieves minor aches and pains due to:
 ■ toothache ■ menstrual cramps ■ headache ■ the common cold ■ minor pain of arthritis ■ muscle aches ■ backache
 ■ temporarily reduces fever

Active ingredient
 Ibuprofen USP, 200 mg (NSAID)
 (in each brown tablet)

Purpose
 ■ nonsteroidal anti-inflammatory drug
 ■ Pain reliever/fever reducer

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
 ■ right before or after heart surgery

Ask a doctor before use if
 ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you are taking a diuretic ■ you have high blood pressure, asthma, or had a stroke ■ you have kidney disease, heart disease, liver cirrhosis, or had a stroke

Ask a doctor or pharmacist before use if you are
 ■ under a doctor's care for heart attack or stroke, because taking aspirin for heart attack or stroke, because pain relievers or fever reducers

When using this product
 ■ take with food or milk if stomach upset occurs ■ taking any other drug ■ ibuprofen may decrease this benefit of aspirin

Stop use and ask a doctor if
 ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ chest pain ■ slurred speech ■ trouble breathing ■ leg swelling ■ weakness in one part or side of body ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear

B-0315-291-12-DB
 ORG122129112



Ibuprofen

Tablets USP, 200 mg
 Pain Reliever/Fever Reducer
 (NSAID)

TWIN PACK • TWIN PACK



Compare to active ingredient of Advil® Tablets**

Ibuprofen

Tablets USP, 200 mg
 Pain Reliever/
 Fever Reducer
 (NSAID)



100 + 100 =

200 mg each

200 Coated Tablets
 Actual Tablet Size



TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts (continued)
 iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?
 1-888-309-9030

**This product is not manufactured or distributed by PF Consumer Healthcare 1 LLC, owner of the registered trademark Advil® Tablets. 50844 ORG122129112

DISTRIBUTED BY OLD EAST MAIN CO.
 100 MISSION RIDGE
 GOODLETTSVILLE, TN 37072


100% Satisfaction Guaranteed!
 If you're not satisfied with this product for any reason, please call us so we can make you 100% satisfied.
 (888) 309-9030

Other information
 ■ store between 20°-25°C (68°-77°F) ■ avoid excessive heat 40°C (104°F) ■ see end flap for expiration date and lot number

Inactive ingredients
 carnauba wax, colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red

Directions
 ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.



Dollar General 44-291

IBUPROFEN

ibuprofen tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55910-791
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	brown	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	44;291
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	NDC 55910			

1	NDC:55910-791-96	2 in 1 CARTON	05/12/2023	
1		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:55910-791-08	1 in 1 CARTON	05/12/2023	
2		24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075010	05/12/2023	

Labeler - DOLGENCORP, LLC (068331990)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(55910-791)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(55910-791)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(55910-791)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(55910-791) , pack(55910-791)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(55910-791)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(55910-791)