

**MUCINEX RAPID CLEAR DAY NIGHT AND CLEAR AND COOL NASAL-  
acetaminophen, guaifenesin, dextromethorphan hydrobromide, triprolidine  
hydrochloride, and oxymetazoline hydrochloride  
RB Health (US) LLC**

-----

**MUCINEX® Rapid Clear® Day Night Caplet + Clear & Cool Nasal Spray Combo**

**DAY**

Acetaminophen 325 mg  
Guaifenesin 200 mg

**NIGHT**

Acetaminophen 325 mg  
Dextromethorphan HBr 10 mg  
Triprolidine HCl 1.25 mg

**SPRAY**

Oxymetazoline hydrochloride 0.05%

**DAY**

Acetaminophen 325 mg ..... Pain reliever/  
fever reducer  
Guaifenesin 200 mg..... Expectorant

**NIGHT**

Acetaminophen 325 mg ..... Pain reliever/fever reducer  
Dextromethorphan HBr 10 mg.....Cough suppressant  
Triprolidine HCl 1.25 mg.....Antihistamine

**SPRAY**

Oxymetazoline hydrochloride 0.05% - Nasal decongestant

**DAY**

**Uses**

■ temporarily relieves these common cold and flu symptoms:

- minor aches and pains ■ headache
- sore throat
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily reduces fever

**NIGHT**

**Uses**

- temporarily relieves these common cold and flu symptoms:
- cough ■ minor aches and pains

- sore throat ■ headache
- runny nose ■ sneezing
- itching of the nose or throat
- itchy, watery eyes due to hay fever
- temporarily reduces fever
- controls cough to help you get to sleep

## **SPRAY**

### **Uses**

temporarily relieves nasal congestion due to:

- a cold
- hay fever or other upper respiratory allergies
- promotes nasal and sinus drainage
- temporarily relieves sinus congestion and pressure
- helps clear nasal passages; shrinks swollen membranes

### **DAY**

#### **Warnings**

Liver warning: This product contains acetaminophen.

Severe liver damage may occur if you take:

- more than 12 caplets in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have

- liver disease
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or

emphysema

■ cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

■ pain or cough gets worse or lasts more than 7 days

■ fever gets worse or lasts more than 3 days

■ redness or swelling is present

■ new symptoms occur

■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## NIGHT

### Warnings

Liver warning: This product contains acetaminophen.

Severe liver damage may occur if you take:

■ more than 8 caplets in 24 hours, which is the maximum daily amount for this product

■ with other drugs containing acetaminophen

■ 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease ■ glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## **SPRAY**

Ask a doctor before use if you have

heart disease  
high blood pressure  
thyroid disease  
diabetes  
difficulty in urination due to enlargement of the prostate gland

When using this product

do not exceed recommended dosage  
do not use this product for more than 3 days. Use only as directed. Frequent or prolonged use may cause nasal congestion to recur or worsen  
this product may cause temporary discomfort such as burning, stinging, sneezing, or an increase in nasal discharge  
the use of this container by more than one person may spread infection

Stop use and ask a doctor if symptoms persist

If pregnant or breast-feeding, ask a health professional before use.

## **DAY**

### **Directions**

- do not take more than directed (see Overdose warning)
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

## **NIGHT**

### **Directions**

- do not take more than directed (see Overdose warning)
- do not take more than 8 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

## **SPRAY**

adults and children 6 to under 12 years of age (with adult supervision): 2 or 3 sprays in each nostril, not more often than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period.

children under 6 years of age: consult a doctor.

Shake well before use.

To open, hold by the white grips then squeeze, push down firmly and turn cap counterclockwise. Before using for the first time, remove the protective cap from the tip and prime metered pump by depressing firmly several times. To spray, hold bottle with thumb at the base and nozzle between first and second fingers. Without tilting head, insert nozzle into nostril. Fully depress pump all the way down with a firm

even stroke and sniff deeply. Wipe nozzle clean after use.

To close, turn cap clockwise.

DO NOT DISCARD CAP.

## **DAY/NIGHT/SPRAY**

### **Other information**

■ store at 20-25°C (68-77°F)

### **DAY**

#### **Inactive ingredients**

croscarmellose sodium, FD&C red no. 40  
aluminum lake, FD&C yellow no. 6 aluminum  
lake, magnesium stearate, microcrystalline  
cellulose, polyethylene glycol, polyvinyl  
alcohol, povidone, talc, titanium dioxide

### **NIGHT**

#### **Inactive ingredients**

croscarmellose sodium, crospovidone, hypromellose,  
microcrystalline cellulose, polyethylene glycol,  
polysorbate 80, povidone, titanium dioxide

### **SPRAY**

benzalkonium chloride,  
benzododecinium chloride,  
camphor

Questions? 1-866-MUCINEX  
(1-866-682-4639)

Keep out of reach of children.

If pregnant or breast-feeding,ask a health professional before use.

# Mucinex®

POWERFUL MULTI-SYMPTOM RELIEF

COLD + SINUS VALUE PACK

NDC 72854-213-03

# Mucinex®

**RAPID+CLEAR™**

**DAY TIME**  
**PAIN, HEADACHE &  
MUCUS CONGESTION**

Acetaminophen »  
Pain Reliever/Fever Reducer  
Guaifenesin » Expectorant

**NIGHT TIME**  
**HEADACHE,  
COUGH & FEVER**

Acetaminophen » Pain Reliever/Fever Reducer  
Dextromethorphan HBr » Cough Suppressant  
Triprolidine HCl » Antihistamine

AGES 12+


AGES 12+

- ✓ Pain & Headache
- ✓ Fever
- ✓ Thins & Loosens Mucus

- ✓ Pain & Headache
- ✓ Fever + Itchy Throat
- ✓ Cough



**RAPID RELEASE**



**12** CAPLETS

TOTAL 20 CAPLETS

**8** CAPLETS

**SINUS MAX®**

**SEVERE NASAL  
CONGESTION RELIEF**  
**CLEAR&COOL**

NASAL SPRAY  
Oxymetazoline HCl 0.05%  
Nasal Decongestant



- ✓ Fast Acting
- ✓ Lasts 12 Hours
- ✓ Cooling Menthol Flavor

**3/4 FL OZ (22 mL)**

**Temper evident:**  
Do not use if carton is  
damaged or if printed seal  
on blister is broken or missing.

**Mucinex**  
**RAPID+CLEAR™**

**DAY TIME**  
**PAIN, HEADACHE &  
MUCUS CONGESTION**  
**NIGHT TIME**  
**HEADACHE, COUGH & FEVER**

Take only as directed.  
Keep carton for full  
information.

**PARENTS:**  
Learn about teen medicine abuse

[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)



reckitt

Dist. by: RB Health (US)  
Parsippany, NJ 07054-0224  
©2025 RB Health  
Patents: [www.reckitt.com/patents](http://www.reckitt.com/patents)  
020525 3325213

**Temper evident:**  
Do not use if printed seal  
over cap is torn or missing.

**Mucinex**  
**SINUS MAX®**

**SEVERE NASAL  
CONGESTION RELIEF**  
**CLEAR&COOL**

To open: hold by the white  
grips then squeeze, push  
down firmly & turn cap.



To close:  
turn cap clockwise

Keep carton for full  
information.



3



Do not take the Mucinex® RAPID+CLEAR™ Daytime Pain, Headache & Mucus Congestion and Mucinex® RAPID+CLEAR™ Nighttime Headache, Cough & Fever caplets at the same time. Always wait at least 4 hours before taking another dose of Mucinex® caplets. Keep carton for complete product information. Do not take more than directed (see Overdose warning).

### Daytime Pain, Headache & Mucus Congestion

### Nighttime Headache, Cough & Fever

### Severe Nasal Congestion Relief Clear & Cool Nasal Spray

#### Drug Facts

##### Active ingredients (in each caplet) Purposes

Acetaminophen 325 mg..... Pain reliever/fever reducer  
Guaifenesin 200 mg..... Expectorant

#### Uses

- temporarily relieves these common cold and flu symptoms:
  - minor aches and pains
  - headache
  - sore throat
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily reduces fever

#### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 caplets in 24 hours, which is the maximum daily amount for this product

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if you have**

- liver disease
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

**When using this product do not use more than directed**

#### Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

#### Drug Facts

##### Active ingredients (in each caplet) Purposes

Acetaminophen 325 mg..... Pain reliever/fever reducer  
Dextromethorphan HBr 10 mg..... Cough suppressant  
Triprolidine HCl 1.25 mg..... Antihistamine

#### Uses

- temporarily relieves these common cold and flu symptoms:
  - cough
  - minor aches and pains
  - headache
  - runny nose
  - sneezing
  - itching of the nose or throat
  - itchy, watery eyes due to hay fever
- temporarily reduces fever
- controls cough to help you get to sleep

#### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 8 caplets in 24 hours, which is the maximum daily amount for this product

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

#### Ask a doctor before use if you have

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

#### Drug Facts

##### Active ingredient Purpose

Oxymetazoline hydrochloride 0.05%... Nasal decongestant

#### Uses

- temporarily relieves nasal congestion due to:
  - a cold
  - hay fever or other upper respiratory allergies
- promotes nasal and/or sinus drainage; temporarily relieves sinus congestion and pressure
- helps clear nasal passages; shrinks swollen membranes

#### Warnings

**Ask a doctor before use if you have**

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- difficulty in urination due to enlargement of the prostate gland

#### When using this product

- do not exceed recommended dosage
- do not use this product for more than 3 days. Use only as directed. Frequent or prolonged use may cause nasal congestion to recur or worsen.
- this product may cause temporary discomfort such as burning, stinging, sneezing, or an increase in nasal discharge
- the use of this container by more than one person may spread infection

**Stop use and ask a doctor if symptoms persist**

**If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

#### Directions

- adults and children 6 to under 12 years of age (with adult supervision): 2 or 3 sprays in each nostril, not more often than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period.
- children under 6 years of age: consult a doctor.

**Shake well before use.**

To open, hold by the white grips then squeeze, push down firmly and turn cap counterclockwise. Before using for the first time, remove the protective cap from the tip and prime metered pump by depressing firmly several times. To spray, hold bottle with thumb at the base and nozzle between first and second fingers. Without tilting head, insert nozzle into nostril. Fully depress pump all the way down with a firm even stroke and sniff deeply. Wipe nozzle clean after use. To close, turn cap clockwise. **DO NOT DISCARD CAP.**

**PEEL CORNER TO READ COMPLETE DRUG FACTS AND INFORMATION**



**Daytime Pain, Headache & Mucus Congestion (cont'd)**

**Drug Facts (continued)**

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children. Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not take more than directed (see Overdose warning)
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

**Other information**

- store at 20-25°C (68-77°F)

**Inactive ingredients**

croscarmellose sodium, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, talc, titanium dioxide

**Questions? 1-866-MUCINEX (1-866-682-4639)**

**Nighttime Headache, Cough & Fever (cont'd)**

**Drug Facts (continued)**

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

**When using this product**

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- use caution when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with a rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children. Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not take more than directed (see Overdose warning)
- do not take more than 8 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

**Other information** ■ store at 20-25°C (68-77°F)

**Inactive ingredients**

croscarmellose sodium, crospovidone, hypromellose, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, titanium dioxide

**Questions? 1-866-MUCINEX (1-866-682-4639)**

**Severe Nasal Congestion Relief Clear & Cool Nasal Spray (cont'd)**

**Drug Facts (continued)**

**Other information**

- store at 20-25°C (68-77°F)

**Inactive ingredients**

benzalkonium chloride, benzododecinium chloride, camphor, cetalkonium chloride, colloidal silicon dioxide, edetate disodium, eucalyptol, glycine, linoleic acid, linolenic acid, menthol, myristalkonium chloride, myristic acid, oleic acid, palmitic acid, palmitoleic acid, polyethylene glycol, polysorbate 80, propylene glycol, purified water, sodium carbonate, sodium chloride, sodium hydroxide, stearic acid

**Questions? 1-866-MUCINEX (1-866-682-4639)**



COATING FREE AREA



COATING FREE AREA



**MUCINEX RAPID CLEAR DAY NIGHT AND CLEAR AND COOL NASAL**

acetaminophen, guaifenesin, dextromethorphan hydrobromide, triprolidine hydrochloride, and oxymetazoline hydrochloride kit

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:72854-213
---------------------	----------------	---------------------------	---------------

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72854-213-03	1 in 1 CARTON	06/01/2025	
1		1 in 1 KIT; Type 1: Convenience Kit of Co-Package		

### Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BLISTER PACK	12
Part 2	1 BLISTER PACK	8
Part 3	0 BOTTLE, PUMP	1 mL

### Part 1 of 3

## MAXIMUM STRENGTH MUCINEX SINUS-MAX PRESSURE, PAIN AND COUGH

acetaminophen, guaifenesin tablet, film coated

### Product Information

Item Code (Source)	NDC:72854-208
Route of Administration	ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

### Inactive Ingredients

Ingredient Name	Strength
FD&C RED NO. 40 ALUMINUM LAKE (UNII: 6T47AS764T)	
FD&C YELLOW NO. 6 ALUMINUM LAKE (UNII: GYP6Z2JR6Q)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
POLYVINYL ALCOHOL (UNII: 532B59J990)	
POVIDONE (UNII: FZ989GH94E)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
TALC (UNII: 7SEV7J4R1U)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	

### Product Characteristics

Color	red	Score	no score
Shape	OVAL	Size	20mm
Flavor		Imprint Code	VV;MSC
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/01/2025	

### Part 2 of 3

### MUCINEX NIGHTSHIFT SINUS MAXIMUM STRENGTH

acetaminophen, dextromethorphan hydrobromide, and triprolidine hydrochloride tablet, coated

### Product Information

Item Code (Source)	NDC:72854-209
Route of Administration	ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>TRIPROLIDINE HYDROCHLORIDE</b> (UNII: YAN7R5L890) (TRIPROLIDINE - UNII:2L8T9S52QM)	TRIPROLIDINE HYDROCHLORIDE	1.25 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>CROSPVIDONE, UNSPECIFIED</b> (UNII: 2S7830E561)	
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	

### Product Characteristics

Color	yellow	Score	no score
-------	--------	-------	----------

Shape	OVAL	Size	19mm
Flavor		Imprint Code	VW;LOGOcrescentmoonplus
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/01/2025	

### Part 3 of 3

## MUCINEX SINUS-MAX SEVERE NASAL CONGESTION RELIEF CLEAR AND COOL NASAL

oxymetazoline hydrochloride solution

### Product Information

Item Code (Source)	NDC:63824-129
Route of Administration	NASAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>OXYMETAZOLINE HYDROCHLORIDE</b> (UNII: K89MJ0S5VY) (OXYMETAZOLINE - UNII:8VLN5B44ZY)	OXYMETAZOLINE HYDROCHLORIDE	0.05 g in 100 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>CAMPHOR (NATURAL)</b> (UNII: N20HL7Q941)	
<b>BENZALKONIUM CHLORIDE</b> (UNII: F5UM2KM3W7)	
<b>BENZODODECINIUM CHLORIDE</b> (UNII: Y5A751G47H)	

### Product Characteristics

Color		Score	
Shape		Size	
Flavor	MENTHOL	Imprint Code	

**Contains****Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		22 mL in 1 BOTTLE, PUMP; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	10/01/2018	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/01/2025	

**Labeler** - RB Health (US) LLC (081049410)

Revised: 6/2025

RB Health (US) LLC