

**MUCINEX RAPID CLEAR DAYTIME PAIN, HEADACHE, AND COUGH-
acetaminophen, dextromethorphan hydrobromide capsule, liquid filled
RB Health (US) LLC**

MUCINEX® Rapid Clear® Liquid Gels Pain, Headache & Cough

Drug Facts

Active ingredients (in each liquid gel) Purposes
Acetaminophen 325 mg Pain reliever/fever reducer
Dextromethorphan HBr 10 mg Cough suppressant

<i>Active ingredients (in each liquid gel)</i>	<i>Purposes</i>
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant

Uses

temporarily relieves these common cold and flu symptoms:

- cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
- the intensity of coughing
- the impulse to cough to help you get to sleep
- minor aches and pains
- sore throat
- headache
- temporarily reduces fever

Warnings

This product contains acetaminophen. Severe liver damage may occur if you take:
more than 12 liquid gels in 24 hours, which is the maximum daily amount
with other drugs containing acetaminophen
3 or more alcoholic drinks daily while using this product

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 liquid gels in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening

- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick

medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 12 liquid gels in any 24-hour period
- adults and children 12 years of age and over: take 2 liquid gels every 4 hours
- children under 12 years of age: do not use

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat

Inactive ingredients

FD&C red no. 40, FD&C yellow no. 6, gelatin, glycerin, hypromellose, isopropyl alcohol, lecithin (sunflower), light mineral oil, mannitol, polyethylene glycol 400, povidone, propylene glycol, purified water, 1,4-sorbitan, sorbitol, titanium dioxide

Questions?

1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.

Dist. by: RB Health (US)
Parsippany, NJ 07054-0224

PRINCIPAL DISPLAY PANEL - 16 Capsule Blister Pack Carton

Fast Dissolving Liquid Gels!

NDC 72854-214

MAXIMUM STRENGTH

MUCINEX®

Acetaminophen – Pain Reliever/Fever Reducer

Dextromethorphan HBr – Cough Suppressant

LIQUID GELS

(Liquid Filled Capsules)

Actual Size

MAXIMUM STRENGTH

Drug Facts

Active ingredients (in each capsule)

Acetaminophen 325 mg.....Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....Cough suppressant

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - the impulse to cough to help you get to sleep
 - minor aches and pains
 - sore throat
 - headache
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 capsules in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

if with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Purposes

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product do not use more than directed

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 12 capsules in any 24-hour period
- adults and children 12 years of age and over: take 2 capsules every 4 hours
- children under 12 years of age: do not use

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat

Inactive ingredients

FD&C red no. 40, FD&C yellow no. 6, gelatin, glycerin, hydroxytoluene, isopropyl alcohol, lecithin (sunflower), light mineral oil, menthol, polyethylene glycol 400, povidone, propylene glycol, purified water, 1,4-sorbitan, sorbitol, titanium dioxide

Questions? 1-866-MUCINEX (1-866-682-4639)

reckitt

Patents:

www.reckitt.com/patents

Dist. by: RB Health (US)

Parlappan, NJ 07054-4224

12824

3321583

©2025 RB Health

PAIN, HEADACHE & COUGH

Mucinex
RAPID+CLEAR

MAXIMUM STRENGTH

Maximum Strength per 4-hour dose
 Do not take more than a total of 12 capsules in any 24-hour period.
 Take only as directed.
 Keep carton for full information.

Scan for FAQs and instructions on proper disposal of medicines or visit www.mucinex.com/fdas
 Please visit our website www.mucinex.com

RAPID RELEASE LIQUID GELS

MAXIMUM STRENGTH

NDC 72854-214-16

Mucinex

RAPID+CLEAR

PAIN, HEADACHE & COUGH

Acetaminophen » Pain Reliever/Fever Reducer
 Dextromethorphan HBr » Cough Suppressant

- ✓ Relieves Pain & Headache
- ✓ Reduces Fever
- ✓ Controls Cough

16

LIQUID GELS (Liquid Filled Capsules)

RAPID RELEASE

Tamper evident:
 Do not use if carton is damaged or if printed seal on blister is broken or missing.

PARENTS:
 Learn about your medicine. Ask your pharmacist.
www.StopMedicineAbuse.org

how2recycle.info

3 63224 9945 6

AGES 12+

acetaminophen, dextromethorphan hydrobromide capsule, liquid filled

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72854-214
Route of Administration	ORAL		

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg

Ingredient Name	Strength
SORBITOL (UNII: 506T60A25R)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C00X)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
ISOPROPYL ALCOHOL (UNII: ND2M416302)	

LIGHT MINERAL OIL (UNII: N6K5787QVP)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
WATER (UNII: 059QF0KO0R)				
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
1,4-SORBITAN (UNII: AV0YTZ4E6J)				
LECITHIN, SUNFLOWER (UNII: 834K0WOS5G)				
MANNITOL (UNII: 3OWL53L36A)				

Product Characteristics			
Color	red	Score	no score
Shape	OVAL	Size	24mm
Flavor		Imprint Code	AR18
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72854-214-16	1 in 1 CARTON	06/01/2025	
1		16 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/01/2025	

Labeler - RB Health (US) LLC (081049410)