

DIUREX MAX- pamabrom tablet, film coated
Kobayashi Healthcare International, Inc.

Diurex Max

Active ingredient (in each caplet)

Pamabrom, 50 mg

Purpose

Diuretic

Uses

For the relief of

- temporary water retention
- bloating
- swelling
- full feeling

associated with the premenstrual and menstrual periods.

Warnings

Ask a doctor or pharmacist before use if you are taking any other medications.

Stop use and ask a doctor if symptoms last more than ten consecutive days.

If pregnant or breastfeeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a poison control center right away.

Directions

- Read all package directions and warnings before use.
- Use only as directed.
- Adults: One (1) pill after breakfast with a full glass of water.
- Dose may be repeated after 6 hours, not to exceed four (4) pills in 24 hours.
- Start taking 5 or 6 days before onset of period and continue until desired relief is obtained or end of period.
- Drink 6 to 8 glasses of water daily.
- For use by normally healthy adults only.
- Persons under 18 years of age should use only as directed by a doctor.

Other information

Calcium content: 160 mg/pill. ****Contents sealed:** Each DIUREX Max, light blue colored, oval shaped pill bears the identifying mark "ALVA" plus a "1" on the pill's reverse side and is sealed in a clear plastic blister with a foil backing. Do not use if seal appears broken or if product contents do not match product description. NOTE: The appearance of golden tinted urine which may occur after taking Diurex Water Pills is a normal and temporary effect. You may report serious side effects to the phone number provided under *Questions?* below.

Inactive ingredients

Calcium sulfate dihydrate, croscarmellose sodium, dicalcium phosphate, FDC Blue No. 1, hypromellose, magnesium stearate, mineral oil, polyethylene glycol, potassium gluconate, riboflavin, sodium lauryl sulfate, stearic acid, talc and titanium dioxide.

Questions? 1-800-792-2582

Datos Farmacológicos

Ingrediente activo (en cada comprimido)	Propósito
Pamabrom, 50 mg.....	Diurético

Usos Para el alivio de la retención temporal de líquidos
■ distensión abdominal ■ hinchazón ■ sensación de saciedad asociadas con los períodos menstruales y premenstruales.

Advertencias

Consulte con su médico o farmacéutico antes de usarlo si está tomando cualquier otro medicamento.

Interrumpa el uso y consulte con su médico si los síntomas duran más de diez días consecutivos.

Si está embarazada o está amamantando, consulte con un profesional de la salud antes de usarlo. ■ **Mantener fuera del alcance de los niños.** En caso de sobredosis, obtenga ayuda médica o comuníquese de inmediato con un centro de toxicología.

Indicaciones ■ Lea todas las indicaciones y advertencias del envase antes de usarlo. ■ Úselo solo de acuerdo con las indicaciones. ■ **Adultos:** Un (1) comprimido después del desayuno con un vaso lleno de agua. ■ La dosis puede repetirse después de 6 horas, sin exceder los cuatro (4) comprimidos en 24 horas. ■ Comience a tomarlo 5 o 6 días antes del inicio del período y

continúe hasta que se obtenga el alivio deseado o hasta el final del período. ■ Tome de 6 a 8 vasos de agua todos los días. ■ Solo para uso en adultos normalmente sanos. ■ Las personas menores de 18 años deben usarlo solo según las indicaciones de un médico. ▼

Datos Farmacológicos (continuación)

Información adicional Contenido de calcio: 160 mg/comprimido.****El contenido está sellado:** Cada comprimido DIUREX, de color azul claro y forma ovalada lleva la marca identificatoria "ALVA" y un "1" en el dorso, y está sellado en un blíster de plástico transparente con un recubrimiento posterior de papel aluminio. No lo use si el sello parece estar roto o si el producto no coincide con la descripción del producto. NOTA: La aparición de orina de color dorado que puede ocurrir después de tomar los comprimidos contra la retención de líquido DIUREX es un efecto normal y temporal. ■ Puede informar efectos secundarios graves al número de teléfono que aparece en la sección *¿Tiene preguntas?* a continuación.

Ingredientes inactivos Sulfato de calcio dihidratado, croscarmelosa sódica, fosfato dicálcico, FD&C azul n.º 1 Lake, hipromelosa, estearato de magnesio, aceite mineral, polietilenglicol, gluconato de potasio, riboflavina, lauril sulfato sódico, ácido esteárico, talco y dióxido de titanio.

¿Tiene preguntas? 1-800-792-2582



diurex[®]

CAFFEINE FREE

Bloat Reliever

Prompt & Effective
Water Bloat Relief

Relieve Discomfort

24

CAPLETS

DIURETIC

DOCTOR RECOMMENDED

CMYK
PMS 286C

maximum strength

diurex[®]

CAFFEINE FREE



Bloat Reliever
Alivio de la distensión abdominal

Vea las instrucciones en español adentro



máxima concentración
diurex[®]
SIN CAFEÍNA

49/24A-14RC,
0824

Drug Facts

Active ingredient (in each caplet) Furosemide, 50 mg. Diuretic

Uses For the relief of ■ temporary water retention ■ bloating ■ swelling ■ full feeling associated with the premenstrual and menstrual periods.

Warnings
Ask a doctor or pharmacist before use if you are taking any other medications.
Stop use and ask a doctor if symptoms last for more than ten consecutive days.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a poison control center right away.

Directions ■ Read all package directions and warnings before use. ■ Use only as directed. ■ Adults: One (1) caplet after breakfast with a full glass of water. ■ Dose may be repeated after 6 hours, not to exceed four (4) caplets in 24 hours. ■ Start taking 5 or 6 days before onset of period and continue until desired relief is obtained or end of period. ■ Drink 6 to 8 glasses of water daily. ■ For use by normally healthy adults only. ■ Persons under 18 years of age should use only as directed by a doctor.

Other information Calcium content: 160 mg/caplet. ■ *Contents sealed: Each Diurex, light blue colored, oval shaped caplet bears the identifying mark "ALVA" plus a "1" on the caplet's reverse side and is sealed in a clear plastic blister with a foil backing. Do not use if seal appears broken or if product contents do not match product description. NOTE: The appearance of golden tinted urine which may occur after taking Diurex Caffeine Free is a normal and temporary effect. ■ You may report serious side effects to the phone number provided under *Questions?* below.

Inactive ingredients Calcium sulfate dihydrate, croscarmellose sodium, dicalcium phosphate, FD&C Blue #1 Lake, hypromellose, magnesium stearate, mineral oil, polyethylene glycol, potassium gluconate, riboflavin, sodium lauryl sulfate, stearic acid, talc and titanium dioxide.

Questions? 1-800-792-2582
www.diurex.com

CONTAINS NO GLUTEN INGREDIENTS

*Based on independent syndicated market research data.

The active ingredient in Diurex[®] Caffeine Free has been recognized to be safe and effective by an FDA expert advisory panel when used as directed.

LOT:
EXP.:



DKT:
KOBAYASHI
© Reg. U.S. Patent and Trademark Office
PROTECT PRODUCT FROM EXCESSIVE HEAT AND MOISTURE.



Prod. No. 49524
**Contents Sealed

KOBAYASHI Consumer Products, LLC
P.O. Box 1191, Dalton, GA 30722, U.S.A.
© 1973-2024



DIUREX MAX

pamabrom tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:52389-249
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PAMABROM (UNII: UA8U0KJM72) (BROMOTHEOPHYLLINE - UNII:FZG87K1MQ6)	PAMABROM	50 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
LIGHT MINERAL OIL (UNII: N6K5787QVP)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POTASSIUM GLUCONATE (UNII: 12H3K5QKN9)	
CALCIUM SULFATE DIHYDRATE (UNII: 4846Q921YM)	
RIBOFLAVIN (UNII: TLM2976OFR)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TALC (UNII: 7SEV7J4R1U)	

Product Characteristics

Color	blue	Score	no score
Shape	OVAL	Size	15mm
Flavor		Imprint Code	ALVA;1
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52389-249-20	1 in 1 CARTON	07/18/2006	04/18/2011
1		30 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:52389-249-40	1 in 1 CARTON	10/01/2015	03/31/2019
2		40 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:52389-249-24	1 in 1 CARTON	04/18/2011	
3		24 in 1 BLISTER PACK; Type 0: Not a Combination Product		
4	NDC:52389-249-48	2 in 1 CARTON	07/18/2013	
4		24 in 1 BLISTER PACK; Type 0: Not a Combination Product		
5	NDC:52389-249-16	2 in 1 CARTON	04/18/2011	05/31/2019
5		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		
6	NDC:52389-249-10	4 in 1 CARTON	11/10/2017	
6		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
7	NDC:52389-249-72	3 in 1 CARTON	06/12/2018	
7		24 in 1 BLISTER PACK; Type 0: Not a Combination Product		
8	NDC:52389-249-30	3 in 1 CARTON	07/12/2021	
8		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
9	NDC:52389-249-50	5 in 1 CARTON	07/18/2022	
9		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M027	12/20/1995	

Labeler - Kobayashi Healthcare International, Inc. (156391729)

Registrant - Kobayashi America Manufacturing, LLC (079852150)