

LOW DOSE ASPIRIUN- aspirin tablet, chewable
AMERIFOODS TRADING COMPANY

1105B-FIR-2025-1120

Drug Facts

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)*

* nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

- for the temporary relief of minor aches and pains or as recommended by your doctor
- ask your doctor about other uses for this product

Warnings

Reye's syndrome

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for:

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

Other information

- store between 20°-25°C (68°-77°F) in a dry place
- retain carton for complete product information and warnings

Inactive ingredients

corn starch, FD&C yellow #6 aluminum lake, flavor, magnesium stearate, silicon dioxide, sucrose

Questions or comments?

1-844-705-4384

PRINCIPAL DISPLAY PANEL

3 PACK 36+36+36=108 TABLETS

FIRST STREET®

NDC 61504-105-03

†COMPARE TO THE ACTIVE INGREDIENT IN BAYER® CHEWABLE ASPIRIN
CHEWABLE

Low Dose Aspirin 81 mg each

Pain Reliever (NSAID)

Aspirin Regimen**

ORANGE FLAVOR

Talk to your doctor or other healthcare provider before using this product for your heart

ACTUAL SIZE

FOR ADULTS

108 TABLETS

NC

Drug Facts

Active ingredient (in each tablet)
Aspirin 81 mg (NSAID)
* nonsteroidal anti-inflammatory drug

Purpose
Pain reliever

Uses
■ for the temporary relief of minor aches and pains or as recommended by your doctor
■ ask your doctor about other uses for this product

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction which may include:
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older
■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed

Do not use
■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
■ for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

Ask a doctor before use if
■ stomach bleeding warning applies to you
■ you have a history of stomach problems, such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
■ you are taking a diuretic
■ you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for:
■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if
■ an allergic reaction occurs. Seek medical help right away.
■ you experience any of the following signs of stomach bleeding:
■ feel faint ■ vomit blood ■ have bloody or black stools
■ have stomach pain that does not get better
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present
■ new symptoms occur
■ ringing in the ears or a loss of hearing occurs
These could be signs of a serious condition.

NC

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
■ drink a full glass of water with each dose
■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
■ children under 12 years: consult a doctor

Other information
■ store between 20°-25°C (68°-77°F) in a dry place
■ retain carton for complete product information and warnings

Inactive ingredients corn starch, FD&C yellow #6 aluminum lake, flavor, magnesium stearate, silicone dioxide, sucrose

Questions or comments?
1-844-705-4384

**Aspirin is not appropriate for everyone, so be sure to talk to your doctor before you begin an aspirin regimen.

DISTRIBUTED BY: AMERIFOODS TRADING CO.
P.O. BOX 512372, LOS ANGELES, CA 90051-0372



CHEWABLE

FIRST STREET

Low Dose Aspirin

3 PACK 36 + 36 + 36 = **108 TABLETS**

FIRST STREET

CHEWABLE

Low Dose Aspirin

Pain Reliever (NSAID)

Aspirin Regimen**

81 mg each

ORANGE FLAVOR

ACTUAL SIZE

108 TABLETS

CHEWABLE

Low Dose Aspirin

Pain Reliever (NSAID)

Aspirin Regimen**

ORANGE FLAVOR

ACTUAL SIZE

108 TABLETS

FIRST STREET

*This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Bayer® Chewable Aspirin.

DO NOT USE IF IMPRINTED SEAL UNDER CAP IS BROKEN OR MISSING

F1105B03F1R_R0




0 41512 18878 7

LOW DOSE ASPIRIUN

aspirin tablet, chewable

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:61504-105
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	

MAGNESIUM STEARATE (UNII: 70097M6I30)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
SUCROSE (UNII: C151H8M554)				
Product Characteristics				
Color	orange	Score	no score	
Shape	ROUND	Size	8mm	
Flavor	ORANGE	Imprint Code	AP;009	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:61504-105-03	3 in 1 CARTON	11/20/2025	
1		36 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M013	11/20/2025	

Labeler - AMERIFOODS TRADING COMPANY (626388680)