

**CHILDRENS ACETAMINOPHEN- childrens acetaminophen oral suspension  
AARNA USA INC.**

-----  
**ACETAMINOPHEN ORAL SUSPENSION GRAPE  
AARNA USA INC.**

**Active Ingredient**

<b>Active ingredient</b> (in each 5mL)	<b>Purpose</b>
Acetaminophen 160 mg .....	Pain reliever/fever reducer

**Uses**

temporarily: reduces fever

relieves minor aches and pains due to:

- the common cold
- flu
- headache
- sore throat
- toothache

**Warnings**

**Liver Warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:**

if sore throat is severe, persists or more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription).
- If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product.

## **Ask a doctor or pharmacist before use if the child has**

liver disease.

Ask doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin. **When using this product do not exceed recommended dose (see overdose warning)**

### **Stop use and ask a doctor if**

- pain gets worse or lasts for more than 5 days
- fever gets worse or last for more than 3 days
- new symptoms occur redness or swelling is present. These could be signs of a serious condition.

Overdose warning:

In case of overdose, get medical help to contact Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

## **Keep out of Reach of children**

### **Directions**

- This product does not contain directions or complete warnings

for adult use

- do not give more than directed (see overdose warning)
- shake well before using
- mL=milliliter
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age.
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

---

<b>Weight (lbs.)</b>	<b>Age (yrs.)</b>	<b>Dose (tsp or mL)</b>
under 24	under 2	ask a doctor
24 to 35	2 to 3	1 tsp or 5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL
48 to 59	6 to 8	2 tsp or 10 mL
60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
72 to 95	11	3 tsp or 15 mL

---

\*or as directed by a doctor

- Attention: use only enclosed dosing cup designed for use with this product. Do not use any other dosing device.

## **Other Information**

- Store at room temperature 20°-25°C (68°-77°F). Protect from freezing. Protect from light.

## Inactive Ingredients

Anhydrous Citric Acid, D&C Red #33, FD&C Blue #1, Grape Flavor, Glycerin, Microcrystalline cellulose and Sodium Methyl Cellulose, Potassium Sorbate, Purified Water, Sorbitol, Sucralose, Sucrose, Xanthan gum.

## Questions or comments?

Call toll-free 1-877-225-6999

## Manufactured for:

AARNA USA INC.

Leland, NC-28451

NDC 82568-0010-4



# Children's Acetaminophen Oral suspension

**160mg per 5mL**  
**PAIN RELIEVER**  
and fever Reducer

For Ages 2 to 11 Years

Alcohol Free  
Ibuprofen Free  
Aspirin Free  
No Parabens



**Grape  
Flavor**

4 FL.OZ (118 mL)

**Drug Facts**

**Active ingredient (in each 5 mL)**  
Acetaminophen 160 mg ..... Pain reliever / fever reducer

**Uses** temporarily relieves fever; relieves minor aches and pains due to: the common cold; flu

**Warnings**  
headache; sore throat; toothache

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:  
more than 5 doses in 24 hours, which is the maximum daily amount; with other drugs containing acetaminophen. **Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening; blisters; rash. If a skin reaction occurs, stop use and seek medical help right away.  
**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. **Do not use** with any other drug containing acetaminophen (prescription or non prescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If your child is allergic to acetaminophen or any of the inactive ingredients in this product.  
**Ask a doctor before use if your child has liver disease.**

**Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.**

**When using this product, do not exceed recommended dose (see overdose warning)**

**Stop use and ask a doctor if:** pain gets worse or lasts for more than 5 days; fever gets worse or lasts for more than 3 days; new symptoms occur; redness or swelling is present. These could be signs of a serious condition. **Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

**Directions** This product does not contain directions or complete warnings for adult use  
do not give more than directed ( see overdose warning) shake well before using  
mL = milliliter find right dose on chart below. If possible, use weight to dose; otherwise, use age.  
repeat dose every 4 hours while symptoms last. do not give more than 5 times in 24 hours

Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)	Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)
under 24	under 2	ask a doctor	48 to 59	6 to 8	2 tsp or 10 mL
24 to 35	2 to 3	1 tsp or 5 mL	60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL	72 to 95	11	3 tsp or 15 mL

\* or as directed by a doctor  
**Attention:** use only enclosed dosing cup designed for use with this product.  
Do not use any other dosing device.

**Other information** store at room temperature 20°-25°C (68°-77°F). Protect from freezing. Protect from Light.

**Inactive ingredients** anhydrous citric acid, D&C red #33, FD&C blue #1, grape flavor, glycerin, microcrystalline cellulose, and sodium carboxymethylcellulose, potassium sorbate, purified water, sorbitol solution, sucralose, sucrose, xanthan gum

**Questions or comments?** call toll-free 1-877-225-6999.

\*\*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Infant's Tylenol Oral Suspension.

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING**



**Manufactured for:**  
AARNA USA Inc.  
Leland, NC-28451



N 82568-00104-1  
3

Exclusively distributed by Aarna Pharma Inc. Fairfield, NJ-07004

# Acetaminophen Oral suspension

## Drug Facts (continued)

- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)*
under 24	under 2	ask a doctor
24 to 35	2 to 3	1 tsp or 5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL
48 to 59	6 to 8	2 tsp or 10 mL
60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
72 to 95	11	3 tsp or 15 mL

\*or as directed by a doctor

**Attention:** use only enclosed dosing cup designed for use with this product. Do not use any other dosing device.

### Other Information

store at room temperature 20-25°C (68-77°F). Protect from Freezing. Protect from Light.

**Inactive ingredients** anhydrous citric acid, D&C red #33, FD&C blue #1, grape flavor, glycerin, microcrystalline cellulose and sodium carboxymethylcellulose, potassium sorbate, purified water, sorbitol solution, sucralose, sucrose, xanthan gum

### Questions or comments?

call toll-free 1-877-225-6999.

\*\*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Infants' Tylenol® Oral Suspension.

**TAMPER EVIDENT: DO NOT USE IF SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

**Aarna USA**  
Manufactured for:  
AARNA USA Inc.  
Leland, NC-28451  
Exclusively distributed by Akron Pharma Inc. Fairfield, NJ-07004



NDC 82568-0010-4



# Children's Acetaminophen Oral suspension

160mg per 5mL

**PAIN RELIEVER**  
and fever Reducer  
For Ages 2 to 11 Years

Alcohol Free  
Ibuprofen Free  
Aspirin Free  
No Parabens



**Grape Flavor**

4 FL.OZ (118 mL)

NDC 82568-0010-4



# Children's Acetaminophen Oral suspension

160mg per 5mL

**PAIN RELIEVER**  
and fever Reducer  
For Ages 2 to 11 Years

Alcohol Free  
Ibuprofen Free  
Aspirin Free  
No Parabens



**Grape Flavor**

4 FL.OZ (118 mL)

## Drug Facts

**Active ingredient (in each 5 mL)** Purpose

Acetaminophen 160 mg Pain reliever / fever reducer

**Uses temporarily:** reduces fever ■ relieves minor aches and pains due to: ■ the common cold ■ flu ■ headache ■ sore throat ■ toothache

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:  
■ more than 5 doses in 24 hours, which is the maximum daily amount  
■ with other drugs containing acetaminophen.

### Allergy alert:

Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** ■ with any other drug containing acetaminophen (prescription or non prescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ If your child is allergic to acetaminophen or any of the inactive ingredients in this product.

**Ask a doctor before use if your child has liver disease.**

**Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.**

**When using this product, do not exceed recommended dose (see overdose warning)**

**Stop use and ask a doctor if** ■ pain gets worse or lasts for more than 5 days ■ fever gets worse or lasts for more than 3 days. ■ new symptoms occur ■ redness or swelling is present These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

■ this product does not contain directions or complete warnings for adult use

- do not give more than directed ( see overdose warning)
- shake well before using
- mL = milliliter
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.

NDC 82568-0010-8



# Children's Acetaminophen Oral suspension

160mg per 5mL

**PAIN RELIEVER**  
and fever Reducer  
For Ages 2 to 11 Years

Alcohol Free  
Ibuprofen Free  
Aspirin Free  
No Parabens



**Grape Flavor**

8 FL.OZ (236 mL)

## Drug Facts

**Active ingredient (in each 5 mL)** Purpose

Acetaminophen 160 mg Pain reliever / fever reducer

**Uses temporarily:** reduces fever ■ relieves minor aches and pains due to: ■ the common cold ■ flu ■ headache ■ sore throat ■ toothache

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:  
■ more than 5 doses in 24 hours, which is the maximum daily amount  
■ with other drugs containing acetaminophen.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. **Do not use** ■ with any other drug containing acetaminophen (prescription or non prescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ If your child is allergic to acetaminophen or any of the inactive ingredients in this product.

**Ask a doctor before use if your child has liver disease.**

**Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.**

**When using this product, do not exceed recommended dose (see overdose warning)**

**Stop use and ask a doctor if** ■ pain gets worse or lasts for more than 5 days ■ fever gets worse or lasts for more than 3 days. ■ new symptoms occur ■ redness or swelling is present These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

### Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed ( see overdose warning)
- shake well before using
- mL = milliliter
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- repeat dose every 4 hours while symptoms last ■ do not give more than 5 times in 24 hours

## Drug Facts (continued)

Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)*
under 24	under 2	ask a doctor
24 to 35	2 to 3	1 tsp or 5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL
48 to 59	6 to 8	2 tsp or 10 mL
60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
72 to 95	11	3 tsp or 15 mL

\*or as directed by a doctor

**Attention:** use only enclosed dosing cup designed for use with this product. Do not use any other dosing device.

### Other Information

store at room temperature 20-25°C (68-77°F). Protect from Freezing. Protect from Light.

**Inactive ingredients** anhydrous citric acid, D&C red #33, FD&C blue #1, grape flavor, glycerin, microcrystalline cellulose and sodium carboxymethylcellulose, potassium sorbate, purified water, sorbitol solution, sucralose, sucrose, xanthan gum

**Questions or comments?** call toll-free 1-877-225-6999.

\*\*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Infants' Tylenol® Oral Suspension.

**TAMPER EVIDENT: DO NOT USE IF SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

**Aarna USA**  
Manufactured for:  
AARNA USA Inc.  
Leland, NC-28451  
Exclusively distributed by Akron Pharma Inc. Fairfield, NJ-07004



# Children's Acetaminophen Oral suspension

NDC 82568-0010-8



## Children's Acetaminophen Oral suspension

160mg per 5mL

**PAIN RELIEVER** and fever Reducer  
For Ages 2 to 11 Years

Alcohol Free  
Ibuprofen Free  
Aspirin Free  
No Parabens



**Grape Flavor**

8 FL.OZ (236 mL)

### Drug Facts (continued)

Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)*
under 24	under 2	ask a doctor
24 to 35	2 to 3	1 tsp or 5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL
48 to 59	6 to 8	2 tsp or 10 mL
60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
72 to 95	11	3 tsp or 15 mL

\*or as directed by a doctor  
**Attention:** use only enclosed dosing cup designed for use with this product. Do not use any other dosing device.

**Other Information**  
store at room temperature 20-25°C (68-77°F). Protect from Freezing. Protect from Light.

**Inactive Ingredients** anhydrous citric acid, D&C red #33, FD&C blue #1, grape flavor, glycerin, microcrystalline cellulose and sodium carboxymethylcellulose, potassium sorbate, purified water, sorbitol solution, sucralose, sucrose, xanthan gum

**Questions or comments?**  
call toll-free 1-877-225-6999.

\*\* This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Infants' Tylenol® Oral Suspension.

**TAMPER EVIDENT: DO NOT USE IF SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

**Aarna USA**  
Manufactured for:  
AARNA USA Inc.  
Leland, NC-28451



Exclusively distributed by Alron Pharma Inc. Fairfield, NJ-07004

NDC 82568-0010-8



## Children's Acetaminophen Oral suspension

160mg per 5mL

**PAIN RELIEVER** and fever Reducer  
For Ages 2 to 11 Years

Alcohol Free  
Ibuprofen Free  
Aspirin Free  
No Parabens



**Grape Flavor**

8 FL.OZ (236 mL)

### Drug Facts

**Active ingredient (in each 5 mL)** Purpose  
Acetaminophen 160 mg ..... Pain reliever / fever reducer

**Uses** temporarily: ■ reduces fever ■ relieves minor aches and pains due to: ■ the common cold ■ flu ■ headache  
■ sore throat ■ toothache

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:  
■ more than 5 doses in 24 hours, which is the maximum daily amount  
■ with other drugs containing acetaminophen.

### Allergy alert:

Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** ■ with any other drug containing acetaminophen (prescription or non prescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
■ if your child is allergic to acetaminophen or any of the inactive ingredients in this product.

**Ask a doctor before use if your child has liver disease.**

**Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.**

**When using this product, do not exceed recommended dose (see overdose warning)**

**Stop use and ask a doctor if** ■ pain gets worse or lasts for more than 5 days ■ fever gets worse or lasts for more than 3 days.  
■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**  
■ this product does not contain directions or complete warnings for adult use

■ do not give more than directed ( see overdose warning)

■ shake well before using

■ mL = milliliter

■ find right dose on chart below. If possible, use weight to dose; otherwise, use age.

■ repeat dose every 4 hours while symptoms last

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

■ do not give more than 5 times in 24 hours

### Drug Facts (continued)

■ mL = milliliter find right dose on chart below. If possible, use weight to dose; otherwise, use age. ■ repeat dose every 4 hours while symptoms last ■ do not give more than 5 times in 24 hours

Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)*
under 24	under 2	ask a doctor
24 to 35	2 to 3	1 tsp or 5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL
48 to 59	6 to 8	2 tsp or 10 mL
60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
72 to 95	11	3 tsp or 15 mL

\*or as directed by a doctor

**Attention:** use only enclosed dosing cup designed for use with this product. Do not use any other dosing device.

### Other Information

store at room temperature 20-25°C (68-77°F). Protect from Freezing. Protect from Light.

**Inactive Ingredients** anhydrous citric acid, D&C red #33, FD&C blue #1, grape flavor, glycerin, microcrystalline cellulose and sodium carboxymethylcellulose, potassium sorbate, purified water, sorbitol solution, sucralose, sucrose, xanthan gum

**Questions or comments?** call toll-free 1-877-225-6999.

\*\* This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark Tylenol® oral suspension

**TAMPER EVIDENT: DO NOT USE IF SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

**Aarna USA**  
Manufactured for:  
AARNA USA Inc.  
Leland, NC-28451



Exclusively distributed by Alron Pharma Inc. Fairfield, NJ-07004

NDC 82568-0010-6



## Children's Acetaminophen Oral suspension

160mg per 5mL

**PAIN RELIEVER** and fever Reducer  
For Ages 2 to 11 Years

Alcohol Free  
Ibuprofen Free  
Aspirin Free  
No Parabens



**Grape Flavor**

16 FL.OZ (473 mL)

### Drug Facts

**Active ingredient (in each 5 mL)** Purpose  
Acetaminophen 160 mg ..... Pain reliever / fever reducer

**Uses** temporarily: ■ reduces fever ■ relieves minor aches and pains due to: ■ the common cold ■ flu ■ headache ■ sore throat ■ toothache

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:  
■ more than 5 doses in 24 hours, which is the maximum daily amount  
■ with other drugs containing acetaminophen.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### Do not use

■ with any other drug containing acetaminophen (prescription or non prescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If your child is allergic to acetaminophen or any of the inactive ingredients in this product.

**Ask a doctor before use if your child has liver disease.**

**Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.**

**When using this product, do not exceed recommended dose (see overdose warning)**

### Stop use and ask a doctor if

■ pain gets worse or lasts for more than 5 days ■ fever gets worse or lasts for more than 3 days. ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

### Directions

■ this product does not contain directions or complete warnings for adult use, ■ do not give more than directed ( see overdose warning)

■ shake well before using

**CHILDRENS ACETAMINOPHEN**

childrens acetaminophen oral suspension

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:82568-0010
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>D&amp;C RED NO. 33</b> (UNII: 9DBA0SBB0L)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>CARBOXYMETHYLCELLULOSE SODIUM</b> (UNII: K679OBS311)	
<b>POTASSIUM SORBATE</b> (UNII: 1VPU26JZZ4)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>SUCROSE</b> (UNII: C151H8M554)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

**Product Characteristics**

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	GRAPE	<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:82568-0010-2	60 mL in 1 BOTTLE; Type 0: Not a Combination Product	06/20/2025	
2	NDC:82568-0010-4	118 mL in 1 BOTTLE; Type 0: Not a Combination Product	06/20/2025	
3	NDC:82568-0010-8	236 mL in 1 BOTTLE; Type 0: Not a Combination Product	06/20/2025	
4	NDC:82568-0010-6	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	06/20/2025	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	06/20/2025	

**Labeler** - AARNA USA INC. (118515992)

Revised: 6/2025

AARNA USA INC.