

FLEET- glycerin liquid
C.B. Fleet Company, Inc.

Fleet Glycerin Liquid 0132-018

FLEET- glycerin liquid
C.B. Fleet Company, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active Ingredient (in each 5.4 mL average delivered dose)

Glycerin 5.4 g

Purpose

Hyperosmotic Laxative

Use

for relief of occasional constipation.
This product usually produces a bowel movement in 15 minutes to 1 hour.

Warnings

May cause rectal discomfort or a burning sensation.

For rectal use only.

Do not use when abdominal pain, nausea, or vomiting are present unless directed by a doctor.

Ask a doctor before using any laxative if you

- have a sudden change in bowel habits lasting more than 2 weeks
- already used a laxative for more than 1 week

Stop using this product and consult a doctor if you have

- rectal bleeding
- no bowel movement within 1 hour of using this product

These symptoms may indicate a serious condition.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Single Daily Dosage (per 24 hours)

Adults and children 6 years and over 1 suppository or as directed by a doctor

Children 2 to under 6 years use Fleet® Pedia-Lax® Liquid

Children under 2 years consult a doctor

Left-side position: Lie on left side with right knee bent and arms resting comfortably.

Knee-chest position: Kneel, then lower head and chest forward until left side of face is resting on surface with left arm folded comfortably

Administering the laxative:

CAUTION: REMOVE ORANGE PROTECTIVE SHIELD BEFORE INSERTING.

Hold unit upright, grasping bulb of unit with fingers. Grasp orange protective shield with other hand, pull gently to remove. With steady pressure, insert tip into rectum with a slight side-to-side movement, with tip pointing toward navel. Do not force the applicator tip into the rectum, as this can cause injury. Insertion may be easier if person receiving suppository bears down as if having a bowel movement. This helps relax the muscles around the anus. Squeeze the bulb until nearly all liquid is expelled. While continuing to squeeze the bulb, remove tip from rectum and discard unit. It is not necessary to empty unit completely. The unit contains more than the amount of liquid needed for effective use. A small amount of liquid will remain in the unit after use.

For a more detailed step-by-step instructional video on how to use Fleet® products, visit our website www.fleetlabs.com to watch the “how to use” video.

Inactive Ingredients

edetate disodium, purified water

Questions?

1-866-255-6960 or www.fleetlabs.com

PRINCIPAL DISPLAY PANEL

Fleet Laxative
 Liquid Glycerin Suppositories
 4 Adult Suppositories
 7.5 ml each



FLEET			
glycerin liquid			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0132-0185
Route of Administration	RECTAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
GLYCERIN (UNII: PDC6A3C00X) (GLYCERIN - UNII:PDC6A3C00X)		GLYCERIN	5.4 g
Inactive Ingredients			
Ingredient Name			Strength
EDETATE DISODIUM (UNII: 7FLD91C86K)			
WATER (UNII: 059QF0K00R)			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0132-0185-82	4 in 1 CARTON; Type 0: Not a Combination Product	06/03/2002	
2	NDC:0132-0185-01	1 in 1 PACKAGE; Type 0: Not a Combination Product	06/03/2002	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	06/03/2002	

Labeler - C.B. Fleet Company, Inc. (003119054)

Revised: 10/2023

C.B. Fleet Company, Inc.