

CLOTRIMAZOLE- clotrimazole cream
Rite Aid Corporation

Clotrimazole Vaginal Cream, USP

Drug Facts

Active ingredients

Clotrimazole, USP 1% (50 mg in each applicatorful)

Clotrimazole, USP 1% (external cream)

Purpose

Vaginal antifungal

Vaginal antifungal

Uses

- treats vaginal yeast infections
- relieves external itching and irritation due to a vaginal yeast infection

Warnings

For vaginal use only

Do not use

if you have never had a vaginal yeast infection diagnosed by a doctor.

Ask a doctor before use if you have

- **vaginal itching and discomfort for the first time**
- **lower abdominal, back or shoulder pain, fever, chills, nausea, vomiting, or foul-smelling vaginal discharge. You may have a more serious condition.**
- vaginal yeast infections often (such as once a month or 3 in 6 months). You could be pregnant, or have a serious underlying medical cause for your symptoms, including diabetes or a weakened immune system.
- been exposed to the human immunodeficiency virus (HIV) that causes AIDS

When using this product

- do not use tampons, douches, spermicides, or other vaginal products. Condoms and diaphragms may be damaged and fail to prevent pregnancy or sexually transmitted

diseases (STDs).

- do not have vaginal intercourse
- mild increase in vaginal burning, itching or irritation may occur
- if you do not get complete relief, ask a doctor before using another product

Stop use and ask a doctor if

- **symptoms do not get better in 3 days**
- **symptoms last more than 7 days.**
- **you get a rash or hives, abdominal pain, fever, chills, nausea, vomiting, or a foul-smelling vaginal discharge**

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- before using this product, read the enclosed educational brochure for complete directions and information
- adults and children 12 years of age and over:
 - **vaginal cream:** insert one applicatorful of cream into the vagina at bedtime for 7 days in a row. Wash applicator after each use.
 - **external cream:** use the same tube of cream if you have itching and irritation on the skin outside the vagina. Squeeze a small amount of cream onto your fingertip. Apply to itchy, irritated skin outside the vagina. Use 2 times daily for up to 7 days as needed.
- children under 12 years of age: ask a doctor

Other information

- to open tube: unscrew cap, lift tab, and pull to remove foil seal prior to use
- do not use if foil seal on tube opening is broken or missing
- do not purchase if carton is open
- store at room temperature 20° to 25°C (68° to 77°F)
- before using any medication, read all label directions. Keep carton, it contains important information

Inactive ingredients

benzyl alcohol, cetostearyl alcohol, cetyl esters wax, 2-octyldodecanol, polysorbate 60, purified water, sodium phosphate monobasic, sorbitan monostearate

Questions?

call **1-877-753-3935** Monday-Friday 9AM-5PM ET

principal display panel

Clotrimazole

Vaginal Cream, USP 1%,

Vaginal Antifungal

7 - DAY TREATMENT

VAGINAL ANTIFUNGAL

Clotrimazole vaginal cream can kill the yeast that may cause vaginal infection

Cure most vaginal yeast infections

1 reusable applicator (7-Day Therapy)

NET WT 1.59 OZ (45g) TUBE OF VAGINAL CREAM (1%) AND

1 REUSABLE PLASTIC APPLICATOR (7-DAY THERAPY)

DISTRIBUTED BY: RITE AID,

200 NEWBERRY COMMONS

ETTERS, PA 17319

Package label

NDC 11822-0220-6

Clotrimazole
Vaginal Cream, USP 1%,
Vaginal Antifungal

**7-DAY TREATMENT
VAGINAL ANTIFUNGAL**

Clotrimazole vaginal cream can kill the yeast that may cause vaginal infection
Cures most vaginal yeast infections
One reusable applicator (7-Day Therapy)

NET WT 1.59 OZ (45 g) TUBE
OF VAGINAL CREAM (1%) &
ONE REUSABLE APPLICATOR
(7-DAY THERAPY)

Educational
Pamphlet
Enclosed

Actual Size

<p>Drug Facts</p> <p>Active Ingredients Clotrimazole, USP 1% (50 mg in each applicator)..... Vaginal antifungal Clotrimazole, USP 1% (external cream)..... Vaginal antifungal</p> <p>Uses ■ treats vaginal yeast infections ■ relieves external itching and irritation due to a vaginal yeast infection</p> <p>Warnings Do not use if you have never had a vaginal yeast infection diagnosed by a doctor. Do not use if you have ever had a vaginal yeast infection diagnosed by a doctor. ■ lower abdominal, back or muscular pain, fever, chills, nausea, vomiting, or fluid-appearing vaginal discharge. You may have a more serious condition. ■ vaginal yeast infections often recur as soon as a month or 3 to 6 months. You could be pregnant, or have a serious underlying medical cause for your symptoms, including diabetes or a weakened immune system. ■ been exposed to the human immunodeficiency virus (HIV) that causes AIDS</p> <p>When using this product ■ do not use tampons, douches, spermicides, or other vaginal products. ■ condoms and diaphragms may be damaged and fail to prevent pregnancy or sexually transmitted diseases (STDs). ■ do not have vaginal intercourse ■ mild increase in vaginal burning, itching or irritation may occur ■ do not use tight clothing ■ ask a doctor before using bubble bath products</p> <p>Keep out of reach of children. ■ symptoms do not get better in 7 days ■ symptoms last more than 7 days ■ you are pregnant or trying to get pregnant ■ have had yeast infections, urinary tract infections, bacterial vaginosis, or other vaginal infections</p>	<p>Drug Facts (continued)</p> <p>Directions If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.</p> <p>Directions ■ before using this product, read the enclosed educational brochure for complete directions and information ■ adults and children 12 years of age and over: ■ vaginal cream: insert one applicatorful of cream into the vagina at bedtime for 7 days in a row. ■ external cream: use the same tube of cream if you have itching and irritation on the skin outside the vagina. Squeeze a small amount of cream onto your fingertip. Apply to itchy, irritated skin outside the vagina. Use 2 times daily for up to 7 days as needed. ■ children under 12 years of age: ask a doctor</p> <p>Other Information ■ to open tube: unscrew cap, lift tab, and pull to remove foil seal prior to use ■ do not use if foil seal on tube opening is broken or missing ■ do not permix if certain to open ■ do not use if you are pregnant or trying to get pregnant ■ before using any medication, read all label directions. Keep track. It contains important information.</p> <p>Inactive Ingredients benzyl alcohol, cetylstearyl alcohol, cetyl stearic wax, 2-ethylhexanoic polyacrylate 60, purified water, sodium phosphate monobasic, sodium monostearate</p> <p>Questions? Call 1-877-765-3635 Monday-Friday 9AM-5PM ET.</p>
--	--

RITE AID Vaginal Antifungal

CLOTRIMAZOLE

clotrimazole cream

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-0220
Route of Administration	VAGINAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CLOTRIMAZOLE (UNII: G07GZ97H65) (CLOTRIMAZOLE - UNII:G07GZ97H65)	CLOTRIMAZOLE	10 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
BENZYL ALCOHOL (UNII: LKG8494WBH)	
CETOSTEARYL ALCOHOL (UNII: 2DMT128M1S)	
CETYL ESTERS WAX (UNII: D072FFP9GU)	
OCTYLDODECANOL (UNII: 461N1O614Y)	
POLYSORBATE 60 (UNII: CAL22UVI4M)	
WATER (UNII: 059QF0KO0R)	
SODIUM PHOSPHATE, MONOBASIC, MONOHYDRATE (UNII: 593YOG76RN)	
SORBITAN MONOSTEARATE (UNII: NVZ4I0H58X)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-0220-6	1 in 1 CARTON	11/22/1996	
1		45 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074165	11/22/1996	01/01/2027

Labeler - Rite Aid Corporation (014578892)

Revised: 10/2025

Rite Aid Corporation