

**AROAMAS COT NICOTINE TRANSDERMAL PATCHES KIT- nicotine transdermal patches patch**  
**PCHEERY LLC**

-----  
**Initial Drug Listing - Aroamas Cot Patch Kit**

Nicotine 21mg, 14mg, 7mg

menthol 0.3%

Stop smoking aid

reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking.

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- an allergy to adhesive tape or have skin problems, because you are more likely to get rashes.
- stomach ulcer or diabetes.
- history of seizures.
- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.
- if you have vivid dreams or other sleep disturbances remove this patch at bedtime.
- skin redness caused by the patch does not go away after four days, or if your skin swells, or you get a rash.
- irregular heartbeat or palpitations occur.
- you get symptoms of nicotine overdose, such as nausea, vomiting, dizziness, weakness and rapid heartbeat.
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Used patches have enough nicotine to poison children and pets. If swallowed, get medical help or contact a Poison Control Center right away. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.

- **if you are under 18 years of age, ask a doctor before use.**
- before using this product, read the enclosed self-help guide for complete directions and other information
- begin using the patch on your quit day
- if you smoke **more than 10 cigarettes per day**, use the following schedule below:

### STEP 1

Use one 21 mg patch/day

Weeks 1-6

### STEP 2

Use one 14 mg patch/day

Weeks 7-8

### STEP 3

Use one 7 mg patch/day

Weeks 9-10

- if you smoke **10 or less cigarettes per day**, do not use **Step 1(21mg)**, start with **Step 2(14mg)** for 6 weeks, then **Step 3(7mg)** for 2 weeks and then stop.
- apply one new patch every 24 hours on skin that is dry, clean and hairless.
- remove backing from patch and immediately press onto skin. Hold for 10 seconds.
- wash hands after applying or removing patch. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.
- the used patch should be removed and a new one applied to a different skin site at the same time each day.
- if you have vivid dreams, you may remove the patch at bedtime and apply a new one in the morning.
- you may wear the patch for 18-24 hours.
- do not wear more than one patch at a time
- do not cut patch in half or into smaller pieces
- do not leave patch on for more than 24 hours because it may irritate your skin and loses strength after 24 hours
- to avoid possible burns, remove patch before undergoing any MRI (magnetic resonance imaging) procedures
- It is important to complete treatment. If you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider.

CODONOPSIS PILOSULA ROOT, CHRYSANTHEMUM INDICUM FLOWER, CLOVE LEAF OIL, TOBACCO LEAF, CAMPHOR OIL

Comments or Questions? Call toll-free: 833-276-2627 - Or visit us at [aroamascot.com](http://aroamascot.com)

PANTONE Black 6 C  
RGB 16 24 32 / CMYK 100 79 49 93

PANTONE P 89-16 C  
CMYK 100 94 0 10

PANTONE P 37-6 C  
CMYK 0 79 94 0

PANTONE P 186-7 C  
CMYK 81 0 92 22



## AROAMAS COT NICOTINE TRANSDERMAL PATCHES KIT

nicotine transdermal patches patch

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:85831-032
Route of Administration	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MENTHOL (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A)	MENTHOL	0.3 g in 100
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	18 g in 100

## Inactive Ingredients

Ingredient Name	Strength
<b>CHRYSANTHEMUM INDICUM FLOWER</b> (UNII: I6OER6U04L)	
<b>CLOVE LEAF OIL</b> (UNII: VCA5491KVF)	
<b>CAMPHOR OIL</b> (UNII: 75IZZ8Y727)	
<b>CODONOPSIS PILOSULA ROOT</b> (UNII: 02VJ3CEY9G)	
<b>TOBACCO LEAF</b> (UNII: 6YR2608RSU)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:85831-032-04	56 in 1 CARTON	06/30/2025	
1		1 in 1 PATCH; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	06/30/2025	

**Labeler** - PCHEERY LLC (061286986)

Revised: 6/2025

PCHEERY LLC