SILACE- docusate sodium liquid Lannett Company, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Silace Liquid

Active Ingredient: Docusate sodium 10 mg (in each 1 mL)

Purpose: Stool Softener

Uses

- for gentle, reliable relief from occasional constipation (irregularity)
- generally produces bowel movement in 12 to 72 hours

Warning

- **Do not use** laxative products for longer than 1 week unless told to do so by a doctor
- if you are presently taking mineral oil unless told to do by a doctor

Ask doctor before use if you have

- stomach pain
- nausea
- vomiting
- noticed a sudden change in bowel habits that last over two weeks

Stop use and ask a doctor if you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take as indicated by doctor
- this product must be given in a 6 to 8 oz. glass of milk or fruit juice or infant's formula to prevent throat irritation. Dose may be taken as a single daily dose or in

divided doses

- dosage should be adjusted to individual response
- higher doses are recommended for initial therapy
- the effect on stools is usually apparent 1 to 3 days after the first dose

Adults and children over 12 years	50 to 200 mg (1 to 4 teaspoonful)
Children 6 to under 12 years	50 to 100 mg (1 to 2 teaspoonful)
Children 3 to under 6 years	25 to 50 mg(1/2 to 1 teaspoonful)
Children under 3 years of age	Ask a doctor

Other information

- store at room temperature 20°-25°C (68°-77°F)
- protect from freezing and excessive heat
- do not use if tamperevident safety seal around cap is broken or missing
- dispense in tight, light-resistant container with a child-resistant closure

Inactive ingredients

citric acid, D&C red no. 33, flavor, methylparaben, Poloxamer, propylene glycol, propylparaben, purified water, sodium citrate.

Questions

888-974-5279

This product is not manufactured or distributed by Purdue Products LP, distributor of Colace® Liquid.

Manufactured by:

Silarx Pharmaceuticals, Inc. 1033 Stoneleigh Ave Carmel, NY 10512 USA

Drug Facis		ISLARX
Active ingredient (in each 1 mL) Docusate sodium 10 mo) Purpose Stool softener	
generally produces bowel movement	ccasional constipation (irregularity)	NDC 54838-116-80
Varnings Do not use - lacative products by a doctor - if you are presently taking min	for longer than 1 week unless told to do so neral oil unless told to do so by a doctor	SILACE
Lak a doctor before use if you have - noticed a sudden change in bowel ha	stomach pain - nausea - vomiting bits that last over two weeks	
Stop use and ask a doctor if you have recta novement after use of a laxative. These	I bleeding or fall to have a bowel e could be signs of a serious condition	LIQUID
f pregnant or breast feeding, ask a h out of reach of children. In case of ov		Docusate Sodium
oison Control Center right away.		(40) Oalutian)
Directions - take as directed by a doctor - this product must be given in a 6 to		(1% Solution)
8 oz. glass of milk or fruit juice to prevent throat irritation. Dose may be taken as a single daily dose or in divided doses - dosage should be adjusted to		Stool Softener
ndividual response - higher doses are		
the effect on stools is usually appare	Stimulant Free	
iduits and children 12 years of age and older shildren 6 to under 12 years of age		
shildren 3 to under 6 years of age	50 to 100 mg (1 to 2 teaspoonful) 25 to 50 mg (1/2 to 1 teaspoonful)	BULK CONTAINER – Not for Household Use
shildren under 3 years of age	ask a doctor	Pharmacist - Dispense in a tight,
Other information - store at room protect from freezing and excessive it	temperature 20°-25° C (68°-77°F) neat	light-resistant container with child-resistant closure.
inactive ingradients citric acid, D&C red	no. 33, lemon-vanilia flavor, methylparaber	
poloxamer, propylene glycol, propylparabe	en, purifed water, sodium citrate	TAMPER EVIDENT:
Questions? © 1-844-834-0530		Do not use if tamper-evident
	This product is not manufactured or distributed by Purdue Products LP, distributor of Colace® Liquid.	safety seal around cap is broken or missing
54838-116-80 9	10-1035 Rev. 04/18	† Compare to the active
		ingredient of Colace® Liquid
		1 Pint (473 mL)
		Manufactured by: Silarx Pharmaceuticals, Inc.

SILACE	
--------	--

docusate sodium liquid **Product Information** Item Code (Source) Product Type HUMAN OTC DRUG NDC:54838-116 **Route of Administration** ORAL Active Ingredient/Active Moiety Basis of Strength Ingredient Name

Strength

	active Ingr	Cuic					
Ingredient Name						Strength	
AN	IHYDROUS CIT	ric a	CID (UNII: XF417D3PSL)				
Dδ	C RED NO. 33	(UNII	9DBA0SBB0L)				
	THYLPARABEI						
PR	OPYLENE GLY	COL	UNII: 6DC9Q167V3)				
	OPYLPARABEN						
	DIUM CITRATI		-				
W	ATER (UNII: 059	QF0K	D0R)				
PO	LOXAMER 407	(UNI	: TUF2IVW3M2)				
Pr	roduct Cha	ract	eristics				
Co	lor				Score		
Sh	аре	e		Size			
Fla	avor	r LEMON (Lemon Vanilla Flavor)			Imprint Code		
Co	ontains						
Pa	ackaging						
#	ltem Code		Package Description	Mai	keting Start Date	Marketing End Date	
	NDC:54838- 116-80		mL in 1 BOTTLE, PLASTIC; Type 0: Not a bination Product	10/05	/1990	11/30/2024	
		Inf	ormation				
Μ	arketing	Im					
M	arketing Marketing Category	Ini	Application Number or Monograph Citation	Mark	eting Start Date	Marketing End Date	

Labeler - Lannett Company, Inc. (002277481)

Revised: 8/2018

Lannett Company, Inc.