

KAOPECTATE- bismuth subsalicylate tablet, coated
Kramer Laboratories

Drug Facts

Active ingredient (per caplet)	Purposes
Bismuth subsalicylate 262 mg	Anti-diarrheal, Upset stomach reliever

Uses relieves:

- traveler's diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
 - heartburn
 - indigestion
 - nausea
 - gas
 - belching
 - fullness

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy Alert: Contains salicylate.

Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use if you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if

- symptoms get worse
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **adults and children 12 years of age and older:** 2 caplets (1 dose)
- swallow caplets with water; do not chew
- repeat dose every 1/2 hour to 1 hour as needed
- do not exceed 8 doses in 24 hours
- use until diarrhea stops but not more than 2 days
- **children under 12 years:** ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

- Each caplet contains very low sodium
- **do not use if carton was opened or if printed foil inner seal under bottle cap is torn or missing**
- store at room temperature 20°-25° C (68°-77° F)

Inactive ingredients

croscarmellose sodium, crospovidone, hydrated silica, hypromellose, magnesium stearate, microcrystalline cellulose, mineral oil, povidone, sodium lauryl sulfate, stearic acid, talc, titanium dioxide, triacetin

Principal Display Panel

FAST & EFFECTIVE RELIEF

Kaopectate®

Bismuth Subsalicylate

• *Anti-Diarrheal* • *Upset Stomach Reliever*

Diarrhea & Upset Stomach

- ✓ Begins controlling symptoms from the first dose
- ✓ Quickly relieves urgency, gas, and cramping
- ✓ Effective on diarrhea from bacteria, viruses, and other causes

CLINICALLY PROVEN MEDICINE

28 Coated Caplets



LOT

EXP

Do not use if foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken.

Kramer Laboratories, Inc.

Bridgewater, NJ 08807

1-800-824-4894

14711-07-20_K0720

Kaopectate[®]

Bismuth Subsalicylate
• Anti-Diarrheal • Upset Stomach Reliever

Drug Facts

Active ingredient (per caplet) Bismuth subsalicylate 262 mg.....**Purposes** Anti-diarrheal, Upset stomach reliever

Uses relieves: • traveler's diarrhea • diarrhea • upset stomach due to overindulgence in food and drink, including: • heartburn • indigestion • nausea • gas • belching • fullness

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy Alert: Contains salicylate. Do not take if you are
• allergic to salicylates (including aspirin) • taking other salicylate products

Drug Facts (continued)

Do not use if you have • an ulcer • a bleeding problem • bloody or black stool

Ask a doctor before use if you have • fever • mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for
• anticoagulation (thinning the blood) • diabetes • gout • arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if • symptoms get worse
• ringing in the ears or loss of hearing occurs • diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- adults and children 12 years of age and older: 2 caplets (1 dose)
- swallow caplets with water; do not chew
- repeat dose every 1/2 hour to 1 hour as needed
- do not exceed 8 doses in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

- Each caplet contains very low sodium
- do not use if carton was opened or if printed foil inner seal under bottle cap is torn or missing
- store at room temperature 20°-25° C (68°-77° F)

Drug Facts (continued)


Inactive ingredients

croscarmellose sodium, crospovidone, hydrated silica, hypromellose, magnesium stearate, microcrystalline cellulose, mineral oil, povidone, sodium lauryl sulfate, stearic acid, talc, titanium dioxide, triacetin

Kramer Laboratories, Inc.
Bridgewater, NJ 08807
1-800-824-4894

Do not use if foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken.

LOT
EXP



Kaopectate®

Bismuth Subsalicylate

- **Anti-Diarrheal**
- **Upset Stomach Reliever**

Diarrhea & Upset Stomach

28 Coated Caplets

Do not use if foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken.

KEEP CARTON AS IT CONTAINS IMPORTANT INFORMATION

Kramer Laboratories, Inc.

Bridgewater, NJ 08807

1-800-824-4894

14712-07-20_K0720



Active ingredient (in each caplet)
Bismuth subsalicylate 262 mg

Purposes Anti-diarrheal, Upset stomach
reliever **Uses relieves:** • traveler's
diarrhea • diarrhea • upset stomach
due to overindulgence in food and drink,
including: • heartburn • indigestion •
nausea • gas • belching • fullness

Warnings **Reye's syndrome:** Children
and teenagers who have or are recovering
from chicken pox or flu-like symptoms
should not use this product. When using
this product, if changes in behavior with
nausea and vomiting occur, consult a
doctor because these symptoms could
be an early sign of Reye's syndrome, a
rare but serious illness. **Allergy alert:**
Contains salicylate. Do not take if you
are • allergic to salicylates (including
aspirin) • taking other salicylate products
Do not use if you have • an
ulcer • a bleeding problem
• bloody or black stool **Ask**
a doctor before use if you
have • fever • mucus in the stool **Ask a**
doctor or pharmacist before use if you
are taking any drug for • anticoagulation
(thinning the blood) • diabetes • gout •

PEEL HERE ▼

arthritis

When using this product a temporary,
but harmless, darkening of the stool
and/or tongue may occur.

Stop use and ask a doctor if
• symptoms get worse • ringing in the
ears or loss of hearing occurs
• diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a
health professional before use.

Keep out of reach of children. In case
of overdose, get medical help or contact
a Poison Control Center right away.

Directions • adults and children 12 years
of age and older: 2 caplets (1 dose)
• swallow caplets with water; do not chew
• repeat dose every 1/2 hour to 1 hour
as needed • do not exceed 8 doses in 24
hours • use until diarrhea stops but not
more than 2 days • children under 12
years: ask a doctor • drink plenty of clear
fluids to help prevent dehydration caused
by diarrhea

Other information • Each caplet contains

Continued ▼

very low sodium • **do not use if carton
was opened or if printed foil inner seal
under bottle cap is torn or missing**
• store at room temperature 20°-25° C
(68°-77° F)

Inactive ingredients croscarmellose
sodium, crospovidone, hydrated silica,
hypromellose, magnesium stearate,
microcrystalline cellulose, mineral oil,
povidone, sodium lauryl sulfate, stearic
acid, talc, titanium dioxide, triacetin

**Do not use if foil seal under
cap, printed with "SEALED for
YOUR PROTECTION"
is missing, open or broken.**

**KEEP CARTON AS IT CONTAINS
IMPORTANT INFORMATION**

Kramer Laboratories, Inc.
Bridgewater, NJ 08807
1-800-824-4894

14712-07-20_K0720

Kaopectate®
Bismuth Subsalicylate

- **Anti-Diarrheal**
- **Upset Stomach Reliever**

Diarrhea & Upset Stomach

84 Coated Caplets

Do not use if foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken.

K0625

LOT:

EXP:

R54039

Kramer Laboratories, Inc. Bridgewater, NJ 08807
1-800-824-4894



<p>Drug Facts (continued)</p> <p>Do not use if you have • an ulcer • a bleeding problem • bloody or black stool</p> <p>Ask a doctor before use if you have • fever • muscle in the stool</p> <p>Ask a doctor or pharmacist before use if you are taking any drug for • anticoagulation (thinning the blood) • diabetes • gout • arthritis</p> <p>When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur</p> <p>Stop use and ask a doctor if • symptoms get worse • ringing in the ears or loss of hearing occurs • diarrhea lasts more than 2 days</p> <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none"> adults and children 12 years of age and older: 2 caplets (1 dose) every 1/2 hour to 1 hour as needed swallow caplets with water, do not chew do not exceed 8 doses (16 caplets) in 24 hours use until diarrhea stops but not more than 2 days children under 12 years: ask a doctor drink plenty of clear fluids to help prevent dehydration caused by diarrhea <p>Other information</p> <ul style="list-style-type: none"> each caplet contains: salicylate 102mg and very low sodium store at room temperature 20°-25° C (68°-77° F) <p>Inactive ingredients</p> <p>croscarmellose sodium, crospovidone, hydrated silica, hypromellose, magnesium stearate, microcrystalline cellulose, mineral oil, povidone, sodium lauryl sulfate, stearic acid, talc, titanium dioxide, triacetin</p>	<p>Kramer Laboratories, Inc., Bridgewater, NJ 08807 1-800-824-4894</p>
---	--

KAOPECTATE

bismuth subsalicylate tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55505-197
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Bismuth Subsalsicylate (UNII: 62TEY51RR1) (Bismuth Cation - UNII:ZS9CD18YE)	Bismuth Subsalsicylate	262 mg

Inactive Ingredients

Ingredient Name	Strength
Croscarmellose Sodium (UNII: M28OL1HH48)	
Crospovidone (UNII: 2S7830E561)	
Hydrated Silica (UNII: Y6O7T4G8P9)	
Hypromellose, Unspecified (UNII: 3NXW29V3WO)	
Magnesium Stearate (UNII: 70097M6I30)	
Microcrystalline Cellulose (UNII: OP1R32D61U)	
Mineral Oil (UNII: T5L8T28FGP)	
Povidone, Unspecified (UNII: FZ989GH94E)	
Sodium Lauryl Sulfate (UNII: 368GB5141J)	
Stearic Acid (UNII: 4ELV7Z65AP)	
Talc (UNII: 7SEV7J4R1U)	
Titanium Dioxide (UNII: 15FIX9V2JP)	
Triacetin (UNII: XHX3C3X673)	

Product Characteristics

Color	WHITE	Score	no score
Shape	OVAL (caplet)	Size	17mm
Flavor		Imprint Code	K

Contains**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55505-197-07	1 in 1 CARTON	10/01/2020	
1		28 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:55505-197-70	1 in 1 CARTON	02/01/2021	
2		42 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:55505-197-78	2 in 1 PACKAGE	04/23/2025	
3		1 in 1 CARTON		
3		42 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:55505-197-77	1 in 1 CARTON	11/15/2025	
4		84 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	10/01/2020	

Labeler - Kramer Laboratories (122720675)

Revised: 12/2025

Kramer Laboratories