

**GUAIFENESIN AND DEXTROMETHORPHAN HYDROBROMIDE- guaifenesin and dextromethorphan hydrobromide syrup**  
**Method Pharmaceuticals**

-----

**Guaifenesin and Dextromethorphan Hydrobromide Syrup**

**Active ingredients**

Dextromethorphan HBr, USP 10 mg  
Guaifenesin, USP 100 mg

**Purpose**

Cough suppressant  
Expectorant

**Uses**

- temporarily relieves cough due to minor throat and bronchial irritation
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive

**Warnings**

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- persistent cough or chronic cough such as occurs with smoking, asthma, chronic bronchitis, emphysema
- cough accompanied by excessive phlegm (mucus)

**Stop use and ask a doctor if**

- cough lasts more than 7 days or occurs with fever, rash, or headaches that lasts. This could be a sign of a serious condition
- hypersensitive to any ingredients

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not take more than 6 doses in any 24-hour period. **This adult product is not intended for use in children under 12 years of age**

adults and children 12 years and over	2 teaspoonfuls (TSP) every 4 hours
children under 12 years	<b>DO NOT USE</b>

**Inactive ingredients**

Citric Acid Anhydrous, USP; Sodium Citrate Dihydrate, USP; Sodium Saccharin, USP; Propylene Glycol, USP; Grape Flavor; Sorbitol Solution, USP; Glycerin, USP; Maltitol Solution, NF

**Other information**

- store at room temperature 20°-25°C (68°-77°F)

**Questions**

1-877-250-3427

**Principal Display Panel**

Method Pharmaceuticals

NDC 58657-505-08

Guaifenesin and Dextromethorphan Hydrobromide Syrup

100 mg/10 mg/5 mL

Expectorant

Cough Suppressant

Grap Flavor

**Non Drowsy • Alcohol Free**

FOR AGES 12 AND UP

**8 fl. oz. (237 mL)**

Manufactured For:

Method Pharmaceuticals, LLC

Fort Worth, TX 76118 Rev. 06/2016

 <p><b>METHOD</b> PHARMACEUTICALS</p> <p>NDC 58657-505-08</p> <p><b>Guaifenesin and Dextromethorphan Hydrobromide Syrup</b></p> <p>100 mg/10 mg/5 mL</p> <p><b>Expectorant Cough Suppressant</b></p> <p>Grape Flavor</p> <p>Non Drowsy • Alcohol Free</p> <p>FOR AGES 12 AND UP</p> <p><b>8 fl. oz. (237mL)</b></p>	<p><b>Drug Facts</b></p> <p><b>Active ingredient (in each 5 mL)</b>      <b>Purpose</b></p> <p>Dextromethorphan Hydrobromide 10 mg ..... Cough Suppressant</p> <p>Guaifenesin 100 mg ..... Expectorant</p> <p><b>Uses</b></p> <ul style="list-style-type: none"> <li>temporarily relieves cough due to minor throat and bronchial irritation</li> <li>helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive</li> </ul> <p><b>Warnings</b></p> <p>Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> <li>persistent cough or chronic cough such as occurs with smoking, asthma, chronic bronchitis, emphysema</li> <li>cough accompanied by excessive phlegm (mucus)</li> </ul> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> <li>cough lasts more than 7 days or occurs with fever, rash, or headaches that lasts. This could be a sign of a serious condition</li> <li>hypersensitive to any ingredients</li> </ul> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>	<p><b>Drug Facts (continued)</b></p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>do not take more than 6 doses in any 24-hour period. This adult product is not intended for use in children under 12 years of age</li> </ul> <table border="1"> <tr> <td>adults and children 12 years and over</td> <td>2 teaspoonfuls (TSP) every 4 hours</td> </tr> <tr> <td>children under 12 years</td> <td><b>DO NOT USE</b></td> </tr> </table> <p><b>Inactive ingredients</b></p> <p>Citric Acid Anhydrous, USP; Glycerin, USP; Grape Flavor; Maltitol Solution, NF; Propylene Glycol, USP; Purified Water; Sodium Citrate Dihydrate, USP; Sodium Saccharin, USP; Sorbitol Solution, USP</p> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>store at room temperature 20°-25°C (68°-77°F)</li> </ul> <p><b>Questions</b> 1-877-250-3427</p> <p>Manufactured For: Method Pharmaceuticals, LLC Fort Worth, TX 76118</p> <p>Rev. 06/2016</p>  <p>3 58657-50508 0</p> <p>Lot: _____ Exp.: _____</p>	adults and children 12 years and over	2 teaspoonfuls (TSP) every 4 hours	children under 12 years	<b>DO NOT USE</b>
	adults and children 12 years and over	2 teaspoonfuls (TSP) every 4 hours				
children under 12 years	<b>DO NOT USE</b>					
<p><b>GUAIFENESIN AND DEXTROMETHORPHAN HYDROBROMIDE</b> guaifenesin and dextromethorphan hydrobromide syrup</p>						

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58657-505
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)		DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 5 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)		GUAIFENESIN	100 mg in 5 mL
Inactive Ingredients			
Ingredient Name			Strength
GLYCERIN (UNII: PDC6A3C0OX)			
MALTITOL (UNII: D65DG142WK)			
SACCHARIN SODIUM (UNII: SB8ZUX40TY)			

<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>TRISODIUM CITRATE DIHYDRATE</b> (UNII: B22547B95K)	
<b>SORBITOL</b> (UNII: 506T60A25R)	

### Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	GRAPE (Grape Flavor)	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58657-505-08	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/14/2016	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/14/2016	

**Labeler** - Method Pharmaceuticals (060216698)

Revised: 10/2025

Method Pharmaceuticals