

CHILDRENS ACETAMINOPHEN- acetaminophen suspension

Major Pharmaceuticals

Children's Acetaminophen Oral Suspension

Ibuprofen Free/Alcohol Free/Aspirin Free

Drug Facts

Active ingredient ***(in each 5 mL)***

Acetaminophen 160 mg.....Pain reliever/fever reducer

Uses

temporarily:

- reduces fever
- relieves minor aches and pain due to:
 - the common cold
 - flu
 - headache
 - sore throat
 - toothache

Warnings

Liver warning: This product contains contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly

Do not Use

- with any other drug containing acetaminophen (prescription or nonprescription). If

you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product, do not exceed the recommended dose (see overdose warning)

Stop use and ask a doctor if:

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Keep out of the reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). Quick medical attention is critical for adults as well as children even if you do not notice any sign or symptoms

Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not take more than directed (see overdose warning)**
- **shake well before using**
- mL = milliliter
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL) *
Under 24	Under 2	Ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL
72-95	11 years	15 mL

*or as directed by a doctor

Other information

- **each 5 mL contains:** sodium 2 mg
- Store at 20° to 25°C (68° to 77°F)
- a red, strawberry flavored suspension supplied in the following oral dosage form:
NDC 0904-7278-41: 5 mL unit dose cup, in a tray of ten cups

Inactive ingredients:

acesulfame K, butylparaben, citric acid, FD&C Red No. 40, flavoring, glycerin, high fructose corn syrup, polysorbate 80, propylene glycol, purified water, sodium benzoate, sorbitol solution, veegum and xanthan gum.

Questions or comments?

Call 1-800-845-8210.

R08/22

Distributed by:

MAJOR® PHARMACEUTICALS

Indianapolis, IN 46268

PRINCIPAL DISPLAY PANEL - 5 mL Unit Dose Cup

MAJOR®

0904-7278-41

Children's Acetaminophen Oral Suspension

160 mg per 5 mL

Pain Reliever • Fever Reducer

Ibuprofen Free / Alcohol Free / Aspirin Free

SHAKE WELL

Delivers 5 mL • See insert

For Institutional Use Only

MAJOR® PHARMACEUTICALS

Indianapolis, IN 46268

F1781C050822



CHILDRENS ACETAMINOPHEN

acetaminophen suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0904-7278(NDC:0121-1781)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
MAGNESIUM ALUMINUM SILICATE (UNII: 6M3P64V0NC)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
BUTYLPARABEN (UNII: 3QPI1U3FV8)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	red	Score	
Shape		Size	
Flavor	BERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0904-7278-70	10 in 1 CASE	06/02/2023	
1		10 in 1 TRAY		
1	NDC:0904-7278-41	5 mL in 1 CUP, UNIT-DOSE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	06/02/2023	

Labeler - Major Pharmaceuticals (191427277)

Revised: 1/2025

Major Pharmaceuticals