QUIT SMOKING NICOTINE PATCHES- nicotine transdermal patches patch Shenzhen Paxi Trading Co., Ltd.

85835-001

Active Ingredient

Nicotine 28mg

Purpose

Stopsmoking aid

Use

Apply one new patch every 24 hours on skin that is dry, clean and hairless.

Wash hands after applying or removing patch. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.

The used patch should be removed and a new one applied to a different skin site at the same time each day.

If you have vivid dreams, you may remove the patch at bedtime and apply a new one in the morning.

Do not wear more than one patch at a time.

Do not cut patch in half or into smaller pieces.

Do not leave patch on for more than 24 hours because it may irritate your skin and loses strength after 24 hours.

To avoid possible burns, remove patch before undergoing any MRI (magnetic resonance imaging) procedures.

It is important to complete treatment. If you feel you need to use the patch for a longer period to keep from smoking, talk to your healthcare provider.

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider.

Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Do not use

Ask a doctor before use if you have

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.

- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.

- an allergy to adhesive tape or have skin problems, because you are more likely to get rashes.

- stomach ulcer or diabetes.
- history of seizures.

When Using

When using this product

- if you have vivid dreams or other sleep disturbance are removed this patch at bedtime.

Stop Use

Stop use and ask a doctor if

- skin redness caused by the patch does not go away after four days, or if your skin swells, or you get a rash.

- irregular heartbeat or palpitations occur.

- you get symptoms of nicotine overdose, such as persistent nausea, vomiting, dizziness, weakness and rapid heartbeat.

- you have symptoms of an allergic reaction (such as difficulty breathing or rash).

Ask Doctor

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug.

- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Keep Out Of Reach Of Children

Keep out of reach of children and pets

Used patches have enough nicotine to poison children and pets. If swallowed, get medical help or contact a Poison Control Center right away. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.

Directions

If you are under 18 years of age, ask a doctor before use.

Before using this product, read the enclosed self-help guide for complete directions and other information.

Begin using the patch on your quit day.

If you smoke more than 20 cigarettes per day, use the following schedule below:

Weeks 1 and 2 STEP 1+ use1¹/₂ patch/day

Weeks 3 and 6

STEP 1 use 1 patch/day Weeks 7 and 8 STEP 2 use I/2 patch/day Weeks 9 and 10 STEP 3 use I/4 patch/day

If you smoke more than 10 but not more than 20 cigarettes, start with Step 1, if you smoke 10 or less cigarettes per day, start with Step 2 for 4 weeks (Start with 1 patch or 3/4 patch for the first week), then on to Step 3 for 2 weeks. (Read the enclosed self-help guide for additional information.)

Apply one new patch every 24 hours on skin that is dry, clean and hairless.

Wash hands after applying or removing patch. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch. The used patch should be removed and a new one applied to a different skin site at the

same time each day.

If you have vivid dreams, you may remove the patch at bedtime and apply a new one in the morning.

Do not wear more than one patch at a time.

Do not cut patch in half or into smaller pieces.

Do not leave patch on for more than 24 hours because it may irritate your skin and loses strength after 24 hours.

To avoid possible burns, remove patch before undergoing any MRI (magnetic resonance imaging) procedures.

It is important to complete treatment. If you feel you need to use the patch for a longer period to keep from smoking, talk to your healthcare provider.

Other information

Store at room temperature

Inactive ingredients

Acrylate adhesive, aluminized polyester, cellulose paper, methacrylic acid copolymer

Questions

After-salesservice contactinformation. If you have any doubts with our products, please contact us. Paxitrading@outlook.com

PRINCIPAL DISPLAY PANEL



QUIT SMOKING NICOTINE PATCHES

nicotine transdermal patches patch

Pr	oduct Inforn	nation						
Pro	oduct Type		HUMAN OTC DRUG	Item Coo	em Code (Source) ND		DC:85835-001	
Ro	ute of Adminis	tration	TOPICAL					
Ac	tive Ingredie	ent/Active	Moiety					
		Ingredi	ent Name		Basis of Streng	gth	Strength	
VIC	OTINE (UNII: 6M3	3C89ZY6R) (NI	COTINE - UNII:6M3C89ZY6R)	NICOTINE		28 mg in 100	
na	active Ingred	lients						
			Ingredient Name	2			Strengt	
	RYLIC ACID (UNII:	(94PBK7X85)						
		. je .: <u>2</u> ,,						
	LYESTER-7 (UNII:	-						
PO		0841698D2F)						
PO PO	LYESTER-7 (UNII: WDERED CELLUI	0841698D2F) L OSE (UNII: SI		1ER (1:1) ((UNII: 74G4R6TH13)			
PO PO	LYESTER-7 (UNII: WDERED CELLUI	0841698D2F) L OSE (UNII: SI	MD1X3XO9M)	1ER (1:1) ((UNII: 74G4R6TH13)			
PO PO	LYESTER-7 (UNII: WDERED CELLUI	0841698D2F) L OSE (UNII: SI	MD1X3XO9M)	1ER (1:1) ((UNII: 74G4R6TH13)			
PO PO ME	LYESTER-7 (UNII: WDERED CELLUI	0841698D2F) L OSE (UNII: SI	MD1X3XO9M)	1ER (1:1) ((UNII: 74G4R6TH13)			
PO PO ME	LYESTER-7 (UNII: WDERED CELLUI THACRYLIC ACIE	0841698D2F) LOSE (UNII: SI D - METHYL M	MD1X3XO9M)		(UNII: 74G4R6TH13) arketing Start Date	Mar	rketing End Date	
PO PO ME Pa	LYESTER-7 (UNII: WDERED CELLUI THACRYLIC ACID Ckaging Item Code NDC:85835-001-	0841698D2F) LOSE (UNII: SI D - METHYL M Pac	MD1X3XO9M) IETHACRYLATE COPOLYM	Ma	arketing Start	Ma		
	LYESTER-7 (UNII: WDERED CELLUI THACRYLIC ACID Ckaging Item Code NDC:85835-001- 01	0841698D2F) LOSE (UNII: SI D - METHYL M Pac 15 in 1 BOX; Product	MD1X3XO9M) IETHACRYLATE COPOLYM Kage Description	Ma 06/30	arketing Start Date	Mar		
	LYESTER-7 (UNII: WDERED CELLUI THACRYLIC ACID Ckaging Item Code NDC:85835-001- 01 NDC:85835-001-	0841698D2F) LOSE (UNII: SI D - METHYL M Pac 15 in 1 BOX; Product 30 in 1 BOX;	MD1X3XO9M) IETHACRYLATE COPOLYM Kage Description Type 0: Not a Combination	Ma 06/30	arketing Start Date 0/2025	Ma		
	LYESTER-7 (UNII: WDERED CELLUI THACRYLIC ACID Ckaging Item Code NDC:85835-001- 01 NDC:85835-001-	0841698D2F) LOSE (UNII: SI D - METHYL M Pac 15 in 1 BOX; Product 30 in 1 BOX;	MD1X3XO9M) IETHACRYLATE COPOLYM Kage Description Type 0: Not a Combination	Ma 06/30	arketing Start Date 0/2025	Ma		
Pa #	LYESTER-7 (UNII: WDERED CELLUI THACRYLIC ACID Ckaging Item Code NDC:85835-001- 01 NDC:85835-001-	0841698D2F) LOSE (UNII: SI D - METHYL M Pac 15 in 1 BOX; Product 30 in 1 BOX; Product	MD1X3XO9M) IETHACRYLATE COPOLYM Kage Description Type 0: Not a Combination Type 0: Not a Combination	Ma 06/30	arketing Start Date 0/2025	Ma		
	LYESTER-7 (UNII: WDERED CELLUI THACRYLIC ACID ckaging Item Code NDC:85835-001- 01 NDC:85835-001- 02	0841698D2F) LOSE (UNII: SI D - METHYL M Pac 15 in 1 BOX; " Product 30 in 1 BOX; " Product	MD1X3XO9M) IETHACRYLATE COPOLYM Kage Description Type 0: Not a Combination Type 0: Not a Combination	Ma 06/30 06/30	arketing Start Date 0/2025			

Labeler - Shenzhen Paxi Trading Co., Ltd. (633886766)

Establishment								
Name	Address	ID/FEI	Business Operations					
Shenzhen Paxi Trading Co., Ltd.		633886766	manufacture(85835-001)					

Revised: 6/2025