

**NIGHTTIME COLD AND FLU MULTI SYMPTOM RELIEF- acetaminophen,  
dextromethorphan hbr, doxylamine succinate liquid  
Shield Pharmaceuticals Corp**

---

**Nighttime Cold and Flu Multi-Symptom Relief**

**Drug Facts**

**Active ingredient (in each softgel)**

Acetaminophen 650 mg  
Dextromethorphan HBr 30 mg  
Doxylamine succinate 12.5 mg

**Purpose**

Pain reliever/fever reducer  
Cough suppressant  
Antihistamine

**Uses**

temporarily relieves common cold/flu symptoms:

- cough due to minor throat & bronchial irritation
- sore throat
- headache • minor aches & pains • fever • runny nose & sneezing

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 Softgels in 24 hrs, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Allergy Alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to enlarged prostate gland

**Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

**When using this product**

- do not use more than directed n excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, & tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults & for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

---

adults & children 12 yrs & over

30 mL every 6 hrs

children 4 to under 12 yrs

ask a doctor

children under 4 yrs

do not use

---

**Other information**

- **each 30 mL contains:** sodium 24 mg
- store between 20-25°C (68-77°F) and do not refrigerate.

**Inactive ingredients**

alcohol, citric acid, FD&C Blue No. 1, FD&C Red No. 40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose,

xanthan gum

## Questions or comments?

1-800-373-6981(toll-free)

Distributed by:

**Shield Pharmaceuticals Corp.**


Ronkonkoma, NY 11779

## PRINCIPAL DISPLAY PANEL

ValuRx

Nighttime Cold & Flu  
Multi-Symptom Relief

Label



### Drug Facts (continued)

**Sore throat warning:** If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, see a doctor promptly.

#### Do not use

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have** ■ liver disease ■ glaucoma ■ cough that occurs with too much phlegm (mucus) ■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema ■ trouble urinating due to enlarged prostate gland

**When using this product** ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery ■ alcohol, sedatives, and tranquilizers may increase drowsiness.

**Stop use and ask a doctor if** ■ pain or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults & for children even if you do not notice any signs or symptoms.

#### Directions

|                                 |                   |
|---------------------------------|-------------------|
| adults & children 12 yrs & over | 30 mL every 6 hrs |
| children 4 to under 12 yrs      | ask a doctor      |
| children under 4 yrs            | do not use        |

#### Other information

■ each 30 mL contains: sodium 24 mg  
■ store between 20-25°C (68-77°F) and do not refrigerate.  
**Inactive ingredients** alcohol, citric acid, FD&C Blue No. 1, FD&C Red No. 40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucrose, xanthan gum

## Questions or comments?

1-800-373-6981

098 06/2004



Lot#: EXP:..



\* This product is not manufactured or distributed by the Procter & Gamble Company, owner of the registered trademarks Vicks® NyQuil® Cold & Flu.  
Made in India

**ValuRx**Compare to the  
Active Ingredients in  
Vicks® NyQuil® Cold & Flu\*

# Nighttime \$125 Cold & Flu

## Multi-Symptom Relief

**Acetaminophen-**  
Pain reliever/Fever reducer**Dextromethorphan HBr-**  
Cough suppressant**Doxylamine succinate-**  
Antihistamine

4 fl. oz. (118 mL)

**TAMPER EVIDENT: DO NOT USE IF SAFETY SEAL AROUND THE BOTTLE OR  
UNDER THE CAP IS BROKEN OR MISSING****Drug Facts****Active Ingredients (in each 15 mL)**Acetaminophen 650 mg ..... Pain reliever/fever reducer  
Dextromethorphan HBr 30 mg ..... Cough suppressant  
Doxylamine succinate 12.5 mg ..... Antihistamine**Uses**

temporarily relieves common cold/flu symptoms: ■ cough due to minor throat &amp; bronchial irritation ■ sore throat ■ minor aches &amp; pains ■ fever ■ runny nose and sneezing

**Warnings****Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4 doses in 24 hours, which is the maximum daily amount for this product ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product**Allergy Alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away.**Drug Facts (continued under label)**

\* This product is not manufactured or distributed by The Procter &amp; Gamble Company, owner of the registered trademarks Vicks®, NyQuil® Cold &amp; Flu.



Made in India



ORIG 06/2024

Distributed by:  
Shield Pharmaceuticals Corp.  
Hauppauge, NY 11788LOT:  
EXP.:

## NIGHTTIME COLD AND FLU MULTI SYMPTOM RELIEF

acetaminophen, dextromethorphan hbr, doxylamine succinate liquid

**Product Information**

|                                |                |                           |                |
|--------------------------------|----------------|---------------------------|----------------|
| <b>Product Type</b>            | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:83059-0125 |
| <b>Route of Administration</b> | ORAL           |                           |                |

**Active Ingredient/Active Moiety**

| Ingredient Name  | Basis of Strength             | Strength             |
|--|-------------------------------|----------------------|
| <b>ACETAMINOPHEN</b> (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)                    | ACETAMINOPHEN                 | 650 mg<br>in 118 mL  |
| <b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 30 mg<br>in 118 mL   |
| <b>DOXYLAMINE SUCCINATE</b> (UNII: V9B19B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)                | DOXYLAMINE SUCCINATE          | 12.5 mg<br>in 118 mL |

**Inactive Ingredients**

| Ingredient Name                               | Strength |
|---|----------|
| <b>CITRIC ACID</b> (UNII: 2968PHW8QP)         |          |
| <b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD) |          |
| <b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA) |          |
| <b>GLYCERIN</b> (UNII: PDC6A3C00X)            |          |
| <b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)    |          |
| <b>WATER</b> (UNII: 059QF0KO0R)               |          |
| <b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)    |          |
| <b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)     |          |
| <b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)     |          |

**SORBITOL** (UNII: 506T60A25R)

**SUCRALOSE** (UNII: 96K6UQ3ZD4)

**XANTHAN GUM** (UNII: TTV12P4NEE)

### Product Characteristics

|                 |        |                     |  |
|-----------------|--------|---------------------|--|
| <b>Color</b>    | green  | <b>Score</b>        |  |
| <b>Shape</b>    |        | <b>Size</b>         |  |
| <b>Flavor</b>   | CHERRY | <b>Imprint Code</b> |  |
| <b>Contains</b> |        |                     |  |

### Packaging

| # | Item Code        | Package Description                                   | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:83059-0125-4 | 118 mL in 1 BOTTLE; Type 0: Not a Combination Product | 01/24/2026           |                    |

### Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M012                                     | 01/24/2026           |                    |

**Labeler** - Shield Pharmaceuticals Corp (118724924)

**Registrant** - Shield Pharmaceuticals Corp (118724924)

Revised: 1/2026

Shield Pharmaceuticals Corp