

GUAIFED- guaifenesin liquid
Aldama Pharmaceuticals, Inc

Guaifed

Active Ingredients:(per 10ml= 2 teaspoonfuls) **Purpose**

Guaifenesin 200 mg Expectorant

Purpose

Expectorant

Uses:

- Helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive.

Warnings

- **Do not use: if you ever had an allergic reaction to any of the ingredients in this product**

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- cough that last or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Stop use and ask a doctor if

- cough lasts more than 7 days, comes back or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding,ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take every 4 hours as needed, or as directed by doctor
- do not take more than 6 doses in 24 hours
- do not exceed recommended dose

Age	Dose
adults and children 12 years and over	10 mL to 20 mL (2 to 4 teaspoonfuls)
children under 12 years of age	ask a doctor

Other information

- **TAMPER - EVIDENT FEATURE:** Do not use if foil seal over bottle opening is torn, broken or missing.
- each 10 mL contains: Sodium 20 mg
- store at room temperature 15° - 30°C (59° - 86°F).

- Protect from freezing
- Protect from light

Inactive Ingredients:

Artificial and natural cherry flavor, citric acid, menthol, Methylparaben, Propylene glycol, Propylparaben, purified water, sodium citrate, sucralose

Questions or Comments? 305-592-9216 or info@aldamapharm.com

DRUG FACTS	
Active ingredient (per 10mL = 2 teaspoonfuls)	Purpose
Guaifenesin, USP 200mg.....	Expectorant
Uses • helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive	
Warnings	
Do not use • if you ever had an allergic reaction to any of the ingredients in this product	
Ask a doctor before use if you have • cough that occurs with too much phlegm (mucus) • cough that last or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema.	
Stop use and ask a doctor if • cough lasts more than 7 days, comes back or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.	
If pregnant or breast-feeding , ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. A	

NDC 73564-801-04

GUAIFED

GUAIFENESIN ORAL SOLUTION

EXPECTORANT



LOOSENS AND RELIEFS CHEST CONGESTION

Alcohol-Dye-Sugar Free

FOR ADULTS AND CHILDREN OVER 12 YEARS

COMPARE TO ACTIVE INGREDIENT OF ORIGINAL ROBITUSSIN®

4 fl.oz (118mL)

DRUG FACTS	
<i>continued</i>	
Directions • take every 4 hours, as needed, or as directed by a doctor • do not take more than 6 doses in 24 hours • do not exceed recommended dose	
adults and children 12 years and over	10 to 20mL (2 to 4 teaspoonfuls)
children under 12 years	ask a doctor
Other information • TAMPER-EVIDENT: Do not use if foil seal over bottle opening is torn, broken, or missing • each 10mL contains Sodium 20mg • store at room temperature 15°-30°C (59°-86°F) • protect from freezing • protect from light	
Inactive Ingredients: Artificial and natural cherry flavor, citric acid, menthol, methylparaben, propylene glycol, propylparaben, purified water, sodium citrate, sucralose	
Questions or Comments? (305)592-9216 or info@aldamapharm.com	
*This product is not manufactured or distributed by the owner of the registered trademark Robitussin®	
 Miami, FL 33122 USA	Code #L-9 Rev: 05/25 Lot#/Exp. Date
	

DRUG FACTS	
Active ingredient (per 10mL = 2 teaspoonfuls)	Purpose
Guaifenesin, USP 200mg.....	Expectorant
Uses • helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive	
Warnings	
Do not use • if you ever had an allergic reaction to any of the ingredients in this product	
Ask a doctor before use if you have • cough that occurs with too much phlegm (mucus) • cough that last or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema.	
Stop use and ask a doctor if • cough lasts more than 7 days, comes back or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.	
If pregnant or breast-feeding , ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. A	

NDC 73564-801-08

GUAIFED

GUAIFENESIN ORAL SOLUTION

EXPECTORANT

LOOSENS AND RELIEFS CHEST CONGESTION

Alcohol-Dye-Sugar Free

FOR ADULTS AND CHILDREN OVER 12 YEARS

COMPARE TO ACTIVE INGREDIENT OF ORIGINAL ROBITUSSIN®

8 fl.oz (237mL)

DRUG FACTS	
<i>continued</i>	
Directions • take every 4 hours, as needed, or as directed by a doctor • do not take more than 6 doses in 24 hours • do not exceed recommended dose	
adults and children 12 years and over	10 to 20 mL (2 to 4 teaspoonfuls)
children under 12 years	ask a doctor
Other information • TAMPER-EVIDENT: Do not use if foil seal over bottle opening is torn, broken, or missing • Each 10mL contains Sodium 20mg • store at room temperature 15°-30°C (59°-86°F) • protect from freezing • protect from light	
Inactive Ingredients: Artificial and natural cherry flavor, citric acid, menthol, methylparaben, propylene glycol, propylparaben, purified water, sodium citrate, sucralose	
Questions or Comments? (305)592-9216 or info@aldamapharm.com	
*This product is not manufactured or distributed by the owner of the registered trademark Robitussin®	
Code #L-11 Rev: 05/25	Lot#/Exp. Date
 Miami, FL 33122 USA	

DRUG FACTS	
Active ingredient (per 10mL= 2 teaspoonfuls)	Purpose
Guaifenesin, USP 200mg	Expectorant
Uses • helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive	
Warnings	
Do not use • if you ever had an allergic reaction to any of the ingredients in this product	
Ask a doctor before use if you have • cough that occurs with too much phlegm (mucus) • cough that last or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema.	
Stop use and ask a doctor if • cough lasts more than 7 days, comes back or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.	
If pregnant or breast-feeding , ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	

NDC 73564-801-16

GUAIFED

GUAIFENESIN ORAL SOLUTION

EXPECTORANT

LOOSENS AND RELIEFS CHEST CONGESTION

Alcohol-Dye-Sugar Free
FOR ADULTS AND CHILDREN OVER 12 YEARS

COMPARE TO ACTIVE INGREDIENT OR ORIGINAL ROBITUSSIN®

16 fl.oz (473mL)

DRUG FACTS <i>continued</i>	
Directions	
<ul style="list-style-type: none"> • take every 4 hours, as needed, or as directed by a doctor • do not take more than 6 doses in 24 hours • do not exceed recommended dose 	
adults and children 12 years and over	10 to 20 mL (2 to 4 teaspoonfuls)
children under 12 years	ask a doctor
Other information	
<ul style="list-style-type: none"> • each 10 mL contains: sodium 20 mg • TAMPER EVIDENT: Do not use if foil seal over bottle opening is torn, broken or missing. • store at room temperature between 15°-30°C (59°-86°F) • protect from freezing • protect from light 	
Inactive ingredients artificial and natural cherry flavor, citric acid, menthol, methylparaben, propylene glycol, propylparaben, purified water, sodium citrate, sucralose	
Questions or Comments? (305)592-9216 or info@aldamapharm.com	

*This product is not manufactured or distributed by the owner of the registered trademark Robitussin®

THIS IS A BULK CONTAINER NOT INTENDED FOR DISPENSING

Code #L-13
Rev: 05/25

Lot#/Exp. Date



GUAIFED

guaifenesin liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:73564-801
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
SUCRALOSE (UNII: 96K6UQ3ZD4)	
MENTHOL (UNII: L7T10EIP3A)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
CITRIC ACID (UNII: 2968PHW8QP)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73564-801-16	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	07/01/2025	
2	NDC:73564-801-08	237 mL in 1 BOTTLE; Type 0: Not a Combination Product	07/01/2025	
3	NDC:73564-801-04	118 mL in 1 BOTTLE; Type 0: Not a Combination Product	07/01/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/01/2025	

Labeler - Aldama Pharmaceuticals, Inc (119484030)

Revised: 10/2025

Aldama Pharmaceuticals, Inc