

VICKS NYQUIL SEVERE COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, and phenylephrine hydrochloride solution
The Procter & Gamble Manufacturing Company

Vicks[®] NyQuil[™] Severe Cold & Flu Original Liquid

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 650 mg
Dextromethorphan HBr 20 mg
Doxylamine succinate 12.5 mg
Phenylephrine HCl 10 mg

Purpose

Pain reliever/fever reducer
Cough suppressant
Antihistamine
Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- cough to help you sleep
- minor aches & pains
- headache
- fever
- sore throat
- runny nose & sneezing
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

Warnings

Liver warning: This product contains acetaminophen.

Severe liver damage may occur if you take

- more than 4 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions.

Symptoms may include: • skin reddening • blisters • rash
If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days

- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- **each 30 mL contains:** sodium 81 mg
- store at no greater than 25°C and do not refrigerate

Inactive ingredients

alcohol, citric acid, D&C Yellow No. 10, FD&C Yellow No. 6, FD&C Green No. 3, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose

Questions?

1-800-362-1683

TAMPER EVIDENT: Do not use if printed shrinkband is broken or missing.

**DIST. BY PROCTER & GAMBLE,
CINCINNATI OH 45202**

PRINCIPAL DISPLAY PANEL

VICKS®

NyQuil™

SEVERE

COLD & FLU




Acetaminophen, Phenylephrine HCl, Doxylamine Succinate, Dextromethorphan HBr

- Headache, Fever, Sore Throat, Minor Aches & Pains
- Nasal Congestion, Sinus Pressure
- Sneezing, Runny Nose
- Cough

Nighttime Relief

Alcohol 10%

12 FL OZ (354 ml)



VICKS NYQUIL SEVERE COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, and phenylephrine hydrochloride solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37000-815
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	green	Score	
Shape		Size	
Flavor	ANISE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37000-815-08	236 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/21/2014	
2	NDC:37000-815-12	354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/21/2014	
3	NDC:37000-815-36	3 in 1 PACKAGE, COMBINATION	07/21/2014	
3	NDC:37000-815-12	354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End
-----------	---------------------------------	-----------------	---------------

Category	Citation	Date	Date
OTC Monograph Drug	M012	07/21/2014	

Labeler - The Procter & Gamble Manufacturing Company (004238200)

Revised: 7/2024

The Procter & Gamble Manufacturing Company