

**CHILDRENS ACETAMINOPHEN ORAL LIQUID GRAPE FLAVOR-**  
**acetaminophen liquid**  
**AARNA USA INC.**

-----  
**Children's Acetaminophen Oral Liquid**  
**Grape Flavor 160mg per 5mL**

**Pain Reliever**  
**Alcohol Free**  
**Aspirin Free**

**Active Ingredient**

<b>Active ingredient</b> (in each 5mL)	<b>Purpose</b>
Acetaminophen 160 mg .....	Pain reliever/fever reducer

**Uses**

temporarily:

- reduces fever

relieves minor aches and pains due to:

- the common cold
- flu
- headache
- sore throat
- toothache

**Warnings**

**Liver Warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:**Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:**

if sore throat is severe, persist or more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription).

- If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or anyof the inactive ingredients in this product.

**Ask a doctor or pharmacist before use if the child has**  
liver disease.

Ask doctor or pharmacist before use if your child is  
taking the blood thinning drug warfarin. **When using this product do not exceed recommended dose (see overdose warning)**

**Stop use and ask a doctor if**

- pain gets worse or lasts for more then 5 days
- fever gets worse or last for more then 3 days
- new symptoms occur redness or swelling is present. These could be signs of a serious condition.

Overdose warning:

In case of overdose, get medical help to contact Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

**Keep out of Reach of children**

**Directions**

- this product does not contain directions or complete warnings for adult use
- shake well before using
- mL = milliliter
- find right dose on chart below. if possible, use weight to dose, otherwise use age
- use only the enclosed dosing cup designed for use with product.
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours.

Weight (lbs)	Age (yrs.)	Dose (tsp or mL)	Weight (lbs)	Age (yrs.)	Dose (tsp or mL)
under 24	under 2	ask a doctor	48 to 59	6 to 8	2 tsp or 10 mL
25 to 35	2 to 3	1 tsp or 5 mL	60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL	72 to 95	11	3 tsp or 15 mL

**Other Information**

- Store at room temperature 20°-25°C (68°-77°F). Protect from freezing. Protect from light.

Anhydrous Citric Acid, D&C Red No 33, FD&C Blue No 1, Glycerin, Grape Flavor, Polyethylene Glycol 400, Purified Water, Sodium Benzoate, Sodium Citrate, Sorbitol Solution, Sucralose.

call toll-free 1-877-225-6999

**AARNA USA INC.**

**Leland, NC-28451**

**This product is not manufactured by or distributed by McNeil Consumer Healthcare, owner of the registered trademark Tylenol Elixir.**

NDC: 82 568-0009-4

# Children's Acetaminophen Oral Liquid


160mg per 5mL

## PAIN RELIEVER and fever Reducer

Alcohol Free  
Aspirin Free

For Ages  
2 to 11 Years

**Grape Flavor**



**Drug Facts**

**Active ingredient (in each 5 mL):**  
Acetaminophen 160 mg

**Purpose:**  
Pain reliever / fever reducer

**Uses temporarily:** reduces fever • relieves minor aches and pains due to: • the common cold • flu • headaches • sore throat • toothache

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:  
• more than 5 doses in 24 hours, which is the maximum daily amount with other liquids containing acetaminophen. Always alert. Acetaminophen may cause severe skin reactions. Symptoms may include:  
• skin redness • blisters • rash. If a skin reaction occurs, stop use and seek medical help right away.  
**Severe threat warning:** If one throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. Do not use with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • If your child is allergic to acetaminophen or any of the inactive ingredients in this product.

**Ask a doctor before use if your child has a liver disease.**

**Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.**

**When using this product, do not exceed recommended dose (see overdose warning).**

**Stop use and ask a doctor if pain gets worse or lasts for more than 5 days • if fever gets worse or lasts for more than 3 days • if new symptoms occur • if rash or swelling is present. These could be signs of a serious condition. Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

**Directions:** This product does not contain directions or complete warnings for adult use • always follow below using • mL • milliliter • and right dose on chart below. If possible, use weight to dose; otherwise, use age. • Use only the enclosed dosing cup designed for use with product.

Weight (lb.)	Age (yrs.)	Dose ( tsp or mL )*	Weight (lb.)	Age (yrs.)	Dose ( tsp or mL )*
under 24	under 2	ask a doctor	48 to 59	6 to 8	2 tsp or 10 mL
24 to 35	2 to 3	1 tsp or 5 mL	60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL	72 to 95	11	3 tsp or 15 mL

\*or as directed by a doctor

**Other Information:** Store at room temperature 20°-25°C (68°-77°F). Product from Freezing & Light.

**Inactive Ingredients:** Anytolene Citric Acid, D-60 Red No. 33, FD&C Blue No. 1, Glycerin, Grape Flavour, Polyethylene Glycol 400, Purified Water, Sodium Benzoate, Sodium Citrate, Sorbitol Solution, Sucralose.

**Questions or comments?** call toll-free: 1-877-225-6999.

\*This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark "Mylan Effect".

**Sample Evidence: Do Not Use If Safety Seal Under Cap is Broken or Missing**

**Aarna USA**  
 Ma not actare to r:  
 AARNA USA Inc.  
 Leeland, NC-28451  
 (825) 688-0009/4-5

**Drug Facts**

**Active ingredient (in each 5 mL)** **Purpose**  
Acetaminophen 160 mg Pain reliever / fever reducer

**Uses** temporarily: ■ reduces fever ■ relieves minor aches and pains due to: ■ the common cold ■ flu ■ headache ■ sore throat ■ toothache

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:  
■ more than 5 doses in 24 hours, which is the maximum daily amount  
■ with other drugs containing acetaminophen.  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.  
**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.  
**Do not use** ■ with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If your child is allergic to acetaminophen or any of the inactive ingredients in this product.  
**Ask a doctor before use if your child has liver disease.**  
**Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.**  
**When using this product, do not exceed recommended dose (see overdose warning)**  
**Stop use and ask a doctor if** ■ pain gets worse or lasts for more than 5 days ■ fever gets worse or lasts for more than 3 days ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition.  
**Keep out of reach of children.**  
**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.  
**Directions**  
■ this product does not contain directions or complete warnings for adult use, shake well before using  
■ mL = milliliter  
■ find right dose on chart below. If possible, use weight to dose; otherwise, use age.  
■ use only the enclosed dosing cup designed for use with product. Do not use any other dosing device.  
■ if needed, repeat dose every 4 hours while symptoms last

NDC: 82568-0009-4

**Children's Acetaminophen Oral Liquid**

**160mg per 5mL**

**PAIN RELIEVER and fever Reducer**

**Alcohol Free Aspirin Free**

**For Ages 2 to 11 Years**

**Grape Flavor**

**4 FL.OZ (118 mL)**

NDC: 82568-0009-4

**Children's Acetaminophen Oral Liquid**

**160mg per 5mL**

**PAIN RELIEVER and fever Reducer**

**Alcohol Free Aspirin Free**

**For Ages 2 to 11 Years**

**Grape Flavor**

**4 FL.OZ (118 mL)**

**Acetaminophen Oral Liquid**

**Drug Facts (continued)**

■ do not give more than 5 times in 24 hours  
■ do not give more than 5 days unless directed by doctor  
■ do not give more than directed (see overdose warning)

Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)*
under 24	under 2	ask a doctor
24 to 35	2 to 3	1 tsp or 5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL
48 to 59	6 to 8	2 tsp or 10 mL
60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
72 to 95	11	3 tsp or 15 mL

\*or as directed by a doctor

**Other Information**  
store at room temperature 20°-25°C (68°-77°F). Protect from Freezing. Protect from Light.

**Inactive Ingredients** Anhydrous Citric Acid, D&C Red No 33, FD&C Blue No 1, Glycerin, Grape Flavor, Polyethylene Glycol 400, Purified Water, Sodium Benzoate, Sodium Citrate, Sorbitol Solution, Sucralose.

**Questions or comments?**  
call toll-free 1-877-225-6999.

\*\*This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark Tylenol® Elkin®.

**Tamper Evident: Do Not Use if Safety Seal Under Cap is Broken or Missing**

Exclusively distributed by: Aarna Pharma Inc. Fairfield, NJ-07004

**Aarna USA**  
Manufactured for:  
AARNA USA Inc.  
Leland, NC-28451

N 3 82568 00094 5

NDC: 82568-0009-8

**Children's Acetaminophen Oral Liquid**

**160mg per 5mL**

**PAIN RELIEVER and fever Reducer**

**Alcohol Free Aspirin Free**

**For Ages 2 to 11 Years**

**Grape Flavor**

**8 FL.OZ (236 mL)**

**Drug Facts**

**Active ingredient (in each 5 mL)** **Purpose**  
Acetaminophen 160 mg Pain reliever / fever reducer

**Uses** temporarily: ■ reduces fever ■ relieves minor aches and pains due to: ■ the common cold ■ flu ■ headache ■ sore throat ■ toothache

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:  
■ more than 5 doses in 24 hours, which is the maximum daily amount  
■ with other drugs containing acetaminophen.  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. Do not use ■ with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If your child is allergic to acetaminophen or any of the inactive ingredients in this product.  
**Ask a doctor before use if your child has liver disease.**  
**Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.**  
**When using this product, do not exceed recommended dose (see overdose warning)**  
**Stop use and ask a doctor if** ■ pain gets worse or lasts for more than 5 days ■ fever gets worse or lasts for more than 3 days ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition.  
**Keep out of reach of children.**  
**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.  
**Directions**  
■ this product does not contain directions or complete warnings for adult use, shake well before using  
■ mL = milliliter  
■ find right dose on chart below. If possible, use weight to dose; otherwise, use age.  
■ use only the enclosed dosing cup designed for use with product.

**Drug Facts (continued)**

Do not use any other dosing device. ■ if needed, repeat dose every 4 hours while symptoms last ■ do not give more than 5 times in 24 hours.

Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)*
under 24	under 2	ask a doctor
24 to 35	2 to 3	1 tsp or 5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL
48 to 59	6 to 8	2 tsp or 10 mL
60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
72 to 95	11	3 tsp or 15 mL

\*or as directed by a doctor

**Other Information**  
store at room temperature 20°-25°C (69°-77°F). Protect from Freezing. Protect from Light.

**Inactive Ingredients** Anhydrous Citric Acid, D&C Red No 33, FD&C Blue No 1, Glycerin, Grape Flavor, Polyethylene Glycol 400, Purified Water, Sodium Benzoate, Sodium Citrate, Sorbitol Solution, Sucralose.

**Questions or comments?** call toll-free 1-877-225-6999.

\*\*This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark Tylenol® Elkin®.

**Tamper Evident: Do Not Use if Safety Seal Under Cap is Broken or Missing**

Exclusively distributed by: Aarna Pharma Inc. Fairfield, NJ-07004

**Aarna USA**  
Manufactured for:  
AARNA USA Inc.  
Leland, NC-28451

N 3 82568 00098 3



### Drug Facts

**Active ingredient (in each 5 mL)** **Purpose**

Acetaminophen 160 mg ..... Pain reliever / fever reducer

**Uses** temporarily: ■ reduces fever ■ relieves minor aches and pains due to: ■ the common cold ■ flu ■ headache ■ sore throat ■ toothache

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes: ■ more than 5 doses in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen.

### Allergy alert:

Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** ■ with any other drug containing acetaminophen (prescription or non prescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If your child is allergic to acetaminophen or any of the inactive ingredients in this product.

**Ask a doctor before use** if your child has liver disease.

**Ask a doctor or pharmacist before use** if your child is taking the blood thinning drug warfarin.

**When using this product, do not exceed recommended dose** (see overdose warning)

**Stop use and ask a doctor if** ■ pain gets worse or lasts for more than 5 days ■ fever gets worse or lasts for more than 3 days. ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

### Directions

■ this product does not contain directions or complete warnings for adult use, shake well before using

■ mL = milliliter

■ find right dose on chart below. If possible, use weight to dose; otherwise, use age.

■ use only the enclosed dosing cup designed for use with product. Do not use any other dosing device.

■ if needed, repeat dose every 4 hours while symptoms last

NDC: 82568-0009-8



## Children's Acetaminophen Oral Liquid

160mg per 5mL

**PAIN RELIEVER**  
and fever Reducer

**Alcohol Free**  
**Aspirin Free**

**For Ages**  
**2 to 11 Years**



Grape  
Flavor

8 FL.OZ (236 mL)

NDC: 82568-0009-8



## Children's Acetaminophen Oral Liquid

160mg per 5mL

**PAIN RELIEVER**  
and fever Reducer

**Alcohol Free**  
**Aspirin Free**

**For Ages**  
**2 to 11 Years**



Grape  
Flavor

8 FL.OZ (236 mL)

## Acetaminophen Oral Liquid

### Drug Facts (continued)

■ do not give more than 5 times in 24 hours  
■ do not give more than 5 days unless directed by doctor  
■ do not give more than directed (see overdose warning)

Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)*
under 24	under 2	ask a doctor
24 to 35	2 to 3	1 tsp or 5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL
48 to 59	6 to 8	2 tsp or 10 mL
60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
72 to 95	11	3 tsp or 15 mL

\*or as directed by a doctor

### Other Information

store at room temperature 20°-25°C (68°-77°F). Protect from Freezing. Protect from Light.

**Inactive Ingredients** Anhydrous Citric Acid, D&C Red No. 33, FD&C Blue No. 1, Glycerin, Grape Flavor, Polyethylene Glycol 400, Purified Water, Sodium Benzoate, Sodium Citrate, Sorbitol Solution, Sucralose.

### Questions or comments?

call toll-free 1-877-225-6999.

\*\*This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark Tylenol® Elixir®

**Tamper Evident: Do Not Use if Safety Seal Under Cap is Broken or Missing**  
Exclusively distributed by: Akron Pharma Inc. Fairfield, NJ-07004



**Aarna USA**  
Manufactured for:  
**AARNA USA Inc.**  
Leland, NC-28451



### Drug Facts

**Active ingredient (in each 5 mL)** **Purpose**

Acetaminophen 160 mg ..... Pain reliever / fever reducer

**Uses** temporarily: ■ reduces fever ■ relieves minor aches and pains due to: ■ the common cold ■ flu ■ headache ■ sore throat ■ toothache

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes: ■ more than 5 doses in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### Do not use

■ with any other drug containing acetaminophen (prescription or non prescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If your child is allergic to acetaminophen or any of the inactive ingredients in this product.

**Ask a doctor before use** if your child has liver disease.

**Ask a doctor or pharmacist before use** if your child is taking the blood thinning drug warfarin.

**When using this product, do not exceed recommended dose** (see overdose warning)

**Stop use and ask a doctor if**

■ pain gets worse or lasts for more than 5 days ■ fever gets worse or lasts for more than 3 days. ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

### Directions

■ this product does not contain directions or complete warnings for adult use, shake well before using ■ mL = milliliter ■ find right dose on chart below.

If possible, use weight to dose; otherwise, use age.

NDC: 82568-0009-6



## Children's Acetaminophen Oral Liquid

160mg per 5mL

**PAIN RELIEVER**  
and fever Reducer

**Alcohol Free**  
**Aspirin Free**

**For Ages**  
**2 to 11 Years**



Grape  
Flavor

16 FL.OZ (473 mL)

### Drug Facts (continued)

■ if needed, repeat dose every 4 hours while symptoms last  
■ do not give more than 5 times in 24 hours  
■ do not give more than 5 days unless directed by doctor.

Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)*
under 24	under 2	ask a doctor
24 to 35	2 to 3	1 tsp or 5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL
48 to 59	6 to 8	2 tsp or 10 mL
60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
72 to 95	11	3 tsp or 15 mL

\*or as directed by a doctor

### Other Information

store at room temperature 20°-25°C (68°-77°F). Protect from Freezing. Protect from Light.

**Inactive Ingredients** Anhydrous Citric Acid, D&C Red No. 33, FD&C Blue No. 1, Glycerin, Grape Flavor, Polyethylene Glycol 400, Purified Water, Sodium Benzoate, Sodium Citrate, Sorbitol Solution, Sucralose.

**Questions or comments?** call toll-free 1-877-225-6999.

\*\*This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark Tylenol® Elixir®

**Tamper Evident: Do Not Use if Safety Seal Under Cap is Broken or Missing**

Exclusively distributed by: Akron Pharma Inc. Fairfield, NJ-07004



**Aarna USA**  
Manufactured for:  
**AARNA USA Inc.**  
Leland, NC-28451



CHILDRENS ACETAMINOPHEN ORAL LIQUID GRAPE FLAVOR

acetaminophen liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:82568-0009
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

Inactive Ingredients	
Ingredient Name	Strength
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics			
Color		Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:82568-0009-4	1 in 1 CARTON	07/01/2024	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:82568-0009-8	1 in 1 CARTON	07/01/2024	
2		236 mL in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:82568-0009-6	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	07/01/2024	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	07/01/2024	

**Labeler -** AARNA USA INC. (118515992)

Revised: 7/2025

AARNA USA INC.