

**ASPIRIN- aspirin tablet, film coated**  
**L.N.K. International, Inc.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Sound Body 44-157**

**Active ingredient (in each tablet)**

Aspirin 325 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever/fever reducer

**Uses**

temporarily relieves:

- minor pain of arthritis
- headache
- muscle pain
- menstrual pain
- toothache
- pain and fever of colds

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- facial swelling
- shock
- hives
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- take more or for a longer time than directed
- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer.

- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if**

- you are taking a diuretic
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma

### **Ask a doctor or pharmacist before use if you are**

taking a prescription drug for

- gout
- diabetes
- arthritis

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away.
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or loss of hearing occurs

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

### ***Directions***

- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours
- children under 12 years: ask a doctor

### ***Other information***

- use by expiration date on package
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

### ***Inactive ingredients***

corn starch, hypromellose, polyethylene glycol, propylene glycol

**Questions or comments?**

**1-800-426-9391**

***Principal Display Panel***

**SOUNDBODY™**

†Compare to the active ingredient in  
Genuine Bayer® Aspirin

NDC 50844-957-29

**Aspirin**

**Pain reliever/Fever reducer  
(NSAID) 325 mg**

ANALGESIC

Lite Coated • Caffeine Free

**150 TABLETS**

†This product is not manufactured or distributed by Bayer HealthCare LLC, owner  
of the registered trademark Genuine Bayer® Aspirin.

50844 REV0915B15729

Distributed by **LNK INTERNATIONAL, INC.**, 60 Arkay Drive,  
Hauppauge, NY 11788 USA V#733000 Item# 022715729UPC

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY  
SEAL UNDER CAP IS BROKEN OR MISSING**

Adhesive Only

STOP PEELING

Questions or comments? 1-800-426-9391

**Inactive ingredients** corn starch, hypromellose, polyethylene glycol, propylene glycol

**Other information** ■ use by expiration date on package ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

■ children under 12 years: ask a doctor

**Directions** ■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222), right away.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor, because it may cause problems in the unborn child or complications during delivery.

■ ringing in the ears or loss of hearing occurs ■ new symptoms occur ■ redness or swelling is present ■ fever gets worse or lasts more than 3 days ■ pain gets worse or lasts more than 10 days ■ allergic reaction occurs. Seek medical help right away. ■ have stomach pain that does not get better ■ vomit blood ■ have bloody or black stools ■ feel faint

**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ arthritis ■ gout ■ diabetes ■ asthma

**Ask a doctor or pharmacist before use if you are taking a prescription drug for** ■ heart disease ■ high blood pressure ■ kidney disease ■ liver disease ■ stomach bleeding ■ you have asthma

**Ask a doctor before use if** ■ you are taking a diuretic ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver

**Do not use** ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ if you have ever had an allergic reaction to this product or any of its ingredients

**Drug Facts (continued)** ■ have had stomach ulcers or bleeding problems ■ have 3 or more alcoholic drinks every day while using this product

■ you have ever had an allergic reaction to this product or any of its ingredients

■ you are taking a diuretic ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver disease, or kidney disease ■ you have asthma



<sup>†</sup>Compare to the active ingredient in Genuine Bayer® Aspirin

NDC 50844-957-29

**Aspirin**

Pain reliever/Fever reducer (NSAID) 325 mg

ANALGESIC  
Lite Coated • Caffeine Free

**150** TABLETS



50844 75129 4

This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Genuine Bayer® Aspirin.

50844 REV0615815729  
Distributed by LNK INTERNATIONAL, INC., 60 Arkay Drive,  
Hauppauge, NY 11788 USA V#733000 ITEM# 022715729UPC

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

**Drug Facts**

**Active ingredient (in each tablet)**

Aspirin 325 mg (NSAID)\* ..... Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

**Uses**

temporarily relieves: ■ minor pain of arthritis ■ headache ■ muscle pain ■ menstrual pain ■ toothache ■ pain and fever of colds

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include: ■ facial swelling ■ shock ■ hives ■ asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older

■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ take more or for a longer time than directed

No print/No varnish  
Lot & Exp date

PEEL HERE FOR MORE DRUG FACTS

Sound Body 44-157

**ASPIRIN**

aspirin tablet, film coated

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:50844-957
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

**Inactive Ingredients**

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	

<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	

### Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	Aspirin;44;157
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50844-957-29	150 in 1 BOTTLE; Type 0: Not a Combination Product	02/10/1996	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part343	02/10/1996	

**Labeler** - L.N.K. International, Inc. (038154464)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	PACK(50844-957)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	MANUFACTURE(50844-957)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	PACK(50844-957)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	PACK(50844-957)