

ANTI DIARRHEAL ANTI GAS- loperamide hydrochloride, simethicone tablet
HyVee Inc

Hy-Vee, Inc. Anti-Diarrheal/Anti-Gas Drug Facts

Active ingredients (in each caplet)

Loperamide HCl 2 mg

Simethicone 125 mg

Purposes

Anti-diarrheal

Anti-gas

Uses

relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly referred to as gas

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Heart alert: Taking more than directed can cause serious heart problems or death

Do not use

- if you have bloody or black stool
- if you have difficulty swallowing

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.
- you have difficulty swallowing the caplet

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- take only on an empty stomach (1 hour before or 2 hours after a meal)
- take with a full (8 oz.) glass of water
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information

- **each caplet contains:** calcium 165 mg, sodium 3 mg
- store between 20° -25° C (68° -77° F). Protect from light.
- **do not use if printed blister unit is broken or torn**

Inactive ingredients

acesulfame potassium, croscarmellose sodium, dibasic calcium phosphate, maltodextrin, microcrystalline cellulose, natural and artificial flavor, propylene glycol, stearic acid

Questions or comments?

1-800-289-8343

Package/Label Principal Display Panel

Hy•Vee®

Compare To Imodium® Multi-Symptom Relief Active Ingredients

Loperamide Hydrochloride and Simethicone Tablets, 2 mg/125 mg

Anti-Diarrheal/Anti-Gas

Multi-Symptom Relief Caplet

actual size

Relieves Symptoms of Diarrhea Plus

- Cramps & Pressure
- Bloating
- Gas

24 CAPLETS*

*Capsule-Shaped Tablets

OPEN OTHER END

OR162 2H C1

Drug Facts
Active ingredients (in each caplet) Loperamide HCl 2 mg, Simethicone HCl 125 mg. **Purposes** Anti-diarrheal, Anti-gas.
Uses Relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly referred to as gas.
Warnings Allergy alert: Do not use if you have ever had rash or other allergic reaction to loperamide HCl.
 Heart alert: Taking more than directed can cause serious heart problems or death.
 Do not use if you have a blood clot or blockage.
 If you have difficulty swallowing.
 Ask a doctor before use if you have:
 ■ a history of abnormal heart rhythm
 ■ liver ■ muscle ■ in the stomach ■ a history of liver disease
 A doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.
 When using this product, dizziness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.
 Stop use and ask a doctor if:
 ■ symptoms get worse
 ■ diarrhea lasts for more than 2 days
 ■ you get abdominal pain or bloating. These may be signs of a serious condition.
 ■ you have difficulty swallowing the caplet

Drug Facts (continued)
Inactive ingredients acacia, magnesium stearate, croscarmellose sodium, dibasic calcium phosphate, methylcellulose, hydroxypropyl methylcellulose, polyethylene glycol, stearic acid.
Questions or comments? 1-800-289-8343
 **This product is not manufactured or distributed by the owner of the registered trademark in outline.

Drug Facts (continued)
 If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (nightway) (1-800-222-1222).
Directions
 ■ Drink plenty of clear fluids to help prevent dehydration caused by diarrhea.
 ■ Take only on an empty stomach (1 hour before or 2 hours after a meal).
 ■ Take with a full (8 oz) glass of water.
 ■ First night dose on a half dose if possible, use weight to dose; otherwise, use age.
 Adults and children:
 1. Take after the first loose stool, but no more than 4 capsules in 24 hours.
 2. Years and over:
 (a) 5-11 years: 1 caplet after the first loose stool, but no more than 3 capsules in 24 hours.
 (b) 5-9 lbs: 1 caplet after the first loose stool, but no more than 2 capsules in 24 hours.
 (c) 9-11 years: 1 caplet after the first loose stool, but no more than 2 capsules in 24 hours.
 Children 2-5 years: ask a doctor (34 to 47 lbs).
 Children under 2 years: do not use (up to 33 lbs).

Other information
 ■ Each caplet contains calcium 155 mg, sodium 3 mg, store between 20°-25°C (68°-77°F). Protect from light.
 ■ Do not use if pain bed blister units are broken or torn.



NDC 42507-699-62

Loperamide Hydrochloride and Simethicone Tablets, 2 mg/125 mg

Anti-Diarrheal/ Anti-Gas

Multi-Symptom Relief

Caplet

Relieves Symptoms of Diarrhea *Plus*

- Cramps & Pressure
- Bloating
- Gas



actual size

24
CAPLETS*

*Capsule-Shaped Tablets


DISTRIBUTED BY HY-VEE, INC.
 5820 WESTTOWN PARKWAY
 WEST DES MOINES, IA 50266 PERA1025
 SATISFACTION GUARANTEED
 OR PURCHASE PRICE REFUNDED
 FOR INFORMATION CALL
 1-800-289-8343
 www.hy-vee.com

smartlabel®
 Scan here for more information or call 1-800-289-8343



GLUTEN FREE

CONVENIENT RECLOSEING TAB





ANTI DIARRHEAL ANTI GAS

loperamide hydrochloride, simethicone tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:42507-699
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	125 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
ANHYDROUS DIBASIC CALCIUM PHOSPHATE (UNII: L11K75P92J)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	WHITE	Score	2 pieces
Shape	OVAL	Size	17mm
Flavor		Imprint Code	L0R1;2;125
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:42507-699-62	24 in 1 CARTON	02/25/2026	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:42507-699-53	12 in 1 CARTON	02/25/2026	
3		1 in 1 BLISTER PACK; Type 0: Not a Combination		

4

Product

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA209837	02/25/2026	

Labeler - HyVee Inc (006925671)

Revised: 3/2026

HyVee Inc