

VITASTEM FIRST AID ANTIBIOTIC- tetracycline hydrochloride ointment
VIADERMA DISTRIBUTION INC

Vitastem First Aid Antibiotic

Drug Facts

Active ingredient (in each gram)

Tetracycline HCl 30 mg

Purpose

First aid antibiotic

Use

First aid to help prevent skin infection in minor cuts, scrapes, and burns

Warnings

For external use only.

Do not use

- longer than 1 week unless directed by a doctor
- in the eyes
- over large areas of the body

Ask a doctor before use if you have

deep or puncture wounds, animal bites, or serious burns.

Stop use and ask a doctor if

condition persists or gets worse.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- clean the affected area
- apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
- may be covered with a sterile bandage

Other information

- store between 68-77°F (20-25°C)
- may stain cloth

Inactive ingredients

Acetic acid, ascorbic acid, chlorhexidine gluconate, cholecalciferol, dimethyl sulfoxide, dipropylene glycol, glucono delta lactone, glycerin, histidine, hydroxethyl cellulose, magnesium stearate, methylparaben, sodium hydroxide, sorbic acid, water

Questions or comments?

Call 1-800-585-8685 or visit: www.viadermainc.com

Package Labeling:

Drug Facts

Active ingredient (in each gram) Purpose
Tetracycline HCl 30 mg First aid antibiotic

Use

First aid to help prevent skin infection in minor cuts, scrapes, and burns

Warnings

For external use only.

Do not use

- longer than 1 week unless directed by a doctor
- in the eyes
- over large areas of the body

Ask a doctor before use if you have deep or puncture wounds, animal bites, or serious burns.

Stop use and ask a doctor if condition persists or gets worse. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- clean the affected area
- apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
- may be covered with a sterile bandage

Drug Facts

Other information

- store between 68-77°F (20-25°C)
- may stain cloth

Inactive ingredients

Acetic acid, ascorbic acid, chlorhexidine gluconate, cholecalciferol, dimethyl sulfoxide, dipropylene glycol, glucono delta lactone, glycerin, histidine, hydroxyethyl-cellulose, magnesium stearate, methyparaben, sodium hydroxide, sorbic acid, water

Questions or comments?

Call 1-800-585-8685 or visit:
www.viadermainc.com

LICENSED TO: VIADERMA DISTRIBUTION INC
BY: VIADERMA, INC.

DISTRIBUTED BY: VIADERMA DISTRIBUTION INC.

REPORT ANY SERIOUS ADVERSE EVENTS TO:

VIADERMA DISTRIBUTION INC.
1050 E FLAMINGO RD., SUITE 107,
LAS VEGAS, NV 89119

TEL: 1-800-585-8685 WWW.VIADERMAINC.COM

MADE IN THE USA

Tetracycline Hydrochloride 3%

Topical Ointment

First Aid Antibiotic

VITASTEM™

VIADERMA®

VIADERMA®
Pharma

VITASTEM™
First Aid Antibiotic

Topical Ointment

Tetracycline
Hydrochloride 3%



VIADERMA®
Pharma

VITASTEM™
First Aid Antibiotic

Topical Ointment

Tetracycline
Hydrochloride 3%



Net Wt. 0.5 oz (15 g)



CONTAINS NO ALCOHOL

NO ANIMAL INGREDIENTS



WARNINGS: FOR EXTERNAL USE ONLY. DO NOT USE LONGER THAN THREE WEEKS. DIRECTED BY A DOCTOR IN THE TREATMENT OF SERIOUS BURNS, LACERATIONS OF THE BODY, ASK A DOCTOR BEFORE USE IF YOU HAVE DEEP OR PUNCTURE WOUNDS, ANIMAL BITES, OR SERIOUS BURNS. STOP USE AND ASK A DOCTOR IF CONDITION PERSISTS OR GETS WORSE. KEEP OUT OF REACH OF CHILDREN. IF SWALLOWED, GET MEDICAL HELP OR CONTACT A POISON CONTROL CENTER RIGHT AWAY.

DIRECTIONS: CLEAN THE AFFECTED AREA. APPLY A SMALL AMOUNT OF THIS PRODUCT ON THE AREA. TO 3 TIMES ONLY MAY BE COVERED WITH A STERILE DRESSING. DISTRIBUTED BY: VIADERMA DISTRIBUTION, INC. REPORT ANY SERIOUS ADVERSE EVENTS TO: VIADERMA DISTRIBUTION, INC. 1080 FERNANDO SUITE 107, LAS VEGAS, NV 89119 TEL: 1-800-595-8085 WWW.VIADERMA.COM MADE IN THE USA

VIADERMA[®]
Pharma
VITASTEM[™]
First Aid Antibiotic

Topical Ointment

Tetracycline Hydrochloride 3%

Net Wt. 0.5 oz (15 g)



VITASTEM FIRST AID ANTIBIOTIC

tetracycline hydrochloride ointment

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:71262-012 |
| Route of Administration | TOPICAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|----------------------------|--------------|
| TETRACYCLINE HYDROCHLORIDE (UNII: P6R62377KV) (TETRACYCLINE - UNII:F8VB5M810T) | TETRACYCLINE HYDROCHLORIDE | 30 mg in 1 g |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| ACETIC ACID (UNII: Q40Q9N063P) | |
| ASCORBIC ACID (UNII: PQ6CK8PD0R) | |
| CHLORHEXIDINE GLUCONATE (UNII: MOR84MUD8E) | |
| CHOLECALCIFEROL (UNII: 1C6V77QF41) | |
| DIMETHYL SULFOXIDE (UNII: YOW8V9698H) | |
| DIPROPYLENE GLYCOL (UNII: E107L85C40) | |
| GLUCONOLACTONE (UNII: WQ29KQ9POT) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| HISTIDINE (UNII: 4QD397987E) | |
| HYDROXYETHYL CELLULOSE, UNSPECIFIED (UNII: T4V6TWG28D) | |
| MAGNESIUM STEARATE (UNII: 70097M6130) | |
| METHYLPARABEN (UNII: A2I8C7HI9T) | |
| SODIUM HYDROXIDE (UNII: 55X04QC32I) | |
| SORBIC ACID (UNII: X045WJ989B) | |

WATER (UNII: 059QF0KO0R)

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:71262-012-01 | 1 in 1 CARTON | 03/10/2025 | |
| 1 | | 15 g in 1 BOTTLE, DROPPER; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M004 | 03/10/2025 | |

Labeler - VIADERMA DISTRIBUTION INC (081113521)

Registrant - VIADERMA DISTRIBUTION INC (081113521)

Revised: 12/2025

VIADERMA DISTRIBUTION INC