

**NASAL- oxymetazoline hcl spray, metered**  
**Meijer Distribution Inc**

-----  
**Meijer Distribution, Inc. Nasal Spray Drug Facts**

**Active ingredient**

Oxymetazoline hydrochloride 0.05%

**Purpose**

Nasal decongestant

**Uses**

- temporarily relieves nasal congestion due to:
- common cold
- hay fever
- upper respiratory allergies
- temporarily relieves sinus congestion and pressure
- shrinks swollen nasal membranes so you can breathe more freely

**Warnings**

**Ask a doctor before use if you have**

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland

**When using this product**

- **do not use more than directed**
- do not use for more than 3 days. Use only as directed. Frequent or prolonged use may cause nasal congestion to recur or worsen.
- temporary discomfort such as burning, stinging, sneezing or an increase in nasal discharge may occur
- use of this container by more than one person may spread infection

**Stop use and ask a doctor if**

symptoms persist

**If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### **Directions**

- adults and children 6 to under 12 years of age (with adult supervision): 2 or 3 sprays in each nostril not more often than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period.
- children under 6 years of age: ask a doctor

**To Use:** Shake well before use. Hold white tabs, SQUEEZE grooved area of cap FIRMLY and turn counter clockwise. Before using the first time, prime metered pump by depressing pump firmly several times. To spray, hold bottle with thumb at base and nozzle between first and second fingers. Without tilting head, insert nozzle into nostril. Fully depress rim with a firm, even stroke and sniff deeply. Wipe nozzle clean after use. Secure cap after use

### **Other information**

- store at 20-25°C (68-77°F)
- retain carton for future reference on full labeling

### **Inactive ingredients**

benzalkonium chloride solution, benzyl alcohol, dibasic sodium phosphate, edetate disodium, microcrystalline cellulose and carboxymethylcellulose sodium, monobasic sodium phosphate, polyethylene glycol, povidone, purified water

### **Questions or comments?**

**1-800-719-9260**

### **Principal Display Panel**

meijer®

COMPARE TO AFRIN® NO DRIP ACTIVE INGREDIENT

MAXIMUM STRENGTH

Nasal Spray

Oxymetazoline Hydrochloride

12 HOUR NASAL DECONGESTANT

NO DRIP

Fast, Powerful Congestion Relief

Steroid Free

For Colds & Allergies

1 FL OZ (30 mL)

**meijer.** NDC 41250-388-10  
MAXIMUM STRENGTH  
**Nasal Spray**  
Oxymetazoline Hydrochloride  
12 HOUR NASAL DECONGESTANT  
Fast, Powerful Congestion Relief  
1 FL OZ (30 mL)

COMPARE TO AFRIN® NO DRIP ACTIVE INGREDIENT\*

**NO DRIP**

Steroid Free

For Colds & Allergies

**meijer.** NDC 41250-388-10  
MAXIMUM STRENGTH  
**Nasal Spray**  
Oxymetazoline Hydrochloride  
12 HOUR NASAL DECONGESTANT  
**NO DRIP**  
For Colds & Allergies  
1 FL OZ (30 mL)  
Actual Size

DO NOT USE IF PRINTED NECK AND IS BROKEN OR MISSING

OUR QUALITY GUARANTEE

www.meijer.com/satisfaction  
www.meijer.com  
GRAND RAPIDS, MI 49544  
DISTRIBUION, INC.  
DIST. BY MEIJER

**Drug Facts**  
**Active ingredient** Oxymetazoline hydrochloride 0.05%...  
**Purpose** Nasal decongestant

**Uses**  
temporarily relieves nasal congestion due to:  
■ common cold ■ hay fever  
■ upper respiratory allergies  
temporarily relieves sinus congestion and pressure  
shrinks swollen nasal membranes so you can breathe more freely

**Warnings**  
Ask a doctor before use if you have  
■ heart disease ■ high blood pressure  
■ thyroid disease ■ diabetes  
■ trouble urinating due to an enlarged prostate gland

**When using this product**  
do not use more than directed  
do not use for more than 3 days. Use only as directed. Frequent or prolonged use may cause nasal congestion to recur or worsen.  
temporary discomfort such as burning, stinging, sneezing or an increase in nasal discharge may occur  
use of this container by more than one person may spread infection

**Stop use and ask a doctor if symptoms persist**  
If pregnant or breast-feeding, ask a health professional before use.  
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**  
adults and children 6 to under 12 years of age (with adult supervision): 2 or 3 sprays in each nostril not more often than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period.  
children under 6 years of age: ask a doctor  
**To Use:** Shake well before use. Hold white tabs, **SQUEEZE** grooved area of cap **FIRMLY** and turn counter clockwise. Before using the first time, prime metered pump by depressing pump firmly several times. To spray, hold bottle with thumb at base and nozzle between first and second fingers. Without tilting head, insert nozzle into nostril. Fully depress rim with a firm, even stroke and sniff deeply. Wipe nozzle clean after use. Secure cap after use.

**Other information**  
store at 20-25°C (68-77°F)  
retain carton for future reference on full labeling

**Drug Facts (continued)**  
**Inactive ingredients**  
benzalkonium chloride solution, benzyl alcohol, dibasic sodium phosphate, edetate disodium, microcrystalline cellulose and carboxymethylcellulose sodium, monobasic sodium phosphate, polyethylene glycol, povidone, purified water

**Questions or comments?**  
1-800-719-9260

\*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Afrin®.

**How to use:**  
Hold white tabs, **SQUEEZE** grooved area of cap **FIRMLY** and turn counter clockwise. Secure cap after use.  
**SQUEEZE** grooves **FIRMLY** while turning

PID 344736  
0 41250-04750 6

PAPER BOX PLASTIC BOTTLE  
how2recycle.info

38810 6E C7

NASAL

oxymetazoline hcl spray, metered

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:41250-388
<b>Route of Administration</b>	NASAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>OXYMETAZOLINE HYDROCHLORIDE</b> (UNII: K89MJ0S5VY) (OXYMETAZOLINE - UNII:8VLN5B44ZY)	OXYMETAZOLINE HYDROCHLORIDE	0.05 g in 100 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>BENZYL ALCOHOL</b> (UNII: LKG8494WBH)	
<b>CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED</b> (UNII: K679OBS311)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>BENZALKONIUM CHLORIDE</b> (UNII: F5UM2KM3W7)	
<b>SODIUM PHOSPHATE, DIBASIC, UNSPECIFIED FORM</b> (UNII: GR686LBA74)	
<b>SODIUM PHOSPHATE, MONOBASIC, UNSPECIFIED FORM</b> (UNII: 3980JIH2SW)	

## Product Characteristics

<b>Color</b>	WHITE (to off white, viscous)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-388-10	1 in 1 CARTON	09/11/2002	
1		30 mL in 1 BOTTLE, SPRAY; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/11/2002	

