

SENNA S- docusate sodium, sennosides tablet
TOP CARE (Topco Associates LLC)

Drug Facts

Active ingredients (in each tablet)

Docusate sodium 50 mg
Sennosides 8.6 mg

Purpose

Stool softener
Laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in 6-12 hours

Warnings

Do not use

- if you are now taking mineral oil, unless directed by a doctor
- laxative products for longer than 1 week unless directed by a doctor

Ask a doctor before use if you have

- stomach pain
- nausea
- vomiting
- noticed a sudden change in bowel habits that continues over a period of 2 weeks

Stop use and ask a doctor if

you have rectal bleeding or fail to have a bowel movement after use of a laxative. These may indicate a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take preferably at bedtime or as directed by a doctor

age	starting dosage	maximum dosage
adults and children 12 years of age and over	2 tablets once a day	4 tablets twice a day
children 6 to under 12 years	1 tablet once a day	2 tablets twice a day
children 2 to under 6 years	1/2 tablet once a day	1 tablet twice a day
children under 2 years	ask a doctor	ask a doctor

Other information

- **each tablet contains:** calcium 30 mg
- **each tablet contains:** sodium 6 mg LOW SODIUM
- store at 25°C (77°F); excursions permitted between 15-30°C (59-86°F)

Inactive ingredients

croscarmellose sodium, dibasic calcium phosphate dihydrate, FD&C yellow #6 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, silicon dioxide, talc, titanium dioxide

Questions or comments?

Call toll free **1-888-423-0139**

Principal Display Panel

COMPARE TO SENOKOT-S® ACTIVE INGREDIENTS†

Vegetable Laxative

+ Stool Softener

DOCUSATE SODIUM 50 mg • STOOL SOFTENER

SENNOSIDES 8.6 MG • LAXATIVE

TABLETS

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

†This product is not manufactured or distributed by Avrio Health L.P., distributor of Senokot-S®.

DISTRIBUTED BY TOPCO ASSOCIATES LLC

ELK GROVE VILLAGE, IL 60007

topcare@topco.com

www.topcarebrand.com

Product Labeling

PLD-05994
FC006101
Lot No.:
Exp. Date:

0 36800 43856 9





COMPARE TO SENOKOT-S® ACTIVE INGREDIENTS†

Vegetable Laxative + Stool Softener

DOCUSATE SODIUM 50 mg • STOOL SOFTENER
SENNOSIDES 8.6 mg • LAXATIVE

120 TABLETS

actual
size



DISTRIBUTED BY TOPCO ASSOCIATES LLC
ELK GROVE VILLAGE, IL 60007
©TOPCO PLVA0521
QUESTIONS? 1-888-423-0139
topcare@topco.com
www.topcarebrand.com



Scan here for more
information or
call 1-888-423-0139

This product is not manufactured or distributed by Avro Health L.P., distributor of Senokot-S®.

Warnings

Do not use if you are now taking mineral oil, unless directed by a doctor

laxative products for longer than 1 week unless directed by a doctor

Ask a doctor before use if you have

- stomach pain ■ nausea ■ vomiting
- noticed a sudden change in bowel habits that continues over a period of 2 weeks

Stop use and ask a doctor if you have rectal bleeding or fail to have a bowel movement after use of a laxative. These may indicate a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

take preferably at bedtime or as directed by a doctor

age	starting dosage	maximum dosage
adults and children 12 years of age and over	2 tablets once a day	4 tablets twice a day
children 6 to under 12 years	1 tablet once a day	2 tablets twice a day
children 2 to under 6 years	1/2 tablet once a day	1 tablet twice a day
children under 2 years	ask a doctor	ask a doctor

Drug Facts

Active ingredients (in each tablet)

Docusate sodium 50 mg

Sennosides 8.6 mg

Stool softener

Laxative

Uses

relieves occasional constipation (regularly) generally produces bowel movement in 6-12 hours

Warnings

Do not use if you are now taking mineral oil, unless directed by a doctor

laxative products for longer than 1 week unless directed by a doctor

Ask a doctor before use if you have

- stomach pain ■ nausea ■ vomiting
- noticed a sudden change in bowel habits that continues over a period of 2 weeks

Stop use and ask a doctor if you have rectal bleeding or fail to have a bowel movement after use of a laxative. These may indicate a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Other information

- each tablet contains: calcium 30 mg
- each tablet contains: sodium 6 mg
- store at 25°C (77°F); excursions permitted between 15-30°C (59-86°F)

Inactive ingredients croscarmellose sodium, dibasic calcium phosphate dinaryrate, FD&C yellow #6 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, silicon dioxide, talc, titanium dioxide

Questions or comments? Call toll free 1-888-423-0139

Drug Facts (continued)

Other information

- each tablet contains: calcium 30 mg
- each tablet contains: sodium 6 mg
- store at 25°C (77°F); excursions permitted between 15-30°C (59-86°F)

Inactive ingredients croscarmellose sodium, dibasic calcium phosphate dinaryrate, FD&C yellow #6 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, silicon dioxide, talc, titanium dioxide

Questions or comments? Call toll free 1-888-423-0139

TOPCARE HEALTH Stool Softener

SENNAS

docusate sodium, sennosides tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:36800-783
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DOCUSATE SODIUM (UNII: F05Q2T2JA0) (DOCUSATE - UNII:M7P27195AG)	DOCUSATE SODIUM	50 mg
SENNOSIDES (UNII: 3FYP5M0IJX) (SENNOSIDES - UNII:3FYP5M0IJX)	SENNOSIDES	8.6 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
ALUMINUM OXIDE (UNII: LMI26O6933)	

Product Characteristics

Color	orange	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	PSD22
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-783-60	1 in 1 BOX	07/31/2019	
1		60 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:36800-783-12	1 in 1 BOX	07/31/2019	
2		120 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

OTC Monograph Drug 505G(a)(3)

07/31/2019

Labeler - TOP CARE (Topco Associates LLC) (006935977)

Revised: 8/2024

TOP CARE (Topco Associates LLC)