

ORMALVI- dichlorphenamide tablet

Cycle Pharmaceuticals Ltd.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use ORMALVI™ safely and effectively. See full prescribing information for ORMALVI™.

ORMALVI™ (dichlorphenamide) tablets, for oral use
Initial U.S. Approval: 1958

INDICATIONS AND USAGE

ORMALVI is an oral carbonic anhydrase inhibitor indicated for the treatment of primary hyperkalemic periodic paralysis, primary hypokalemic periodic paralysis, and related variants (1)

DOSAGE AND ADMINISTRATION

- Initiate dosing at 50 mg by mouth once or twice daily (2.1)
- Titrate up or down dose based on individual response (2.1)
- The minimum recommended dosage is 50 mg daily, and the maximum recommended dosage is 200 mg daily (2.1)
- Evaluate response to ORMALVI after 2 months of treatment (2.2)

DOSAGE FORMS AND STRENGTHS

Tablets: 50 mg (3)

CONTRAINDICATIONS

- Hepatic insufficiency (4)
- Severe pulmonary obstruction (4)
- Hypersensitivity to dichlorphenamide or other sulfonamides (4)
- Concomitant use with high dose aspirin (4)

WARNINGS AND PRECAUTIONS

- Hypersensitivity and Other Life-Threatening Reactions: discontinue ORMALVI at the first appearance of skin rash or any sign of immune-mediated or idiosyncratic adverse reaction (5.1)
- Hypokalemia: baseline and periodic measurements of serum potassium are recommended; if hypokalemia develops or persists, consider reducing the dose or discontinuing ORMALVI and correcting potassium levels (5.3)
- Metabolic acidosis: baseline and periodic measurements of serum bicarbonate are recommended; if metabolic acidosis develops or persists, consider reducing the dose or discontinuing ORMALVI (5.4)
- Falls: consider reducing the dose or discontinuing ORMALVI in patients who experience falls (5.5)

ADVERSE REACTIONS

Most common adverse reactions (incidence at least 10% and greater than placebo) include paresthesias, cognitive disorder, dysgeusia, and confusional state (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Cycle Pharmaceuticals Ltd at 1-857-437-3969 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

Aspirin: anorexia, tachypnea, lethargy, and coma have been reported with concomitant use of dichlorphenamide and high-dose aspirin. The concomitant use of ORMALVI and high-dose aspirin is contraindicated. ORMALVI should be used with caution in patients receiving lower doses of aspirin (4, 5.2, 7.1)

USE IN SPECIFIC POPULATIONS

Pregnancy: Based on animal data, may cause fetal harm (8.1)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 7/2025

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Dosage Information
- 2.2 Monitoring to Assess Effectiveness
- 2.3 Monitoring to Assess Safety

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Hypersensitivity and Other Life-threatening Reactions
- 5.2 Concomitant Use of Aspirin or Other Salicylates
- 5.3 Hypokalemia
- 5.4 Metabolic Acidosis
- 5.5 Falls

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

7 DRUG INTERACTIONS

- 7.1 Aspirin and Other Salicylates
- 7.2 Drugs that are Substrates of Organic Anion Transporter1 (OAT1)
- 7.3 Drugs that Cause Hypokalemia
- 7.4 Drugs that Cause Metabolic Acidosis
- 7.5 Drugs that are Inhibitors of OAT1 or OAT3

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

ORMALVI is indicated for the treatment of primary hyperkalemic periodic paralysis, primary hypokalemic periodic paralysis, and related variants.

2 DOSAGE AND ADMINISTRATION

2.1 Dosage Information

Initiate dosing at 50 mg by mouth once or twice daily. The dosage may be increased or decreased based on individual response, at weekly intervals (or sooner in case of adverse reaction). The minimum recommended total daily dosage is 50 mg, and the maximum recommended total daily dosage is 200 mg.

2.2 Monitoring to Assess Effectiveness

Primary hyperkalemic periodic paralysis, primary hypokalemic periodic paralysis, and related variants are a heterogeneous group of conditions, for which the response to ORMALVI may vary. Therefore, prescribers should evaluate the patient's response to ORMALVI after 2 months of treatment to decide whether ORMALVI should be continued.

2.3 Monitoring to Assess Safety

Baseline and periodic measurements of serum potassium and sodium bicarbonate during ORMALVI treatment is recommended [*see Warnings and Precautions (5.3, 5.4)*].

3 DOSAGE FORMS AND STRENGTHS

White to off-white, round shaped, flat faced beveled edged tablet, scored on one side, engraved with 'ZD' above the score and '50' below the score, the other side is plain.

4 CONTRAINDICATIONS

ORMALVI is contraindicated in the following circumstances:

- Hypersensitivity to dichlorophenamide or other sulfonamides [*see Warnings and Precautions (5.1)*]
- Concomitant use of ORMALVI and high dose aspirin [*see Warnings and Precautions (5.2) and Drug Interactions (7.1)*]
- Severe pulmonary disease, limiting compensation to metabolic acidosis caused by ORMALVI [*see Warnings and Precautions (5.4)*]
- Hepatic insufficiency: ORMALVI may aggravate hepatic encephalopathy.

5 WARNINGS AND PRECAUTIONS

5.1 Hypersensitivity and Other Life-threatening Reactions

Fatalities associated with the administration of sulfonamides have occurred because of adverse reactions including Stevens-Johnson syndrome, toxic epidermal necrolysis, fulminant hepatic necrosis, agranulocytosis, aplastic anemia, and other blood dyscrasias. Pulmonary involvement can occur in isolation or as part of a systemic reaction.

ORMALVI should be discontinued at the first appearance of skin rash or any sign of

immune-mediated or other life-threatening adverse reaction.

5.2 Concomitant Use of Aspirin or Other Salicylates

Carbonic anhydrase inhibitors, including ORMALVI, can cause metabolic acidosis [see *Warnings and Precautions (5.4)*], which can increase the risk of salicylate toxicity. Anorexia, tachypnea, lethargy, and coma have been reported with concomitant use of dichlorophenamide and high-dose aspirin. Therefore, the concomitant use of ORMALVI and high-dose aspirin is contraindicated. Patients with concomitant use of ORMALVI and low-dose aspirin should be carefully monitored.

5.3 Hypokalemia

ORMALVI increases potassium excretion and can cause hypokalemia. The risk of hypokalemia is greater when ORMALVI is used in patients with conditions associated with hypokalemia (e.g., adrenocortical excess, renal tubular acidosis type 1 and 2), and in patients receiving other drugs that may cause hypokalemia [see *Drug Interactions (7.3)*].

Baseline and periodic measurements of serum potassium during ORMALVI treatment is recommended.

If hypokalemia develops or persists, consideration should be given to reducing the dose or discontinuing ORMALVI and correction of potassium levels.

5.4 Metabolic Acidosis

ORMALVI can cause hyperchloremic non-anion gap metabolic acidosis. Concomitant use of ORMALVI with other drugs that cause metabolic acidosis may increase the severity of acidosis. Concomitant use of ORMALVI in compensated patients with respiratory acidosis, such as in advanced lung diseases, may lead to respiratory decompensation.

Baseline and periodic measurements of serum bicarbonate during ORMALVI treatment are recommended.

If metabolic acidosis develops or persists, consideration should be given to reducing the dose or discontinuing ORMALVI [see *Drug Interactions (7.4)*].

5.5 Falls

ORMALVI increases the risk of falls. The risk of falls is greater in the elderly and with higher doses of ORMALVI. Consider dose reduction or discontinuation of ORMALVI in patients who experience falls while treated with ORMALVI.

6 ADVERSE REACTIONS

The following serious adverse reactions are described elsewhere in labeling:

- Hypersensitivity and Other Life-Threatening Reactions [see *Warnings and Precautions (5.1)*]
- Hypokalemia [see *Warnings and Precautions (5.3)*]
- Metabolic Acidosis [see *Warnings and Precautions (5.4)*]
- Falls □ [see *Warnings and Precautions (5.5)*]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

In a 9-week randomized controlled trial in adults with hyperkalemic or hypokalemic periodic paralysis (Study 1), the most common adverse reactions in patients treated with dichlorphenamide, with rates greater than placebo, were paresthesia, cognitive disorder, dysgeusia, and confusional state. The mean dose of dichlorphenamide was 94 mg/day in patients with hypokalemic periodic paralysis and 82 mg/day in patients with hyperkalemic periodic paralysis.

Table 1 lists the incidence of adverse reactions that occurred in 5% of patients treated with dichlorphenamide and more commonly than in patients treated with placebo in Study 1.

Table 1: Adverse Reactions in Patients Treated with Dichlorphenamide with Incidence 5% and more common than in Patients Treated with Placebo in Study 1

	Adverse Reaction	Dichlorphenamide N = 36 (%)	Placebo N = 29 (%)
Nervous system disorders	Paresthesia	44	14
	Cognitive disorder ¹	14	7
	Dysgeusia	14	0
	Confusional state	11	0
	Headache	8	7
	Hypoesthesia	8	0
	Lethargy	8	0
	Dizziness	6	0
Gastrointestinal disorders	Diarrhea	6	3
	Nausea	6	0
General disorders and administration site conditions	Fatigue	8	0
	Malaise	6	0
Investigations	Weight decreased	6	0
Musculoskeletal and connective tissue disorders	Muscle spasms	8	0
	Arthralgia	6	3
	Muscle twitching	6	0
Respiratory	Dyspnea	6	0
	Pharyngolaryngeal pain	6	0
Skin	Rash	8	0
	Pruritus	6	0

¹Cognitive disorder combined cases with the preferred terms of cognitive disorder, disturbance in attention, and mental impairment.

6.2 Postmarketing Experience

Adverse reactions have been identified during postapproval use of dichlorphenamide. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

The following are adverse reactions which have been reported during postapproval use of dichlorphenamide and were serious or are not reported in the previous section of labeling [see *Clinical Trials Experience (6.1)*]: amnesia, cardiac failure, condition aggravated, convulsion, hallucination, nephrolithiasis, pancytopenia, psychotic disorder, renal tubular necrosis, stupor, syncope, tremor.

7 DRUG INTERACTIONS

7.1 Aspirin and Other Salicylates

Carbonic anhydrase inhibitors, including ORMALVI, can cause metabolic acidosis [see *Warning and Precautions (5.2, 5.4)*], which can increase the risk of salicylate toxicity. Anorexia, tachypnea, lethargy, and coma have been reported with concomitant use of dichlorphenamide and high-dose aspirin. Therefore, concomitant use of ORMALVI and high-dose aspirin is contraindicated. Patients with concomitant use of ORMALVI and low-dose aspirin should be carefully monitored [see *Contraindications (4) and Warnings and Precautions (5.2)*].

7.2 Drugs that are Substrates of Organic Anion Transporter1 (OAT1)

In vitro, dichlorphenamide is an inhibitor of OAT1 transporters. The concomitant administration of ORMALVI may increase the plasma exposures of OAT1 substrates. Use of ORMALVI with drugs that are sensitive to OAT1 inhibition (e.g., methotrexate, famotidine, oseltamivir) is not recommended [see *Clinical Pharmacology (12.3)*].

7.3 Drugs that Cause Hypokalemia

The risk of hypokalemia is greater with coadministration of ORMALVI and other drugs that can cause hypokalemia (e.g., loop diuretics, thiazide diuretics, laxatives, antifungals, penicillins, and theophylline) [see *Warnings and Precautions (5.3)*].

7.4 Drugs that Cause Metabolic Acidosis

Coadministration of ORMALVI and other drugs that can cause metabolic acidosis may increase the severity of the acidosis [see *Warnings and Precautions (5.4)*].

7.5 Drugs that are Inhibitors of OAT1 or OAT3

An *in vitro* transporter study indicated that dichlorphenamide is a substrate of human transporters OAT1 and OAT3 [see *Clinical Pharmacology (12.3)*]. Therefore, signs of dichlorphenamide toxicity should be monitored when administered with OAT1 or OAT3 inhibitors.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

There are no adequate data on the developmental risk associated with the use of ORMALVI in pregnant women. A no-effect dose has not been established. Dichlorphenamide was teratogenic when administered orally to pregnant rats.

The background risk of major birth defects and miscarriage for the indicated population is unknown. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4%, and 15% to 20%, respectively.

Clinical Considerations

Fetal/Neonatal adverse reactions

ORMALVI treatment can cause metabolic acidosis [see *Warnings and Precautions (5.4)*]. The effect of dichlorphenamide-induced metabolic acidosis has not been studied in pregnancy; however, metabolic acidosis in pregnancy (due to other causes) can cause decreased fetal growth, decreased fetal oxygenation, and fetal death, and may affect the fetus' ability to tolerate labor. Pregnant patients should be monitored for metabolic acidosis and treated as in the nonpregnant state. Newborns of mothers treated with ORMALVI should be monitored for metabolic acidosis because of possible occurrence of transient metabolic acidosis following birth.

Labor or Delivery

Although the effect of ORMALVI on labor and delivery in humans has not been established, the development of dichlorphenamide-induced metabolic acidosis in the mother and/or in the fetus might affect the fetus' ability to tolerate labor.

Data

Animal Data

Teratogenic effects (fetal limb reduction defects) were reported following oral administration of dichlorphenamide to pregnant rats during organogenesis at 350 mg/kg, or 17 times the maximum recommended human dose (200 mg/day) on a body surface area (mg/m²) basis. A no-effect dose for adverse effects on embryofetal development has not been established.

8.2 Lactation

Risk Summary

There are no data on the presence of dichlorphenamide in human milk, the effects on the breastfed infant, or the effects on milk production.

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for ORMALVI and any potential adverse effects on the breastfed infant from ORMALVI or from the underlying maternal condition.

8.4 Pediatric Use

Safety and effectiveness of ORMALVI in pediatric patients have not been established.

8.5 Geriatric Use

The risk of falls and of metabolic acidosis are greater in elderly patients [see *Warnings and Precautions* (5.4, 5.5)].

10 OVERDOSAGE


Symptoms of overdose or toxicity may include drowsiness, anorexia, nausea, vomiting, dizziness, ataxia, tremor, and tinnitus.

In the event of overdose, induce emesis or perform gastric lavage. The electrolyte disturbances most likely to be encountered from overdose are hypokalemia and hyperchloremic metabolic acidosis.

11 DESCRIPTION

ORMALVI tablets, USP contain dichlorphenamide, an oral carbonic anhydrase inhibitor. Dichlorphenamide, USP a dichlorinated benzenedisulfonamide, is known chemically as 4,5-dichloro-1,3-benzenedisulfonamide.

Its empirical formula is $C_6H_6Cl_2N_2O_4S_2$ and its structural formula is:



Dichlorphenamide USP is a white or practically white, crystalline powder with a molecular weight of 305.16. It is very slightly soluble in water but soluble in dilute solutions of alkali carbonate and hydroxide.

ORMALVI (dichlorphenamide) tablets, USP are supplied as tablets, for oral administration, each containing 50 mg dichlorphenamide. Inactive ingredients are lactose monohydrate, partially pregelatinized starch and magnesium stearate.

FDA approved dissolution test specifications differ from USP.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Dichlorphenamide is a carbonic anhydrase inhibitor. However, the precise mechanism by which dichlorphenamide exerts its therapeutic effects in patients with primary periodic paralysis is unknown.

12.2 Pharmacodynamics

ORMALVI can cause metabolic acidosis, which can increase the risk of salicylate toxicity

with coadministration [see *Warnings and Precautions (5.2)*]. ORMALVI-induced metabolic acidosis can also increase in severity with coadministration of other drugs that cause metabolic acidosis [see *Warnings and Precautions (5.4)*].

12.3 Pharmacokinetics

After single-dose administration in healthy subjects in fasted state, dichlorphenamide C_{max} and AUC increased in a dose-proportional manner within the range of 25 mg to 400 mg (2 times the maximum recommended dose). The steady-state is expected to be achieved within 10 days of twice-daily dosing.

Absorption

The median time to reach maximum concentration (T_{max}) of dichlorphenamide was about 1.5 to 3 hours postdose after both single and multiple dose administrations.

Distribution

The plasma protein binding of dichlorphenamide is approximately 88%.

Elimination

Following a single-dose administration, mean terminal half-life was in the range of 32 to 66 hours.

Metabolism

Dichlorphenamide is not a substrate for CYP1A2, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP2D6 and CYP3A4 isoforms when tested *in vitro*.

Drug Interaction Studies

In vitro Assessment of Drug Interactions

Drug-Metabolizing Enzyme Inhibition

Dichlorphenamide is not an inhibitor for CYP1A2, 2B6, 2C8, 2C9, 2C19, 2D6, or 3A4 enzymes when tested *in vitro*.

Drug-Metabolizing Enzyme Induction

Dichlorphenamide is not an inducer for CYP1A2, 2B6, or 3A4 enzymes when tested *in vitro*.

In vitro Assessment of Transporter-Drug Interactions

Dichlorphenamide is neither a substrate nor inhibitor for p-gp, BCRP, OATP1B1, OATP1B3, OAT2, OAT4, OCT1, OCT2, MATE1, or MATE2-K when tested *in vitro*.

Dichlorphenamide is not an inhibitor of OAT3, but is an inhibitor of OAT1 based on *in vitro* studies [see *Drug Interactions (7.4)*].

Dichlorphenamide is a substrate for transporters OAT1 and OAT3 based on *in vitro* studies [see *Drug Interactions (7.2)*].

In Vivo Drug Interactions

The use of dichlorphenamide in combination with high-dose aspirin is contraindicated as it may lead to salicylate toxicity. The mechanism(s) of this interaction is not known.

Specific Populations

Geriatrics

The pharmacokinetics of dichlorphenamide in the elderly has not been determined.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Studies to assess the carcinogenic potential of dichlorphenamide have not been conducted.

Mutagenesis

Studies to assess the genotoxicity of dichlorphenamide have not been conducted.

Impairment of Fertility

Studies to assess the effects of dichlorphenamide on fertility have not been conducted.

14 CLINICAL STUDIES

The efficacy of dichlorphenamide was evaluated in two clinical studies, Study 1 and Study 2.

Study 1

Study 1 was a 9-week, double-blind, placebo-controlled multi-center study. Study 1 consisted of two substudies: a substudy in patients with hypokalemic periodic paralysis (n=44), and a substudy in patients with hyperkalemic periodic paralysis (n=21). The primary efficacy endpoint in both substudies was the average number of self-reported attacks of muscle weakness per week over the final 8 weeks of the trial. Withdrawal from the study for acute severe worsening (increase in attack frequency or severity) was also assessed as an endpoint.

In Study 1, the dose of dichlorphenamide was 50 mg b.i.d. for treatment-naïve patients. Patients already on dichlorphenamide prior to the study continued on the same dose while on dichlorphenamide during the study. In patients taking acetazolamide prior to the study, the dose of dichlorphenamide was set at 20% of the acetazolamide dose. Dose reduction for tolerability was permitted.

Hypokalemic Periodic Paralysis Substudy of Study 1

In the hypokalemic periodic paralysis substudy, median age of patients was 45 years and 73% of patients were male. Patients treated with dichlorphenamide (n=24) had 2.2 fewer attacks per week than patients (n=20) treated with placebo (p=0.02). None of the patients randomized to dichlorphenamide reached the endpoint of withdrawal from the study for acute worsening, vs. five patients randomized to placebo. The mean dose of dichlorphenamide at Week 9 was 94 mg/day.

Hyperkalemic Periodic Paralysis Substudy of Study 1

In the Hyperkalemic Periodic Paralysis substudy, median age of patients was 43 years and 43% of patients were male. During the double-blind treatment period, patients treated with dichlorphenamide (n=12) had 3.9 fewer attacks per week than patients

anaphylaxis or other life-threatening reactions [see *Warnings and Precautions (5.1)*].

Drug Interactions

Instruct patients to notify their healthcare provider of all of the drugs and over-the-counter medications that they take and to not take aspirin or other salicylates without first discussing with their healthcare provider [see *Drug Interactions (7.1, 7.2, 7.3, 7.4)*].

Metabolic Acidosis

Instruct patients to contact their healthcare provider immediately if they develop possible manifestations of metabolic acidosis (e.g., fast breathing, fatigue/tiredness, loss of appetite, or irregular heartbeat or palpitations) [see *Warnings and Precautions (5.4)*].

Falls

Inform patients that ORMALVI can increase their risk of falls [see *Warnings and Precautions (5.5)*].

Cognitive Impairment

Advise patients to notify their healthcare provider if they experience symptoms of cognitive impairment including confusion and memory lapse [see *Adverse Reactions (6.1)*].

Driving and Operating Machinery

ORMALVI may cause drowsiness/fatigue in some patients [see *Adverse Reactions (6.1)*]. Caution patients on the potential for impaired ability to drive and operate machinery.



Manufactured by:

ZENARA PHARMA PVT. LTD., Hyderabad-500051, INDIA.

Manufactured for:

CYCLE PHARMACEUTICALS LTD

The Broers Building, 21 JJ Thomson Ave, Cambridge, CB3 0FA, United Kingdom

AW000377

Revised: July 2025

ORMALVI™ (dichlorphenamide) 50 mg tablets, 100s Count

Each tablet contains 50 mg of dichlorphenamide, USP
Usual Dosage: See Prescribing Information for dosage information.
 Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature].
 Keep this and all medications out of the reach of children.
 Dispense in a tight container with a child-resistant closure.

Manufactured by: ZENARA PHARMA PVT. LTD. Hyderabad-500051, INDIA.
Manufactured for: Cycle Pharmaceuticals Ltd Cambridge, CB3 0FA, UK

100 Tablets NDC 70709-085-10

ORMALVI™
 (dichlorphenamide)
 tablets, USP

AW000306

50 mg

CYCLE PHARMA

Rx only

3 70709 08510 4

LAB-PS0211/00 Neutral Code: DRUGS/AP/01/2006

Over Printing Zone

ORMALVI

dichlorphenamide tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:70709-085
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DICHLORPHENAMIDE (UNII: VVJ6673MHY) (DICHLORPHENAMIDE - UNII:VVJ6673MHY)	DICHLORPHENAMIDE	50 mg

Inactive Ingredients

Ingredient Name	Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
STARCH, CORN (UNII: O8232NY3SJ)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

Product Characteristics

Color	white (to off-white)	Score	2 pieces
Shape	ROUND (flat faced beveled edged)	Size	6mm
Flavor		Imprint Code	ZD;50
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70709-085-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	03/15/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA215924	03/15/2024	

Labeler - Cycle Pharmaceuticals Ltd. (218215530)

Establishment

Name	Address	ID/FEI	Business Operations
Zenara Pharma Pvt. Ltd.		924839850	manufacture(70709-085) , pack(70709-085) , analysis(70709-085) , label(70709-085)

Revised: 7/2025

Cycle Pharmaceuticals Ltd.