

ASPIRIN LOW DOSE- aspirin tablet, chewable
L.N.K. International, Inc.

Quality Plus 44-218

Active ingredient (in each chewable tablet)

Aspirin 81 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

for the temporary relief of minor aches and pains or as recommended by your doctor

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- are age 60 or older
- take more or for a longer time than directed
- have had stomach ulcers or bleeding problems

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - vomit blood
 - have bloody or black stools
 - feel faint
 - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- drink a full glass of water with each dose
- chew or crush tablets completely before swallowing; do not swallow tablets whole
- adults and children 12 years and over: take 4 to 8 chewable tablets every 4 hours not to exceed 48 chewable tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, dextrans hydrated, FD&C yellow #6 aluminum lake, flavor, maltodextrin, sucralose

Questions or comments?

1-800-426-9391

Principal Display Panel

QUALITY

+PLUS

NDC 50844-218-07

†Compare to active ingredient in
Bayer® Chewable Low Dose Aspirin

LOW DOSE

ASPIRIN

CHEWABLE

TABLETS 81 mg

PAIN RELIEVER

(NSAID)

**Chew or
crush tablets
completely
before
swallowing.**

- ASPIRIN REGIMEN
- ORANGE FLAVORED

ACTUAL SIZE

36

Chewable Tablets

Talk to your doctor or other healthcare provider
before using this product for your heart.

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

†This product is not manufactured or distributed
by Bayer AG, owner of the registered trademark
Bayer® Chewable Low Dose Aspirin.

50844 REV0122B21807

Distributed by

LNK INTERNATIONAL, INC.
 60 Arkay Drive, Hauppauge, NY 11788
 USA

*This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® Chewable Low Dose Aspirin.
 50844 REVO122B21807
 Distributed by
LNK INTERNATIONAL, INC.
 60 Arkay Drive, Hauppauge, NY 11788
 USA

QUALITY PLUS

LOW DOSE

ASPIRIN CHEWABLE TABLETS

81 mg

PAIN RELIEVER (NSAID)

36

Chewable Tablets

NDC 50844-218-07

¹Compare to active ingredient in Bayer® Chewable Low Dose Aspirin


Chew or crush tablets completely before swallowing.

• ASPIRIN REGIMEN
• ORANGE FLAVORED

Talk to your doctor or other healthcare provider before using this product for your heart.

ACTUAL SIZE

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING



3 5084421807 0

No Print Area
Lot no. & Exp. Date

Quality Plus 44-218

Drug Facts (continued)

Do not use

- have had stomach ulcers or bleeding problems
- if you are allergic to aspirin or any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - vomit blood
 - have bloody or black stools
 - feel faint
 - have stomach pain that does not get better
 - ringing in the ears or a loss of hearing occurs
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur

These could be signs of a serious condition.

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Drug Facts

Active ingredient
 (in each chewable tablet)
 Aspirin 81 mg (NSAID)Pain reliever
 *nonsteroidal anti-inflammatory drug

Purpose
 For the temporary relief of minor aches and pains or as recommended by your doctor

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take a blood thinner (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- are age 60 or older
- take more or for a longer time than directed

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- drink a full glass of water with each dose
- chew or crush tablets completely before swallowing; do not swallow tablets whole
- adults and children 12 years and over: take 4 to 8 chewable tablets every 4 hours not to exceed 48 chewable tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients corn starch, dextrates hydrated, FD&C yellow #6 aluminum lake, flavor, maltodextrin, sucralose

Questions or comments?
 1-800-426-9391

Drug Facts (continued)

- you experience any of the following signs of stomach bleeding:
 - vomit blood
 - have bloody or black stools
 - feel faint
 - have stomach pain that does not get better
 - ringing in the ears or a loss of hearing occurs
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur

These could be signs of a serious condition.

Quality Plus 44-218

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Drug Facts

Active ingredient
 (in each chewable tablet)
 Aspirin 81 mg (NSAID)Pain reliever
 *nonsteroidal anti-inflammatory drug

Purpose
 For the temporary relief of minor aches and pains or as recommended by your doctor

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take a blood thinner (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- are age 60 or older
- take more or for a longer time than directed

Quality Plus 44-218

Quality Plus 44-218

ASPIRIN LOW DOSE			
aspirin tablet, chewable			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50844-218

Route of Administration	ORAL
--------------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
DEXTROSE MONOHYDRATE (UNII: LX22YL083G)	
FD&C YELLOW NO. 6 ALUMINUM LAKE (UNII: GYP6Z2JR6Q)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	orange (light)	Score	no score
Shape	ROUND	Size	8mm
Flavor	ORANGE	Imprint Code	44;218
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50844-218-07	1 in 1 CARTON	09/12/1990	
1		36 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	09/12/1990	

Labeler - L.N.K. International, Inc. (038154464)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(50844-218)

Establishment

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

LNK International, Inc.		832867837	manufacture(50844-218)
-------------------------	--	-----------	------------------------

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(50844-218)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(50844-218)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(50844-218)

Revised: 11/2025

L.N.K. International, Inc.