

**ASPIRIN LOW DOSE- aspirin tablet, chewable**  
**L.N.K. International, Inc.**

-----  
**Quality Plus 44-218**

***Active ingredient (in each chewable tablet)***

Aspirin 81 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever

***Uses***

for the temporary relief of minor aches and pains or as recommended by your doctor

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- are age 60 or older
- take more or for a longer time than directed
- have had stomach ulcers or bleeding problems

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug for

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - vomit blood
  - have bloody or black stools
  - feel faint
  - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

***Directions***

- **do not take more than directed**
- drink a full glass of water with each dose
- chew or crush tablets completely before swallowing; do not swallow tablets whole
- adults and children 12 years and over: take 4 to 8 chewable tablets every 4 hours not to exceed 48 chewable tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

***Other information***

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

## ***Inactive ingredients***

corn starch, dextrans hydrated, FD&C yellow #6 aluminum lake, flavor, maltodextrin, sucralose

## ***Questions or comments?***

**1-800-426-9391**

## ***Principal Display Panel***

**QUALITY**

**+PLUS**

NDC 50844-218-07

†Compare to active ingredient in  
Bayer® Chewable Low Dose Aspirin

LOW DOSE

**ASPIRIN**

**CHEWABLE**

**TABLETS 81 mg**

PAIN RELIEVER

**(NSAID)**

**Chew or  
crush tablets  
completely  
before  
swallowing.**

- ASPIRIN REGIMEN
- ORANGE FLAVORED

ACTUAL SIZE

**36**

Chewable Tablets

Talk to your doctor or other healthcare provider  
before using this product for your heart.

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED  
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

†This product is not manufactured or distributed  
by Bayer AG, owner of the registered trademark  
Bayer® Chewable Low Dose Aspirin.

50844      REV0122B21807

Distributed by

**LNK INTERNATIONAL, INC.**  
 60 Arkay Drive, Hauppauge, NY 11788  
 USA

\*This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® Chewable Low Dose Aspirin.  
 50844 REVO122B21807  
 Distributed by  
**LNK INTERNATIONAL, INC.**  
 60 Arkay Drive, Hauppauge, NY 11788  
 USA

**QUALITY PLUS**

LOW DOSE

**ASPIRIN CHEWABLE TABLETS** 81 mg

PAIN RELIEVER (NSAID)

**36**  
Chewable Tablets

NDC 50844-218-07


<sup>1</sup>Compare to active ingredient in Bayer® Chewable Low Dose Aspirin

Chew or crush tablets completely before swallowing.

Talk to your doctor or other healthcare provider before using this product for your heart.

**ACTUAL SIZE**

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING



3 5084421807 0

**No Print Area  
Lot no. & Exp. Date**

**Drug Facts (continued)**

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not take more than directed
- drink a full glass of water with each dose
- chew or crush tablets completely before swallowing; do not swallow tablets whole
- adults and children 12 years and over: take 4 to 8 chewable tablets every 4 hours not to exceed 48 chewable tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

**Other information**

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

**Inactive ingredients** corn starch, dextrates hydrated, FD&C yellow #6 aluminum lake, flavor, maltodextrin, sucralose

**Questions or comments?**  
1-800-426-9391

**Drug Facts (continued)**

■ have had stomach ulcers or bleeding problems

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are** taking a prescription drug for

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - vomit blood
  - have bloody or black stools
  - feel faint
  - have stomach pain that does not get better
  - ringing in the ears or a loss of hearing occurs
  - pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - redness or swelling is present
  - new symptoms occur

These could be signs of a serious condition.

**KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION**

**Drug Facts**

**Active ingredient**  
(in each chewable tablet)  
Aspirin 81 mg (NSAID) .....Pain reliever  
\*nonsteroidal anti-inflammatory drug

**Purpose**  
for the temporary relief of minor aches and pains or as recommended by your doctor

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take a blood thinner (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- are age 60 or older
- take more or for a longer time than directed

**Drug Facts (continued)**

■ stomach bleeding warning applies to you

■ you have a history of stomach problems, such as heartburn

■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

■ you have asthma

■ you are taking a diuretic

**Ask a doctor or pharmacist before use if you are** taking a prescription drug for

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - vomit blood
  - have bloody or black stools
  - feel faint
  - have stomach pain that does not get better
  - ringing in the ears or a loss of hearing occurs
  - pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - redness or swelling is present
  - new symptoms occur

These could be signs of a serious condition.

B-1603-218A-07-R  
REVO122B21807

**Quality Plus 44-218**

**ASPIRIN LOW DOSE**

aspirin tablet, chewable

**Product Information**

**Product Type**

HUMAN OTC DRUG

**Item Code (Source)**

NDC:50844-218

<b>Route of Administration</b>	ORAL
--------------------------------	------

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>DEXTROSE MONOHYDRATE</b> (UNII: LX22YL083G)	
<b>FD&amp;C YELLOW NO. 6 ALUMINUM LAKE</b> (UNII: GYP6Z2JR6Q)	
<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	

### Product Characteristics

<b>Color</b>	orange (light)	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	8mm
<b>Flavor</b>	ORANGE	<b>Imprint Code</b>	44;218
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50844-218-07	1 in 1 CARTON	09/12/1990	
1		36 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	09/12/1990	

**Labeler** - L.N.K. International, Inc. (038154464)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(50844-218)

### Establishment

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

LNK International, Inc.		832867837	manufacture(50844-218)
-------------------------	--	-----------	------------------------

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(50844-218)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(50844-218)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(50844-218)

Revised: 11/2024

L.N.K. International, Inc.