

**RBUGR QUIT SMOKING NICOTINE- nicotine 18%, menthol 0.3% quit smoking
nicotine patch
Jiangxi Hemei Pharmaceutical Co., Ltd**

84010-153

Active Ingredient

Nicotine 18%
menthol 0.3%

Purpose

stop smoking aid

Use

reduces withdrawal symptoms,including nicotine craving,associated with quitting smoking

Warnings

If you are pregnant or breast-feeding, only use this patch on the advice of your health care provider.Smoking can seriously harm your child.

- Try to stop smoking without using any nicotine replacement patch,This patch is considered safer than smoking. However, the risks to your child from this patch are not fully known.
- The patch is effective in helping to alleviate nicotine cravings. It is a health supplement to assist in quitting smoking and is not to be used in place of a nicotine patch.

Do not use

If you are allergic to any of the ingredients.

- Gastric ulcers or diabetes mellitus, history of seizures prohibited.
- If you are under 18 years of age without a doctor's recommendation.
- Do not use on damaged skin or wounds.

When Using

if you have vivid dreams or other sleep disturbances, remove this patch at bedtime. · Stop use and ask a doctor if feeling uncomfortable or experiencing other sensitive dizziness symptoms. · And follow phase recommendations (being adjusted within the recommended range according to your condition and acceptability). · Do not cut the

7mg, patch into halves or smaller pieces, · To avoid possible burns—remove the patch before performing an MRI (Magnetic Resonance Imaging) examination. · Adjust the wearing time according to your body weight and your smoking needs (it is best to wear the patch for more than 12 hours in order to allow the nicotine in the patch to penetrate completely into the skin, and if you feel uncomfortable, you can use the intermittent patch method, i.e., temporarily remove the patch and put it back on once the discomfort subsides). · The used patch should be removed and a new one applied to a different skin site at the same time each day. · Slight dizziness and nausea during use are normal, please stop using temporarily and drink more water. · Reduce the dosage and use it again when it returns to normal. · It is important to complete treatment, if you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider. · It is best to remove hair from the surface of the skin and keep the skin dry before use for best results. · Drink plenty of water during use of the patch to promote absorption and metabolism

Stop Use

you are using a non - nicotine stop smoking drug. —Taking a prescription medicine for depression or asthma, your prescription dose may need to be adjusted. —Skin redness caused by the patch does not go away after 4 days or ·You develop a rash, swelling, or severe skin irritation. —You experience irregular heartbeat or palpitations. —Symptoms of nicotine overdose occur (such as nausea, vomiting, dizziness, or weakness). —It is important to complete treatment, if you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider.

Ask Doctor

Heart disease, recent heart attack, or irregular of heartbeat. · Cardiovascular disorders · History of myocardial infarction Documented arrhythmias · High blood pressure not controlled by medication, · Stomach ulcer or diabetes · Been prescribed medication for depression or asthma · Non-tobacco users or people with skin allergies.

Keep Out Of Reach Of Children

Please store in a cool dry place for best results. (preferably under 20-25°). Keep out of reach of children and pets. Please wrap used patches completely before throwing them away. If inadvertently licked by children or pets, get medical help or contact a Poison Control Center right away.

Directions

If you are under 18 years of age, ask a doctor before use, · Before using this product, read the enclosed self-help guide for complete directions and other information · Begin using the patch on your quit day. · If you smoke more than 10 cigarettes per day, use the following schedule below.

If you smoke no more than ten cigarettes a day, please start with step 2. Keep your skin clean and dry before each use. Immediately after removing the adhesive backing, apply the patch to the skin and press the patch against the skin for ten seconds to ensure a

snug fit. Wash your hands after each patch application and after removing the patch privately.

Inactive ingredients

acrylate adhesive, cellulose paper, methacrylic acid .

PRINCIPAL DISPLAY PANEL

★★★ RBUGR ★★★

QUIT SMOKING

Nicotine Patches

21 Patches

STEP 1+ TRANSDERMAL PATCH 28 mg

STEP 1 21 mg

STEP 2 14 mg

STEP 3 7 mg

Botanical-Derived Actives

Upgrade Detachable Patch

Helps reduce nicotine craving and withdrawal symptoms

PLEASE READ THE INSTRUCTIONS CAREFULLY BEFORE USE!

Manufacturer: Jiangsu Hemei Pharmaceutical Co., Ltd.
Address: 4th Floor, Building 1906, Phase II, Zhongguo Medical Incubation Park, Zhanggao, Yichun, Jiangsu 337208 China (CHN)

★★★ RBUGR ★★★

PLEASE READ THE INSTRUCTIONS CAREFULLY BEFORE USE!

DRUG FACTS

Active ingredients (in each patch) Purpose

Nicotine 18% Stop smoking aid

menthol 0.3% Stop smoking aid

USE: reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking.

WARNINGS

If you are pregnant or breast-feeding, only use this patch on the advice of your health care provider. Smoking can seriously harm your child. • Try to stop smoking without using any nicotine replacement patch. This product is considered safer than smoking. However, the risks to your child from this patch are not fully known. • The patch is effective in helping to alleviate nicotine cravings. It is a health supplement to assist in quitting smoking and is not to be used in place of a nicotine patch.

DO NOT USE

If you are allergic to any of the ingredients. • Gastric ulcers or diabetes mellitus, history of seizures prohibited. • If you are under 18 years of age without a doctor's recommendation. • Do not use on damaged skin or wounds.

ASK A DOCTOR BEFORE USE IF YOU HAVE

Heart disease, recent heart attack, or irregular heartbeat • Cardiovascular disorders • History of myocardial infarction Documented arrhythmias • High blood pressure not controlled by medication • Stomach ulcer or diabetes • Been prescribed medication for depression or asthma • Non-tobacco users or people with skin allergies.

STOP USE AND ASK A DOCTOR IF

you are using a non-nicotine stop smoking drug. • Taking a prescription medicine for depression or asthma, your prescription dose may need to be adjusted. • Skin redness caused by the patch does not go away after 4 days. • You develop a rash, swelling, or severe skin irritation. • You experience irregular heartbeat or palpitations. • Symptoms of nicotine overdose occur (such as nausea, vomiting, dizziness, or weakness). • It is important to complete treatment, if you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider.

WHEN USING THE PATCH

If you have vivid dreams or other sleep disturbances, remove this patch at bedtime. • Stop use and ask a doctor if feeling uncomfortable or experiencing other sensitive dizziness symptoms. • And follow phase recommendations (being adjusted within the recommended range according to your condition and acceptability). • Do not cut the 7 mg patch into halves or smaller pieces. • To avoid possible burns, remove the patch before performing an MRI (Magnetic Resonance Imaging) examination. • Adjust the wearing time according to your body weight and your smoking needs (it is best to wear the patch for more than 12 hours in order to allow the nicotine in the patch to penetrate completely into the skin, and if you feel uncomfortable, you can use the intermittent patch method, i.e., temporarily remove the patch and put it back on once the discomfort subsides). • The used patch should be removed and a new one applied to a different skin site at the same time each day. • Slight dizziness and nausea during use are normal, please stop using temporarily and drink more water. • Reduce the dosage and use it again when it returns to normal. • It is important to complete treatment, if you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider. • It is best to remove hair from the surface of the skin and keep the skin dry before use for best results. • Drink plenty of water during use of the patch to promote absorption and metabolism.

DIRECTIONS

If you are under 18 years of age, ask a doctor before use. • Before using this product, read the enclosed self-help guide for complete directions and other information. • Begin using the patch on your quit day. • If you smoke more than 10 cigarettes per day, use the following schedule below.

| STEP 1+ | STEP 1 | STEP 2 | STEP 3 |
|------------|------------|----------|-----------|
| Week 1-2 | Week 3-6 | Week 7-8 | Week 9-10 |
| 2 patches | 1½ patches | 1 patch | ¾ patch |
| 1½ patches | 1 patch | ¾ patch | ½ patch |

If you smoke no more than ten cigarettes a day, please start with step 2. Keep your skin clean and dry before each use. Immediately after removing the adhesive backing, apply the patch to the skin and press the patch against the skin for ten seconds to ensure a snug fit. Wash your hands after each patch application and after removing the patch privately.

KEEP OUT OF REACH OF

Please store in a cool dry place for best results. (preferably under 20-25°). Keep out of reach of children and pets. Please wrap used patches completely before throwing them away. If inadvertently taken by children or pets, get medical help or contact a Poison Control Center right away.

INACTIVE INGREDIENTS

acrylate adhesive, cellulose paper, methacrylic acid.

MFD EXP

RBUGR QUIT SMOKING NICOTINE

nicotine 18%, menthol 0.3% quit smoking nicotine patch

| Product Information | | | |
|--|----------------|--------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:84010-153 |
| Route of Administration | TOPICAL | | |
| Active Ingredient/Active Moiety | | | |
| Ingredient Name | | Basis of Strength | Strength |
| MENTHOL (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A) | | MENTHOL | 0.3 g in 100 |

NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)

NICOTINE

18 g in 100

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| METHACRYLIC ACID (UNII: 1CS02G8656) | |
| 2-HYDROXYETHYL ACRYLATE (UNII: 25GT92NY0C) | |
| POWDERED CELLULOSE (UNII: SMD1X3XO9M) | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:84010-153-01 | 21 in 1 BOX; Type 0: Not a Combination Product | 07/16/2025 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M017 | 07/16/2025 | |

Labeler - Jiangxi Hemei Pharmaceutical Co., Ltd (724892056)

Establishment

| Name | Address | ID/FEI | Business Operations |
|---------------------------------------|---------|-----------|------------------------|
| Jiangxi Hemei Pharmaceutical Co., Ltd | | 724892056 | manufacture(84010-153) |

Revised: 7/2025

Jiangxi Hemei Pharmaceutical Co., Ltd