

SATOHAP- methyl salicylate, dl-camphor, l-menthol spray
Sato Pharmaceutical Co., Ltd.

Satohap Spray

Active ingredients

dl-Camphor 5%

l-Menthol 4%

Methyl salicylate 10%

Purpose

dl-Camphor External analgesic

l-Menthol External analgesic

Methyl salicylate External analgesic

Uses temporary relieves minor aches and pains of muscles and joints due to
■ backache ■ arthritis ■ strains ■ bruises ■ sprains

Warnings

For external use only

Do not use on wounds or on damaged skin

When using this product

■ avoid contact with the eyes

■ do not bandage tightly

Stop use and ask a doctor if

■ symptoms persist for more than 7 days

■ symptoms clear up and occur again within a few days

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Warnings-contents under pressure

■ do not puncture or incinerate container.

■ do not expose to heat or store at temperatures above 120 degrees F.

Directions

■ adults and children 2 years of age and over: Hold can upright and spray until wet.
Apply to affected area not more than 3 to 4 times daily.

■ children under 2 years of age: Do not use and ask a doctor.

Inactive ingredients

fragrance, phenylethyl alcohol, alcohol, propylene glycol, LPG

NDC 49873-068-02
sato

打撲・ねんざ・肩こりに
SATOHAP®
SPRAY
 EXTERNAL ANALGESIC SPRAY

サトウハップ
 スプレー
 外用鎮痛剤

NO CFC's

2.37 FL OZ
 (70 mL)

Drug Facts

Active ingredients Purpose
 dl-Camphor 5%.....External analgesic
 l-Menthol 4%.....External analgesic
 Methyl salicylate 10%...External analgesic

Uses temporarily relieves minor aches and pains of muscles and joints due to
 ■ backache ■ arthritis ■ strains
 ■ bruises ■ sprains

Warnings

For external use only

Do not use on wounds or damaged skin

When using this product

- avoid contact with the eyes
- do not bandage tightly

Stop use and ask a doctor if

- condition worsens
- symptoms persist for more than 7 days
- symptoms clear up and occur again within a few days

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Warnings-contents under pressure

- do not puncture or incinerate container.
- do not expose to heat or store at temperatures above 120 degrees F.

Directions

- adults and children 2 years of age and over: Hold can upright and spray until wet. Apply to affected area not more than 3 to 4 times daily.
- children under 2 years of age: Do not use and ask a doctor.

Inactive ingredients

fragrance, phenylethyl alcohol, alcohol, propylene glycol, LPG

医薬品表示

有効成分・分量 配合目的
 dl-カンフル 5%.....外用消炎鎮痛剤
 l-メントール 4%.....外用消炎鎮痛剤
 サリチル酸メチル 10%...外用消炎鎮痛剤

効 能 下記に起因する筋肉痛及び関節痛の一時的緩和。

- 腰痛 ■ 関節炎 ■ 筋違え
- 打撲 ■ ねんざ

使用上の注意

外用にのみご使用ください。

傷口及び損傷した皮膚には使用しないでください。

使用の際注意してください。

- 目に入らないようご注意ください。
- きつく包帯しないでください。

使用を中止し、医師にご相談ください。

- 症状が悪化した場合
- 7日以上症状の改善が見られない場合
- 一度改善した症状が数日以内に再発した場合

小児の手の届かない所に保管してください。誤って飲み込んだ場合には、直ちに専門家又はボイズンコントロールセンターへご相談ください。

高圧製品に対する注意

- 容器を壊したり、熱やしたりしないでください。
- 熱にさらしたり120°F以上の場所での保管はしないでください。

用法・用量

- 大人及び2才以上の小児: 容器を傾けないようにして患部がぬれるまで噴霧してください。1日3~4回を限度として患部に噴霧します。
- 2才未満の小児: 使用せず、医師にご相談ください。

その他の成分 香料、フェニルエチルアルコール、アルコール、プロピレングリコール、液化石油ガス

EXP. DATE (使用期限), LOT (製造記号):

See bottom (底面に記載)

US-D

Manufactured by

SATO PHARMACEUTICAL CO., LTD.

1-5-27 MOTOAKASAKA MINATO-KU TOKYO, JAPAN

Reports of serious side effects associated with use of this product can be sent to:

SATO PHARMACEUTICAL, INC

1650 E. Glenn Curtiss St., Unit 2, Carson, CA 90746

製造元

佐藤製薬株式会社

日本国東京都港区元赤坂1丁目5番27号



3

SATOHAP

methyl salicylate, dl-camphor, l-menthol spray

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49873-068
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
METHYL SALICYLATE (UNII: LAV5U5022Y) (SALICYLIC ACID - UNII:O414PZ4LPZ)	METHYL SALICYLATE	10 g in 100 mL
LEVOMENTHOL (UNII: BZ1R15MTK7) (LEVOMENTHOL - UNII:BZ1R15MTK7)	LEVOMENTHOL	4 g in 100 mL
CAMPHOR (SYNTHETIC) (UNII: 5TJD82A1ET) (CAMPHOR (SYNTHETIC) - UNII:5TJD82A1ET)	CAMPHOR (SYNTHETIC)	5 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
PHENYLETHYL ALCOHOL (UNII: ML9LGA7468)	
ALCOHOL (UNII: 3K9958V90M)	
LIQUEFIED PETROLEUM GAS (UNII: 5K616HU99V)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49873-068-02	70 mL in 1 CANISTER; Type 0: Not a Combination Product	11/03/2006	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	11/03/2006	

Labeler - Sato Pharmaceutical Co., Ltd. (690575642)

Establishment

Name	Address	ID/FEI	Business Operations
Daizo Corporation		695312298	pack(49873-068) , label(49873-068)

Establishment

Name	Address	ID/FEI	Business Operations
Sato Pharmaceutical Co., Ltd.		715699133	manufacture(49873-068)

