

**ASPIRIN LOW DOSE- aspirin tablet**  
**NUVICARE LLC**

-----  
**Aspirin Pain reliever**

**Active Ingredient (in each tablet)**

Aspirin 81mg (NSAID)\*

\*Nonsteroidal Anti-Inflammatory Drug

**Purpose**

Pain reliever

**Uses:**

- for the temporary relief of minor aches and pains or as recommended by your doctor.  
**Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief**
- ask your doctor about other uses for Safety Coated 81 mg Aspirin

**Warnings:**

**Reye's syndrome**

Children and teenagers who have or are recovering from chickenpox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert**

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning**

This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug for • gout • diabetes • arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - pain gets worse or lasts for more than 10 days
  - Fever gets worse or lasts more than 3 days
  - redness or swelling is present
  - new symptoms occur
  - ringing in the ears or loss of hearing occurs

**If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**KEEP OUT OF REACH OF CHILDREN.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**DIRECTIONS**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor

- children under 12 years: consult a doctor

## OTHER INFORMATION

- store at room temperature. Avoid excessive heat above 40°C (104°F).

## INACTIVE INGREDIENTS

Microcrystalline Cellulose, Lactose, Croscarmellose Sodium, Colloidal Silicon Dioxide, Acrylcoat L 30 (Methacrylic Acid copolymer), Triethyl citrate, Polysorbate 80, Sodium hydroxide, Talc, Titanium Dioxide, D & C yellow # 10 Aluminum lake, yellow oxide of iron.

## QUESTIONS OR COMMENTS?

Call **1 (718) 337-8733** or visit [support@nuvicare.com](mailto:support@nuvicare.com)



**Aspirin Low Dose**  
Aspirin 81 mg each tablet

✓ Safety Coated Aspirin Regimen

✓ Talk to your doctor or other healthcare provider before using this product for your heart.

**300**  
caplets

not actual size

Compare to  
**BAYER®**  
**Low Dose Aspirin**  
active ingredient



**Drug Facts**

**Active ingredient (in each tablet)**  
Aspirin 81 mg (NSAID)\* ..... Pain reliever  
\*nonsteroidal anti-inflammatory drug

**Uses:** ■ for the temporary relief of minor aches and pains or as recommended by your doctor. **Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.** ■ ask your doctor about other uses for Safety Coated 81 mg Aspirin

**Warnings: Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had

(continued under label)

†This product is not manufactured or distributed by Bayer Healthcare, LLC owner of the Registered Trademark BAYER® Low Dose Aspirin Tablets

**Distributed by:** Nuvicare LLC, Brooklyn, NY 11211, U.S.A.  
[www.nuvicare.com](http://www.nuvicare.com)  
**PL0222-00**



3 84324 00003 9

Lot No. Exp. Dt.

**PEEL HERE**

**Drug Facts** *(continued)*

stomach ulcers or bleeding problems ■ take a blood thinner (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Do not use** ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if** ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma

**Ask a doctor or pharmacist before use if you are** taking a prescription drug for ■ gout ■ diabetes ■ arthritis

**Stop use and ask a doctor if** ■ an allergic reaction occurs. Seek medical help right away ■ you experience any of the following signs of stomach bleeding: • feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ ringing in the ears or a loss of hearing occurs

**If pregnant or breast-feeding**, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.** **KEEP OUT OF REACH OF CHILDREN.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions:** ■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor ■ children under 12 years: consult a doctor

**Other information:** ■ store at room temperature. Avoid excessive heat above 40°C (104°F).

**Other ingredients:** Microcrystalline Cellulose (PH-102), Lactose, Croscarmellose Sodium, Colloidal Silicon Dioxide, Acrylcoat L 30 DIMethacrylic acid copolymer], triethyl citrate, Polysorbate 80, Sodium hydroxide, Talc, Titanium dioxide, D & C yellow#10 Aluminum lake, Yellow oxide of iron

**Questions or comments?**  
Call 1 (718) 337-8733 or visit: [support@nuvicare.com](mailto:support@nuvicare.com)

## ASPIRIN LOW DOSE

aspirin tablet

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:84324-017
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>MICROCRYSTALLINE CELLULOSE 102</b> (UNII: PNR0YF693Y)	
<b>LACTOSE, UNSPECIFIED FORM</b> (UNII: J2B2A4N98G)	
<b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER</b> (UNII: NX76LV5T8J)	
<b>TRIETHYL CITRATE</b> (UNII: 8Z96QXD6UM)	
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>SODIUM HYDROXIDE</b> (UNII: 55X04QC32I)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>D&amp;C YELLOW NO. 10 ALUMINUM LAKE</b> (UNII: CQ3XH3DET6)	
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)	

## Product Characteristics

<b>Color</b>	yellow	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	8mm
<b>Flavor</b>		<b>Imprint Code</b>	81
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84324-017-01	300 in 1 BOTTLE; Type 0: Not a Combination Product	08/01/2024	

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M013	08/01/2024	

**Labeler** - NUVICARE LLC (119257565)

