MUCOSA DM- dextromethorphan hydrobromide, guaifenes in tablet Magno-Humphries, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Mucosa DM

MHL NDC 54257-277-01 immediate Acting

Mucosa DM

EXPECTORANT/COUGH SUPPRESSANT

- Soothes Cough
- Loosens Congestion

Guaifenesin 400 mg
Dextromethorphan HBr 20 mg
50 TABLETS

DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGE

Active ingredients (in each tablet)

Dextromethorphan HBr 20mg Guaifenesin 400 mg

Purpose

Dextromethorphan HBr 20mg......Cough Suppressant Guaifenesin 400 mg.....Expectorant

Uses

helps loosen phlegm (mucus), helps thin bronchial secretions to make coughs more productive, temporarily relieves cough due to minor throat and bronchial irritation associated with the common cold.

Warnings

Do not use for children under 12. Do not use if you are now taking Da prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for two weeks after stopping the MAOI drug. If you do not know if your prescription drug contains MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have Cougha companied with excessive phlegm (mucus), persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema.

When using this product do not exceed recommended dosage do not use for more than 7 days.

Stop use and ask a doctor if cough lasts for more than 7 days, recurs, or is accompanied with fever, rash, or persistent headache. These could be signs of a serious condition.

IIf pregnant or breast-feeding, <code>lask</code> a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

adults and children 12 years & older - take one tablet every 4 hours with a full glass of water while symptoms persist. Do not exceed 6 doses in 24 hours.

children under 12 years of age - do not use

Other information

• store at controlled room temperature 15° to 30°C (59° to 86°F) Keep in a dry place and do not expose to excessive heat.

Inactive ingredients

Croscarmellose sodium, maltodextrin, microcrystalline cellulose, povidone, silicon dioxxide, stearic acid.

Questions?

call toll-free **1-800-935-6737**

Distributed by:

Magno-Humphries, Inc., OR 97223 U.S.A.

Visit our website:

| www.magno-humphries.com

Distributed by:

Magno-Humphries, Inc. Tigard, OR 97223 U.S.A.

Visit our website:

lwww.magno-humphries.com

Package Labeling:





Drug Facts (cont'd)

Directions

children under 12 years of age | do not use

Other information

 store at controlled room temperature 15° to 30°C (59° to 86°F) Keep in a dry place and do not expose to excessive heat.

Inactive ingredients
Croscarmellose sodium, maltodextrin, microcrystalline
cellulose, povidone, silicon dioxide, stearic acid.

Questions? call toll-free 1-800-935-6737

Distributed by: Magno-Humphries Labs Tigard, OR 97223 U.S.A.

Visit our website: www.magno-humphries.com bronchial secretions to make coughs more productive, temporarily relieves cough due to minor throat and bronchial irritation associated with the common cold.

Warnings Do not use for children under 12.

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for two weeks after stopping the MAOI drug. If you do not know if your prescription drug contains MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have cough accompanied with excessive phlegm (mucus), persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema.

When using this product do not exceed recommended dosage do not use for more than 7 days.

Stop use and ask a doctor if cough lasts for more than 7 days, recurs, or is accompanied with fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

adults and children 12 years & older take one tablet every 4 hours with a full glass of water while symptoms persist. Do not exceed 6 doses in 24 hours.



■ PEEL FOR DIRECTIONS G7752-050-01-1

MUCOSA DM

dextromethorphan hydrobromide, guaifenesin tablet

Product Information

Product TypeHUMAN OTC DRUGItem Code (Source)NDC:54257-277

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9 D2RTI9 KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg

Inactive Ingredients	
Ingredient Name	Strength
CROSCARMELLOSE SODIUM (UNII: M28 O L 1HH48)	

MALTO DEXTRIN (UNII: 7CVR7L4A2D)	
MICRO CRYSTALLINE CELLULO SE (UNII: OP1R32D61U)	
PO VIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SILICON DIO XIDE (UNII: ETJ7Z6 XBU4)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics	duct Characteristics			
Color	white	Score	2 pieces	
Shape	OVAL	Size	17mm	
Flavor		Imprint Code	AZ345	
Contains				

	Packaging				
ı	# Item Code	Package Description	Marketing Start Date	Marketing End Date	
	1 NDC:54257-277-01	50 in 1 BOTTLE; Type 0: Not a Combination Product	04/04/2017		

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph final	part341	04/04/2017		

Labeler - Magno-Humphries, Inc. (063251433)

Revised: 12/2017 Magno-Humphries, Inc.