EQUATE LANSOPRAZOLE- lansoprazole capsule, delayed release WALMART INC.

Wal-Mart Lansoprazole Drug Facts

Active ingredient (in each capsule)

Lansoprazole 15 mg

Purpose

Acid Reducer

Use

- treats frequent heartburn (occurs <u>2 or more</u> days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert:

- do not use if you are allergic to lansoprazole
- lansoprazole may cause severe skin reactions. Symptoms may include:
- skin reddening
- blisters
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting

stomach pain

Ask a doctor or pharmacist before use if you are

• taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding,

ask a health care professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information

read the directions and warnings before use

- keep the carton. It contains important information.
- store at 20-25°C (68-77°F)
- keep product out of high heat and humidity
- · protect product from moisture
- close cap tightly after use

Inactive ingredients

D&C red no. 28, D&C yellow no. 10, FD&C blue no. 1, FD&C red no. 40, gelatin, hypromellose, low substituted hydroxypropyl cellulose, mannitol, meglumine, methacrylic acid copolymer, pharmaceutical ink, polyethylene glycol, polysorbate 80, sodium lauryl sulfate, sugar spheres, talc, titanium dioxide

Questions or comments?

1-888-287-1915

Package/Label Principal Display Panel

3 PACK

THREE 14-DAY COURSES OF TREATMENT

equate™

Compare to Prevacid® 24 HR active ingredient

Lansoprazole Delayed-Release Capsules, 15 mg

Acid Reducer

24 HOUR

Treats Frequent Heartburn

- May take 1 to 4 days for full effect
- Sodium-Free

Actual Size


15 mg

42 CAPSULES

Three 14-day courses of treatment

3 Bottles Inside





Drug Facts

Active ingredient (in each capsule)

Purpose Acid Reducer

La nso prazole 15 mg.....

- treats frequent heartbum (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

- Allergy alert: do not use if you are allergic to lansoprazole
- lansoprazole may cause severe skin reactions. Symptoms may include:
- skin reddening blisters rash
- If an allergic reaction occurs, stop use and seek medical help right away.

Do not use if you have:

- trouble or pain swallowing food, vo miting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chestpain

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have ■ liver disease

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with hear burn
- unexplained weight loss nause a or vomiting stomach pain

Ask a doctor or pharmacist before use if you are

■ taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if your heartburn continues or worsens

- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea you develop a rash orjoint pain

If pregnant or breast-feeding, ask a health care professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours 14-Day Course of Treatment
 - swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: a ska doc to r be fore use. Heartburn in children may sometimes be caused by a serious condition.

Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause hear tourn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcoholand even som e acidic fruits and vegetables.
- ■Eatslowly and do not eatbig meals.
- Do note at late at night or justbefore bedtime.
- Do not lie flator bend over soon after eating.
- ■Raise the head of your bed.
- We ar loose-fitting clothing around your stomach.
- ■If you are overweight, lose weight
- If you smoke, quit smoking.

KEEP THE CARTON. IT CONTAINS IMPORTANT INFORMATION.

DO NOT USE IF PRINTED SEAL UNDER CAPIS BROKEN OR MISSING OR BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.

Satisfaction guaranteed Or we'll replace it or give you your money back. For questions or comments or to report an undesired reaction or side effect, please call 1-888-287-1915.

DISTRIBUTED BY: Walmart Inc., Bentonville, AR 72716

*This product is not manufactured or distributed by Takeda Pharmaceuticals U.S.A., Inc., owner of the registered trademark Prevacid®.



Drug Facts (continued)

protect product from moisture

Other information

- read the directions and warnings before use keep the car to n. It contains important information.
- store at 20-25°C(68-77°F)
 - keep product out of high heat and humidity
 - close cap tightly after use

Inactive ingredients D&C red no. 28, D&C yellow no. 10, FD&C blue no. 1, FD&C red no. 40, gelatin, hyprome lose, low substituted hydroxypropyl cellulose, mannibl, meglumine, methacrylic acid copolymer, pharmace utical ink, polyethylene glycol, polysorbate 80, sodium lauryl sulfate, sugar spheres, talc, titanium dioxide

Questions or comments?1-888-287-1915

EQUATE LANSOPRAZOLE

lansoprazole capsule, delayed release

Product Information Product Type HUMAN OTC DRUG Item Code (Source) NDC:79903-068 **Route of Administration ORAL**

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
LANSOPRAZOLE (UNII: 0K5C5T2QPG) (LANSOPRAZOLE - UNII:0K5C5T2QPG)	LANSOPRAZOLE	15 mg	

Inactive Ingredients	
Ingredient Name	Strength
D&C RED NO. 28 (UNII: 767IP0Y5NH)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
LOW-SUBSTITUTED HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 2165RE0K14)	
MANNITOL (UNII: 30WL53L36A)	
MEGLUMINE (UNII: 6HG8UB2MUY)	
METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYSORBATE 80 (UNII: 60ZP39ZG8H)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics				
Color	PINK, GREEN	Score	no score	
Shape	CAPSULE	Size	15mm	
Flavor		Imprint Code	24HR	
Contains				

Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:79903-068- 01	1 in 1 CARTON	07/20/2021		
1		14 in 1 BOTTLE; Type 0: Not a Combination Product			
2	NDC:79903-068- 03	3 in 1 CARTON	08/04/2021		
2		14 in 1 BOTTLE; Type 0: Not a Combination Product			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA202319	07/20/2021	

Labeler - WALMART INC. (051957769)

Revised: 3/2025 WALMART INC.