AGE DEFYING MINERAL PROTECTION- titanium dioxide, zinc oxide cream MARK SCHWARTZ CLINICAL SKINCARE, INC.

Active Ingredients:

Titanium Dioxide 8% Zinc Oxide 3.8%

Purpose:

Sunscreen

Uses:

Helps Prevent Sunburn. If used as directed with other sun protection measures (see "Directions"), decreases the risk of skin cancer and early skin aging caused by the sun.

Warnings:

For external use only. Do not use on broken or damaged skin. Stop use and ask a doctor if rash occurs. When using this product, keep out of eyes. Rinse with water to remove. Keep out of reach of children. If swallowed, get medical help or contact a poison control center right away.

Warnings:

Keep out of reach of children

Directions:

Apply liberally 15 minutes before sun exposure. Use a water-resistant product if swimming or sweating. Reapply at least every 2 hours. Children under 6 months: Ask a doctor. Sun Protection Measures: Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad-Spectrum SPF value of 15 or higher and other sun protection measures including: Limit time in the sun, especially from 10 a.m. - 2 p.m. Wear long-sleeved shirts, pants, hats and sunglasses.

Inactive Ingredients:

Alumina, Cyclohexasiloxane, Cyclopentasiloxane, Dimethicone, Dimethicone Crosspolymer, Dimethicone/Vinyl Dimethicone Crosspolymer, Dimethiconol, Lauryl PEG/PPG-18/18 Methicone, Hydrogen Dimethicone, PEG-10 Dimethicone, Tetrahexyldecyl Ascorbate, Tocopheryl Acetate

Other Information:

- Protect this product from excessive heat and direct sun
- PA++++

DRUG FACTS (cont)

- Sun Protection Measures: Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:
 - Limit time in the sun, especially from 10 am-2 pm
 - Wear long-sleeved shirts, pants, hats, and sunglasses

INACTIVE INGREDIENTS:

Alumina, Cyclohexasiloxane, Cyclopentasiloxane, Dimethicone, Dimethicone Crosspolymer, Dimethicone/Vinyl Dimethicone Crosspolymer, Dimethiconol, Lauryl PEG/PPG-18/18 Methicone, Hydrogen Dimethicone, PEG-10 Dimethicone, Tetrahexyldecyl Ascorbate, Tocopheryl Acetate

OTHER INFORMATION:

- Protect this product from excessive heat and direct sun
 - PA ++++

MANUFACTURED FOR: MARK SCHWARTZ CLINICAL SKINCARE INC. NY, NY 10075 Made in USA www.schwartzmdskincare.com

CLINICALSKINCARE





SUNSCREEN

AGE DEFYING MINERAL PROTECTION

CLEAR ZINC & TITANIUM BROAD SPECTRUM SPF 40 WATER RESISTANT (80 MINUTES)

DRUG FACTS

ACTIVE INGREDIENTS: Titanium Dioxide 8% Zinc Oxide 3.8% PURPOSE: Sunscreen Sunscreen

USES: Helps prevent sunburn. If used as directed with other sun protection measures (see directions), decreases the risk of skin cancer and early skin aging caused by the sun.

WARNINGS: For external use only. Do not use on damaged or broken skin. Stop use and ask a doctor if rash occurs. When using this product, keep out of eyes. Rinse with water to remove. Keep out of the reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

DIRECTIONS: Apply liberally 15 minutes before sun exposure.

- Reapply:
 - After 80 minutes of swimming or sweating
 - Immediately after towel drying
 At least every 2 hours
- Children under 6 months: Ask a doctor



Net Wt. 1.8 OZ / 53 G

AGE DEFYING MINERAL PROTECTION

titanium dioxide, zinc oxide cream

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:85715-202

Route of Administration TOPICAL

Active Ingredient/Active Moiety

,			
Ingredient Name	Basis of Strength	Strength	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	80 mg in 1 g	
ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	38 mg in 1 g	

Inactive Ingredients

Ingredient Name	Strength
ALUMINA (UNII: LMI26O6933)	
HYDROGEN DIMETHICONE (20 CST) (UNII: 12Z59IF64N)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
CYCLOPENTASILOXANE (UNII: 0THT5PCI0R)	
LAURYL PEG/PPG-18/18 METHICONE (UNII: ZJ5S27D9NX)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
ALPHA-TOCOPHEROL ACETATE (UNII: 9E8X80D2L0)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
DIMETHICONE CROSSPOLYMER (UNII: UF7620L1W6)	
TETRAHEXYLDECYL ASCORBATE (UNII: 9LBV3F07AZ)	
CYCLOHEXASILOXANE (UNII: XHK3U310BA)	
DIMETHICONOL (2000 CST) (UNII: T74O12AN6Y)	

Packaging			
# Item Code	Package Description	Marketing Start Date	Marketing End Date
1 NDC:85715- 202-50	53 g in 1 BOTTLE, PUMP; Type 0: Not a Combination Product	12/06/2021	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M020	12/06/2021	

Labeler - MARK SCHWARTZ CLINICAL SKINCARE, INC. (115548819)

Establishment				
Name	Address	ID/FEI	Business Operations	
Fragrance Manufacturing Inc.		793406000	manufacture(85715-202)	

Establishment				
Name	Address	ID/FEI	Business Operations	
Custom Analytics LLC		144949372	analysis (85715-202)	

Revised: 7/2025 MARK SCHWARTZ CLINICAL SKINCARE, INC.