

ASCURIN FAST ACTION- aspirin powder
HealthDoc LLC

AsCurin Fast Action

Drug Facts

Active ingredient (in each sachet)

Aspirin (NSAID*) 81mg

*nonsteroidal anti-inflammatory drug

Purpose

Pain Reliever-Fever Reducer

Use

- For the temporary relief of minor aches and pains and to reduce fever.

Warnings

Reye's syndrome:Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert:

Aspirin may cause a severe allergic reaction which may include:

- Hives
- Facial swelling
- Asthma (wheezing)
- Shock

Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout, or arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - any new symptoms occur
 - ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- Adults: Drink a full glass of water with each dose.
- Adults and children 12 years and over: Take 5-8 sachets every 4 hours not to exceed 49 sachets in 24 hours unless directed by a doctor.
- Children under 12 years of age: consult a doctor.
- See your doctor for other uses of aspirin, but do not use for more than 10 days without consulting your doctor because serious side effects may occur.

Other information

- Store at 77°F (25°C)

■ Excursions permitted between 59-86°F (15-30°C)

Inactive ingredients

Citric Acid, Curcumin, Malic Acid, Neotame, Orange Oil, Polyvinylpyrrolidone K30, Sorbitol, Sucralose.

Package Labeling:

**HealthDoc**
AsCurin
FAST ACTION
Soft Taste, Hard on Pain and Fever

Drug Facts
Active ingredient (in each sachet) Purpose
Aspirin (NSAID) 81mg Pain Reliever-Fever Reducer
Nonsteroidal anti-inflammatory drug

Use
■ For the temporary relief of minor aches and pains and to reduce fever

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction which may include:
■ Hives
■ Facial swelling
■ Asthma (wheezing)
■ Shock
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older
■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed
Do not use
■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
■ for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor
Ask a doctor before use if
■ stomach bleeding warning applies to you
■ you have a history of stomach problems, such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

Drug Facts (continued)
■ you are taking a diuretic
■ you have asthma
Ask a doctor or pharmacist before use if you are
■ taking a prescription drug for diabetes, gout, or arthritis
Stop use and ask a doctor if
■ an allergic reaction occurs. Seek medical help right away.
■ you experience any of the following signs of stomach bleeding:
■ feel faint
■ vomit blood
■ have bloody or black stools
■ have stomach pain that does not get better
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present
■ any new symptoms occur
■ ringing in the ears or loss of hearing occurs
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions
■ Adults: Drink a full glass of water with each dose.
■ Adults and children 12 years and over: Take 5-8 sachets every 4 hours not to exceed 48 sachets in 24 hours unless directed by a doctor.
■ Children under 12 years of age: consult a doctor.
■ See your doctor for other uses of aspirin, but do not use for more than 10 days without consulting your doctor because serious side effects may occur.
Other information
■ Store at 77°F (25°C)
■ Excursions permitted between 59-86°F (15-30°C)
Inactive ingredients
Citric Acid, Curcumin, Malic Acid, Neotame, Orange Oil, Polyvinylpyrrolidone K30, Sorbitol, Sucralose

No Sodium
No Artificial color
No Artificial flavor
No Carbonate

**HealthDoc**
AsCurin
FAST ACTION
Adult's Aspirin (NSAID)
Pain Reliever-Fever Reducer
Powder Sachets

Soft Taste, Hard on Pain and Fever

Dissolve in mouth or mix with water

28 – 0.07 oz (2 g) Sachets
Net Wt. 2 oz (56 g)


Distributed by: HealthDoc LLC
1004 Industrial Dr. Ste 3
Berlin Township, NJ 08091
United States

How to use?

Shake contents down
Tear at Arrow

Pour on tongue
Follow with water

Mix Water & Drink



**Adult's Aspirin (NSAID)
Pain Reliever-Fever Reducer
Powder Sachets**

Soft Taste, Hard on Pain and Fever
Dissolve in mouth or mix with water

Net Wt. 0.07 oz (2 g)

Aspirin (NSAID) 81mg

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed

How to Use?

- Shake contents down and tear at arrow.
- Mix with water & drink.
- Or pour on tongue and follow with water.

Distributed by: HealthDoc LLC
1004 Industrial Dr. Ste 3
Berlin Township, NJ 08091
United States

This Unit Not Labeled For Retail Sale.

Please refer to outer container for complete Drug Facts information.

ASCURIN FAST ACTION

aspirin powder

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:83607-001
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg in 2 g

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
CURCUMIN (UNII: IT942ZTH98)	
MALIC ACID (UNII: 817L1N4CKP)	
NEOTAME (UNII: VJ597D52EX)	
ORANGE OIL (UNII: AKN3KSD11B)	
POVIDONE K30 (UNII: U725QWY32X)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83607-001-00	28 in 1 BOX	10/23/2023	
1		2 g in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	10/23/2023	

Labeler - HealthDoc LLC (013084196)

Revised: 10/2023

HealthDoc LLC