

STOOL SOFTENER PLUS STIMULANT LAXATIVE- docusate sodium and sennosides tablet

TOP CARE (Topco Associates LLC)

Drug Facts

Active ingredients (in each tablet)

Docusate sodium 50 mg

Sennosides 8.6 mg

Purpose

Stool softener

Stimulant laxative

Uses

- for overnight relief from occasional constipation (irregularity)
- generally produces bowel movement in 6 to 12 hours

Warnings

Do not use

- laxative products for longer than 1 week unless directed by a doctor
- if you are presently taking mineral oil, unless directed by a doctor

Ask a doctor before use if you have

- stomach pain
- nausea
- vomiting
- noticed a sudden change in bowel habits that lasts over a period of 2 weeks

Stop use and ask a doctor if

you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- Take only by mouth. Doses may be taken as a single daily dose, preferably in the evening, or in divided doses.

adults and children 12 years and older	take 2-4 tablets daily
children 6 to under 12 years of age	take 1-2 tablets daily
children 2 to under 6 years of age	take up to 1 tablet daily
children under 2	ask a doctor

Other information

- each tablet contains: **calcium 20 mg**
- each tablet contains: **sodium 6 mg** VERY LOW SODIUM
- store at 25°C (77°F); excursions permitted between 15-30°C (59-86°F)

Inactive ingredients

carnauba wax*, croscarmellose sodium, dibasic calcium phosphate dihydrate, FD&C blue #2 aluminum lake, FD&C red #40 aluminum lake, hypromellose, magnesium stearate, maltodextrin*, microcrystalline cellulose, polyethylene glyco*, polyvinyl alcohol*, silicon dioxide, sodium benzoate*, stearic acid, talc*, titanium dioxide

*contains one or more of these ingredients

Questions or comments?

Call toll free **1-888-423-0139**

Principal Display Panel

COMPARE TO COLACE® 2-IN-1 ACTIVE INGREDIENT†

Stool Softener

Plus Stimulant Laxative

DOCUSATE SODIUM 50 mg STOOL SOFTENER

SENNOSIDES 8.6 mg STIMULANT LAXATIVE

For Overnight Relief Of Occasional Constipation

- Effective
- Reliable
- Comfortable

TABLETS

GLUTEN FREE

†This product is not manufactured or distributed by Purdue Products L.P., distributor of Colace® 2-IN-1

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

DISTRIBUTED BY: TOPCO ASSOCIATES LLC

ELK GROVE VILLAGE, IL 60007

topcare@topco.com www.topcarebrand.com

Product Label

NDC 36800-084-01



Stool Softener + Stimulant Laxative

COMPARE TO COLACE® 2-IN-1 ACTIVE INGREDIENTS†

DOCUSATE SODIUM 50 mg
STOOL SOFTENER
SENNOSIDES 8.6 mg
STIMULANT LAXATIVE



For Overnight Relief Of Occasional Constipation:

- Effective
- Reliable
- Comfortable



actual size

100 TABLETS

DISTRIBUTED BY TOPCO ASSOCIATES LLC
ELK GROVE VILLAGE, IL 60007 © TOPCO
PLVA0618 QUESTIONS? 1-888-423-0139
topcare@topco.com www.topcarebrand.com
Visit here for more information:
<http://topbrnds.com/48zr8q>
PLD-H9T
FC004542



GLUTEN FREE

✓ QUALITY GUARANTEED



0 36800 16028 6

Lot No.:

Exp. Date:

Drug Facts

Active ingredients (in each tablet)	Purpose
Docusate sodium 50 mg.....	Stool softener
Sennosides 8.6 mg.....	Stimulant laxative

Uses

- for overnight relief from occasional constipation (irregularity)
- generally produces bowel movement in 6 to 12 hours

Warnings

- Do not use**
- laxative products for longer than 1 week unless directed by a doctor
 - if you are now taking mineral oil, unless directed by a doctor

Ask a doctor before use if you have

- stomach pain ■ nausea ■ vomiting
- noticed a sudden change in bowel habits that lasts over a period of 2 weeks

Stop use and ask a doctor if you have rectal bleeding or fail to have a bowel movement after use of a laxative. These may indicate a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take only by mouth. Doses may be taken as a single daily dose, preferably in the evening, or in divided doses.

Drug Facts (continued)

adults and children 12 years and older	take 2-4 tablets daily
children 6 to under 12 years of age	take 1-2 tablets daily
children 2 to under 6 years of age	take up to 1 tablet daily
children under 2	ask a doctor

Other information

- each tablet contains: calcium 20 mg
- each tablet contains: sodium 6 mg VERY LOW SODIUM
- store at 25°C (77°F); excursions permitted between 15-30°C (59-86°F)

Inactive ingredients

carnauba wax*, croscarmellose sodium, dibasic calcium phosphate dihydrate, FD&C blue #2 aluminum lake, FD&C red #40 aluminum lake, hypromellose*, magnesium stearate, maltodextrin*, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol*, silicon dioxide, sodium benzoate*, stearic acid, talc*, titanium dioxide
*contains one or more of these ingredients

Questions or comments?

Call toll free 1-888-423-0139

†This product is not manufactured or distributed by Purdue Products L.P., distributor of Colace® 2-IN-1.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

TOPCARE HEALTH Stool Softener + Stimulant Laxative

STOOL SOFTENER PLUS STIMULANT LAXATIVE

docusate sodium and sennosides tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:36800-084
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

DOCUSATE SODIUM (UNII: F05Q2T2JA0) (DOCUSATE - UNII:M7P27195AG)	DOCUSATE SODIUM	50 mg
SENNOSIDES (UNII: 3FYP5M0JJX) (SENNOSIDES - UNII:3FYP5M0JJX)	SENNOSIDES	8.6 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	

Product Characteristics

Color	red	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	TCL097;0806;AV;S44
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-084-01	1 in 1 BOX	01/31/2016	
1		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	01/31/2016	

Labeler - TOP CARE (Topco Associates LLC) (006935977)

