


**OXYBUTYNIN CHLORIDE - oxybutynin chloride tablet**  
**ANI Pharmaceuticals, Inc.**

-----  
**Oxybutynin Chloride Tablets, USP**  
**Rx only**

**DESCRIPTION**

Each scored oxybutynin chloride tablet, USP contains 5 mg of oxybutynin chloride. Chemically, oxybutynin chloride is d, l (racemic) 4-diethylamino-2-butynyl phenylcyclohexylglycolate hydrochloride. The molecular formula of oxybutynin chloride is  $C_{22}H_{31}NO_3 \cdot HCl$ . The structural formula appears below:



Oxybutynin chloride, USP is a white crystalline solid with a molecular weight of 393.95. It is readily soluble in water and acids, but relatively insoluble in alkalis.

Oxybutynin chloride Tablets, USP also contains anhydrous lactose, croscarmellose sodium, magnesium stearate, microcrystalline cellulose.

Oxybutynin chloride tablets, USP are for oral administration.

Therapeutic Category: Antispasmodic, anticholinergic.

Meets USP Dissolution Test 2.

**CLINICAL PHARMACOLOGY**

Oxybutynin chloride exerts a direct antispasmodic effect on smooth muscle and inhibits the muscarinic action of acetylcholine on smooth muscle. Oxybutynin chloride exhibits only one fifth of the anticholinergic activity of atropine on the rabbit detrusor muscle, but four to ten times the antispasmodic activity. No blocking effects occur at skeletal neuromuscular junctions or autonomic ganglia (antinicotinic effects).

Oxybutynin chloride relaxes bladder smooth muscle. In patients with conditions characterized by involuntary bladder contractions, cystometric studies have demonstrated that oxybutynin chloride increases bladder (vesical) capacity, diminishes the frequency of uninhibited contractions of the detrusor muscle, and delays the initial

desire to void. Oxybutynin chloride thus decreases urgency and the frequency of both incontinent episodes and voluntary urination.

Antimuscarinic activity resides predominately in the R-isomer. A metabolite, desethyloxybutynin, has pharmacological activity similar to that of oxybutynin in *in vitro* studies.

## Pharmacokinetics

### Absorption


Following oral administration of oxybutynin chloride tablets, oxybutynin is rapidly absorbed achieving  $C_{max}$  within an hour, following which plasma concentration decreases with an effective half-life of approximately 2 to 3 hours. The absolute bioavailability of oxybutynin is reported to be about 6% (range 1.6 to 10.9%) for the tablets. Wide interindividual variation in pharmacokinetic parameters is evident following oral administration of oxybutynin.

The mean pharmacokinetic parameters for R- and S-oxybutynin are summarized in **Table 1**. The plasma concentration-time profiles for R- and S-oxybutynin are similar in shape; **Figure 1** shows the profile for R-oxybutynin.

**Table 1** Mean (SD) R-and S-Oxybutynin Pharmacokinetic Parameters Following Three Doses of Oxybutynin Chloride 5 mg Administered every 8 Hours (n=23)

Parameters (units)	R-Oxybutynin	S-Oxybutynin
$C_{max}$ (ng/mL)	3.6 (2.2)	7.8 (4.1)
$T_{max}$ (h)	0.89 (0.34)	0.65 (0.32)
$AUC_t$ (ng•h/mL)	22.6 (11.3)	35.0 (17.3)
$AUC_{inf}$ (ng•h/mL)	24.3 (12.3)	37.3 (18.7)

**Figure 1.** Mean R-Oxybutynin Plasma Concentrations Following Three Doses of Oxybutynin Chloride 5 mg Administered Every 8 Hours for 1 Day in 23 Healthy Adult Volunteers.



Oxybutynin chloride steady-state pharmacokinetics were also studied in 11 pediatric patients with detrusor overactivity associated with a neurological condition (e.g., spina bifida). These pediatric patients were on oxybutynin chloride tablets with total daily dose ranging from 7.5 mg to 15 mg (0.22 to 0.53 mg/kg). Overall, most patients (86.9%) were taking a total daily oxybutynin chloride dose between 10 mg and 15 mg. Sparse sampling technique was used to obtain serum samples. When all available data are normalized to an equivalent of 5 mg twice daily oxybutynin chloride, the mean pharmacokinetic parameters derived for R- and S-oxybutynin and R- and S-desethyloxybutynin are summarized in **Table 2**. The plasma-time concentration profiles for R- and S-oxybutynin are similar in shape; **Figure 2** shows the profile for R-oxybutynin when all available data are normalized to an equivalent of 5 mg twice daily.

**Table 2** Mean  $\pm$  SD R- and S-Oxybutynin and R- and S-Desethyloxybutynin Pharmacokinetic Parameters in Children Aged 5 to 15 Following Administration of 7.5 mg to 15 mg Total Daily Dose of Oxybutynin Chloride Tablets (N=11)


All Available Data Normalized to an Equivalent of Oxybutynin Chloride Tablets 5 mg BID or TID at Steady State

	<b>R-Oxybutynin</b>	<b>S-Oxybutynin</b>	<b>R-Desethyloxybutynin</b>	<b>S-Desethyloxybutynin</b>
$C_{max}^*$ (ng/mL)	6.1 $\pm$ 3.2	10.1 $\pm$ 7.5	55.4 $\pm$ 17.9	28.2 $\pm$ 10.0
$T_{max}$ (hr)	1.0	1.0	2.0	2.0
$AUC^\dagger$ (ng•hr/mL)	19.8 $\pm$ 7.4	28.4 $\pm$ 12.7	238.8 $\pm$ 77.6	119.5 $\pm$ 50.7

\*Reflects  $C_{max}$  for pooled data

† $AUC_{0-\text{end of dosing interval}}$

**Figure 2.** Mean steady-state ( $\pm$ SD) R-oxybutynin plasma concentrations following administration of total daily Oxybutynin Chloride Tablet dose of 7.5 mg to 15 mg (0.22 mg/kg to 0.53 mg/kg) in children 5 to 15 years of age. – Plot represents all available data normalized to the equivalent of oxybutynin chloride 5 mg BID or TID at steady state



### **Food Effects**

Data in the literature suggests that oxybutynin solution co-administered with food resulted in a slight delay in absorption and an increase in its bioavailability by 25% (n=18).<sup>1</sup>

### **Distribution**

Oxybutynin is widely distributed in body tissues following systemic absorption. The volume of distribution is 193 L after intravenous administration of 5 mg oxybutynin chloride. Both enantiomers of oxybutynin are highly bound (>99%) to plasma proteins. Both enantiomers of desethyloxybutynin are also highly bound (>97%) to plasma proteins. The major binding protein is alpha-1 acid glycoprotein.

### **Metabolism**

Oxybutynin is metabolized primarily by the cytochrome P450 enzyme systems, particularly CYP3A4 found mostly in the liver and gut wall. Its metabolic products include phenylcyclohexylglycolic acid, which is pharmacologically inactive, and desethyloxybutynin, which is pharmacologically active.

### **Excretion**

Oxybutynin is extensively metabolized by the liver, with less than 0.1% of the administered dose excreted unchanged in the urine. Also, less than 0.1% of the

administered dose is excreted as the metabolite desethyloxybutynin

## **CLINICAL STUDIES**

Oxybutynin chloride was well tolerated in patients administered the drug in controlled studies of 30 days' duration and in uncontrolled studies in which some of the patients received the drug for 2 years.

## **INDICATIONS & USAGE**

Oxybutynin chloride tablets, USP are indicated for the relief of symptoms of bladder instability associated with voiding in patients with uninhibited neurogenic or reflex neurogenic bladder (i.e., urgency, frequency, urinary leakage, urge incontinence, dysuria).

## **CONTRAINDICATIONS**

Oxybutynin chloride tablets are contraindicated in patients with urinary retention, gastric retention and other severe decreased gastrointestinal motility conditions, uncontrolled narrow-angle glaucoma and in patients who are at risk for these conditions.

Oxybutynin chloride tablets are also contraindicated in patients who have demonstrated hypersensitivity to the drug substance or other components of the product.

## **WARNINGS**

Angioedema of the face, lips, tongue and/or larynx has been reported with oxybutynin. In some cases, angioedema occurred after the first dose. Angioedema associated with upper airway swelling may be life-threatening. If involvement of the tongue, hypopharynx, or larynx occurs, oxybutynin should be promptly discontinued and appropriate therapy and/or measures necessary to ensure a patent airway should be promptly provided.

## **PRECAUTIONS**

### **Central Nervous System Effects**

Oxybutynin is associated with anticholinergic central nervous system (CNS) effects (See **ADVERSE REACTIONS**). A variety of CNS anticholinergic effects have been reported, including hallucinations, agitation, confusion and somnolence. Patients should be monitored for signs of anticholinergic CNS effects, particularly in the first few months after beginning treatment or increasing the dose. If a patient experiences anticholinergic CNS effects, dose reduction or drug discontinuation should be considered.

Oxybutynin chloride should be used with caution in patients with preexisting dementia treated with cholinesterase inhibitors due to the risk of aggravation of symptoms.

Oxybutynin chloride should be used with caution in patients with Parkinson's disease due to the risk of aggravation of symptoms.

## **General**

Oxybutynin chloride should be used with caution in the frail elderly, in patients with hepatic or renal impairment, and in patients with myasthenia gravis.

Oxybutynin chloride may aggravate the symptoms of hyperthyroidism, coronary heart disease, congestive heart failure, cardiac arrhythmias, hiatal hernia, tachycardia, hypertension, myasthenia gravis, and prostatic hypertrophy.

## **Urinary Retention**

Oxybutynin chloride should be administered with caution to patients with clinically significant bladder outflow obstruction because of the risk of urinary retention (see **CONTRAINDICATIONS**).

## **Gastrointestinal Disorders**

Oxybutynin chloride should be used with caution in patients with autonomic neuropathy due to the risk of aggravation of symptoms of decreased gastrointestinal motility.

Oxybutynin chloride should be administered with caution to patients with gastrointestinal obstructive disorders because of the risk of gastric retention (see **CONTRAINDICATIONS**).

Administration of oxybutynin chloride to patients with ulcerative colitis may suppress intestinal motility to the point of producing a paralytic ileus and precipitate or aggravate toxic megacolon, a serious complication of the disease.

Oxybutynin chloride, like other anticholinergic drugs, may decrease gastrointestinal motility and should be used with caution in patients with conditions such as ulcerative colitis, and intestinal atony.

Oxybutynin chloride should be used with caution in patients who have gastroesophageal reflux and/or who are concurrently taking drugs (such as bisphosphonates) that can cause or exacerbate esophagitis.

## **Information for Patients**

Patients should be informed that oxybutynin may produce angioedema that could result in life-threatening airway obstruction. Patients should be advised to promptly discontinue oxybutynin therapy and seek immediate medical attention if they experience edema of the tongue, edema of the laryngopharynx, or difficulty breathing.

Patients should be informed that heat prostration (fever and heat stroke due to

decreased sweating) can occur when anticholinergics such as oxybutynin chloride are administered in the presence of high environmental temperature.

Because anticholinergic agents such as oxybutynin may produce drowsiness (somnolence), or blurred vision, patients should be advised to exercise caution.

Patients should be informed that alcohol may enhance the drowsiness caused by anticholinergic agents such as oxybutynin.

## **Drug Interactions**

The concomitant use of oxybutynin with other anticholinergic drugs or with other agents which produce dry mouth, constipation, somnolence (drowsiness), and/or other anticholinergic-like effects may increase the frequency and/or severity of such effects.

Anticholinergic agents may potentially alter the absorption of some concomitantly administered drugs due to anticholinergic effects on gastrointestinal motility. This may be of concern for drugs with a narrow therapeutic index. Anticholinergic agents may also antagonize the effects of prokinetic agents, such as metoclopramide.

Mean oxybutynin chloride plasma concentrations were approximately 3 to 4 fold higher when oxybutynin was administered with ketoconazole, a potent CYP3A4 inhibitor.

Other inhibitors of the cytochrome P450 3A4 enzyme system, such as antimycotic agents (e.g., itraconazole and miconazole) or macrolide antibiotics (e.g., erythromycin and clarithromycin), may alter oxybutynin mean pharmacokinetic parameters (i.e.,  $C_{max}$  and AUC). The clinical relevance of such potential interactions is not known. Caution should be used when such drugs are co-administered.

## **Carcinogenesis, Mutagenesis, Impairment of Fertility**

A 24-month study in rats at dosages of oxybutynin chloride of 20, 80, and 160 mg/kg/day showed no evidence of carcinogenicity. These doses are approximately 6, 25, and 50 times the maximum human exposure, based on surface area.

Oxybutynin chloride showed no increase of mutagenic activity when tested in *Schizosaccharomyces pompholiciformis*, *Saccharomyces cerevisiae* and *Salmonella typhimurium* test systems.

Reproduction studies using oxybutynin chloride in the hamster, rabbit, rat, and mouse have shown no definite evidence of impaired fertility.

## **Pregnancy**

### *Teratogenic Effects*

### **Category B.**

Reproduction studies using oxybutynin chloride in the hamster, rabbit, rat, and mouse have shown no definite evidence of impaired fertility or harm to the animal fetus. The safety of oxybutynin chloride administered to women who are or who may become pregnant has not been established. Therefore, oxybutynin chloride should not be given to pregnant women unless, in the judgment of the physician, the probable clinical benefits outweigh the possible hazards.

## **Nursing Mothers**

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when oxybutynin chloride is administered to a nursing woman.

## **Pediatric Use**

The safety and efficacy of oxybutynin chloride administration have been demonstrated for pediatric patients 5 years of age and older (see **DOSAGE AND ADMINISTRATION**).

The safety and efficacy of oxybutynin chloride tablets were studied in 30 children in a 24 week, open-label trial. Patients were aged 5 to 15 years, all had symptoms of detrusor overactivity in association with a neurological condition (e.g., spina bifida), all used clean intermittent catheterization, and all were current users of oxybutynin chloride. Study results demonstrated that the administration of oxybutynin chloride was associated with improvement in clinical and urodynamic parameters.

At total daily doses ranging from 5 mg to 15 mg, treatment with oxybutynin chloride tablets was associated with an increase from baseline in mean urine volume per catheterization from 122 mL to 145 mL, an increase from baseline in mean urine volume after morning awakening from 148 mL to 168 mL, and an increase from baseline in the mean percentage of catheterizations without a leaking episode from 43% to 61%. Urodynamic results in these patients were consistent with the clinical results. Treatment with oxybutynin chloride tablets was associated with an increase from baseline in maximum cystometric capacity from 230 mL to 279 mL, a decrease from baseline in mean detrusor pressure at maximum cystometric capacity from 36 cm H<sub>2</sub>O to 33 cm H<sub>2</sub>O, and a reduction in the percentage of patients demonstrating uninhibited detrusor contractions (of at least 15 cm H<sub>2</sub>O) from 39% to 20%.

As there is insufficient clinical data for pediatric populations under age 5, oxybutynin chloride is not recommended for this age group.

## **Geriatric Use**

Clinical studies of oxybutynin chloride did not include sufficient numbers of subjects age 65 and over to determine whether they respond differently from younger patients. Other reported clinical experience has not identified differences in responses between healthy elderly and younger patients; however, a lower initial starting dose of 2.5 mg given 2 or 3 times a day has been recommended for the frail elderly due to a prolongation of the elimination half-life from 2 to 3 hours to 5 hours.<sup>2,3,4</sup> In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

## **ADVERSE REACTIONS**

The safety and efficacy of oxybutynin chloride was evaluated in a total of 199 patients in three clinical trials. These participants were treated with oxybutynin chloride 5 to 20 mg/day for up to 6 weeks. Table 3 shows the incidence of adverse events judged by investigators to be at least possibly related to treatment and reported by at least 5% of



patients.

**Table 3** Incidence (%) of Adverse Events Reported by  $\geq 5\%$  of Patients Using Oxybutynin Chloride (5 to 20 mg/day)

<b>Body System</b>	<b>Adverse Event</b>	<b>Oxybutynin Chloride (5to 20 mg/day) (n=199)</b>
Infections and Infestations	Urinary tract infection	6.5%
Psychiatric Disorders	Insomnia	5.5%
	Nervousness	6.5%
	Dizziness	16.6%
Nervous System Disorders	Somnolence	14.0%
	Headache	7.5%
	Blurred vision	9.6%
Eye Disorders	Dry mouth	71.4%
	Constipation	15.1%
	Nausea	11.6%
	Dyspepsia	6.0%
Renal and Urinary Disorders	Urinary Hesitation	8.5%
	Urinary Retention	6.0%

The most common adverse events reported by patients receiving oxybutynin chloride 5 to 20 mg/day were the expected side effects of anticholinergic agents. The incidence of dry mouth was dose-related.

In addition, the following adverse events were reported by 1 to <5% of patients using oxybutynin chloride (5 to 20 mg/day) in all studies.

*Infections and Infestations:* nasopharyngitis, upper respiratory tract infection, bronchitis, cystitis, fungal infection; *Metabolism and Nutrition Disorders:* fluid retention; *Psychiatric Disorders:* confusional state; *Nervous System Disorders:* dysgeusia, sinus headache; *Eye Disorders:* kerato conjunctivitis sicca, eye irritation; *Cardiac Disorders:* palpitations, sinus

arrhythmia; *Vascular Disorders*: flushing; *Respiratory, thoracic and Mediastinal Disorders*: nasal dryness, cough, pharyngolaryngeal pain, dry throat, sinus congestion, hoarseness, asthma, nasal congestion; *Gastrointestinal Disorders*: diarrhea, abdominal pain, loose stools, flatulence, vomiting, abdominal pain upper, dysphagia, apyhalism, eructation, tongue coated; *Skin and Subcutaneous Tissue Disorders*: dry skin, pruritis; *Musculoskeletal and Connective Tissue Disorders*: back pain, arthralgia, pain in extremity, flank pain; *Renal and Urinary Disorders*: dysuria, pollakiuria; *General Disorders and Administration Site Conditions*: fatigue, edema peripheral, asthenia, pain, thirst, edema; *Investigations*: blood pressure increased, blood glucose increased, blood pressure decreased; *Injury, Poisoning, and Procedural Complications*: fall.

## **Postmarketing Surveillance**

Because postmarketing adverse events are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure. The following additional adverse events have been reported from worldwide postmarketing experience with oral oxybutynin chloride:

*Psychiatric Disorders*: psychotic disorder, agitation, hallucinations, memory impairment; *Nervous System Disorders*: convulsions; *Eye Disorders*: cycloplegia, mydriasis, glaucoma; *Cardiac Disorders*: tachycardia, QT interval prolongation, chest discomfort; *Gastrointestinal Disorders*: decreased gastrointestinal motility, frequent bowel movements; *Skin and Subcutaneous Tissue Disorders*: rash, decreased sweating; *Renal and Urinary Disorders*: impotence; *Reproductive System and Breast Disorders*: Suppression of lactation; *General Disorders and Administration Site Conditions*: hypersensitivity reactions, including angioedema with airway obstruction, urticaria, and face edema; rare anaphylactic reactions requiring hospitalization for emergency treatment; *Metabolism and Nutrition Disorders*: anorexia; *Respiratory, Thoracic and Mediastinal Disorders*: dysphonia.

## **OVERDOSAGE**

Treatment should be symptomatic and supportive. Activated charcoal as well as a cathartic may be administered.

Overdosage with oxybutynin chloride has been associated with anticholinergic effects including central nervous system excitation (e.g., restlessness, tremor, irritability, convulsions, delirium, hallucinations), flushing, fever, dehydration, cardiac arrhythmia, vomiting, and urinary retention. Other symptoms may include hypotension or hypertension, respiratory failure, paralysis, and coma.

Ingestion of 100 mg oxybutynin chloride in association with alcohol has been reported in a 13-year-old boy who experienced memory loss, and a 34-year-old woman who developed stupor, followed by disorientation and agitation on awakening, dilated pupils, dry skin, cardiac arrhythmia, and retention of urine. Both patients fully recovered with symptomatic treatment.

## **DOSAGE & ADMINISTRATION**

### **Adults**

The usual dose is one 5-mg tablet two to three times a day. The maximum recommended dose is one 5-mg tablet four times a day. A lower starting dose of 2.5 mg two or three times a day is recommended for the frail elderly.

### **Pediatric patients over 5 years of age**

The usual dose is one 5-mg tablet two times a day. The maximum recommended dose is one 5-mg tablet three times a day.

### **HOW SUPPLIED**

Oxybutynin Chloride Tablets, USP 5 mg are white, round, biconvex tablets debossed with “n005” on one side and bisect on the other side. They are supplied as follows:

NDC 70954- <b>005</b> -10	Bottles of 100 Tablets
NDC 70954- <b>005</b> -20	Bottles of 500 Tablets
NDC 70954- <b>005</b> -30	Bottles of 1000 Tablets
NDC 70954- <b>005</b> -50	Unit dose blister packages of 100 (10 cards of 10 tablets each)

**Pharmacist: Dispense in tight, light-resistant container as defined in the USP.**

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

### **REFERENCES**

1. Yong C et al. Effect of Food on the Pharmacokinetics of Oxybutynin in normal subjects. *Pharm Res.* 1991; 8 (Suppl.): S-320.
2. Hughes KM et al. Measurement of oxybutynin and its N-desethyl metabolite in plasma, and its application to pharmacokinetic studies in young, elderly and frail elderly volunteers. *Xenobiotica.* 1992; 22 (7): 859-869.
3. Ouslander J et al. Pharmacokinetics and Clinical Effects of Oxybutynin in Geriatric Patients. *J. Urol.* 1988; 140: 47-50.
4. Yarker Y et al. Oxybutynin: A review of its Pharmacodynamic and Pharmacokinetic Properties, and its Therapeutic Use in Detrusor Instability. *Drugs & Aging.* 1995; 6(3): 243-262.

Manufactured by:

**Novitium Pharma LLC**

70 Lake Drive, East Windsor

New Jersey 08520

**Issued:** 11/2024

LB4461-01

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL**

**PRINCIPAL DISPLAY PANEL - 5 MG CONTAINER LABEL - 100'S COUNT**

NDC 70954-005-10

**100 Tablets**

**Oxybutynin Chloride Tablets, USP 5 mg**

**Novitium Pharma LLC**

**Rx only**

NDC 70954-005-10 100 Tablets

**Oxybutynin Chloride Tablets, USP**

**5 mg**

**novitium pharma** | Rx only

**Each tablet contains:**  
Oxybutynin Chloride, USP ..... 5 mg

**Dosage and administration:** Adults – One tablet two to four times a day. Pediatric patients – (see Precautions). One tablet two to three times a day. See package insert for full prescribing information.

**Store at 20° to 25°C (68° to 77°F)** [see USP Controlled Room Temperature].

**Dispense in tight, light-resistant container as defined in the USP.**

**WARNING:** Keep this and all medications out of the reach of children.

Manufactured by:  
**Novitium Pharma LLC**  
70 Lake Drive, East Windsor  
New Jersey 08520

LB4462-00  
**Issued:** 03/2022

3 70954-005-10 0

**PRINCIPAL DISPLAY PANEL - 5 MG CONTAINER LABEL - 500'S COUNT**

NDC 70954-005-20

**500 Tablets**

**Oxybutynin Chloride Tablets, USP 5 mg**

**Novitium Pharma LLC**

**Rx only**




# Oxybutynin Chloride Tablets, USP 5 mg

Novitium Pharma LLC

Rx only

<p>Oxybutynin Chloride Tablet, USP</p> <p><b>5 mg</b></p> <p>Exp. XXX 0000 Lot# X000000X</p>  <p>NDC 70954-005-40 LB4276-00 Iss: 06/2021 Mfg. by: Novitium Pharma LLC, East Windsor, NJ</p>	<p>Oxybutynin Chloride Tablet, USP</p> <p><b>5 mg</b></p> <p>Exp. XXX 0000 Lot# X000000X</p>  <p>NDC 70954-005-40 LB4276-00 Iss: 06/2021 Mfg. by: Novitium Pharma LLC, East Windsor, NJ</p>
<p>Oxybutynin Chloride Tablet, USP</p> <p><b>5 mg</b></p> <p>Exp. XXX 0000 Lot# X000000X</p>  <p>NDC 70954-005-40 LB4276-00 Iss: 06/2021 Mfg. by: Novitium Pharma LLC, East Windsor, NJ</p>	<p>Oxybutynin Chloride Tablet, USP</p> <p><b>5 mg</b></p> <p>Exp. XXX 0000 Lot# X000000X</p>  <p>NDC 70954-005-40 LB4276-00 Iss: 06/2021 Mfg. by: Novitium Pharma LLC, East Windsor, NJ</p>
<p>Oxybutynin Chloride Tablet, USP</p> <p><b>5 mg</b></p> <p>Exp. XXX 0000 Lot# X000000X</p>  <p>NDC 70954-005-40 LB4276-00 Iss: 06/2021 Mfg. by: Novitium Pharma LLC, East Windsor, NJ</p>	<p>Oxybutynin Chloride Tablet, USP</p> <p><b>5 mg</b></p> <p>Exp. XXX 0000 Lot# X000000X</p>  <p>NDC 70954-005-40 LB4276-00 Iss: 06/2021 Mfg. by: Novitium Pharma LLC, East Windsor, NJ</p>
<p>Oxybutynin Chloride Tablet, USP</p> <p><b>5 mg</b></p> <p>Exp. XXX 0000 Lot# X000000X</p>  <p>NDC 70954-005-40 LB4276-00 Iss: 06/2021 Mfg. by: Novitium Pharma LLC, East Windsor, NJ</p>	<p>Oxybutynin Chloride Tablet, USP</p> <p><b>5 mg</b></p> <p>Exp. XXX 0000 Lot# X000000X</p>  <p>NDC 70954-005-40 LB4276-00 Iss: 06/2021 Mfg. by: Novitium Pharma LLC, East Windsor, NJ</p>



## OXYBUTYNIN CHLORIDE

oxybutynin chloride tablet

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:70954-005
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OXYBUTYNIN CHLORIDE (UNII: L9F3D9RENQ) (OXYBUTYNIN - UNII:K9P6MC7092)	OXYBUTYNIN CHLORIDE	5 mg

### Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)

### Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	2 pieces
<b>Shape</b>	ROUND	<b>Size</b>	8mm
<b>Flavor</b>		<b>Imprint Code</b>	n005
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70954-005-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	11/15/2017	
2	NDC:70954-005-20	500 in 1 BOTTLE; Type 0: Not a Combination Product	11/15/2017	
3	NDC:70954-005-30	1000 in 1 BOTTLE; Type 0: Not a Combination Product	11/15/2017	
4	NDC:70954-005-50	100 in 1 CARTON	08/20/2021	
4	NDC:70954-005-40	1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA209823	11/15/2017	

**Labeler** - ANI Pharmaceuticals, Inc. (145588013)

### Establishment

Name	Address	ID/FEI	Business Operations
Novitium Pharma LLC		080301870	ANALYSIS(70954-005) , LABEL(70954-005) , MANUFACTURE(70954-005) , PACK(70954-005)

Revised: 11/2024

ANI Pharmaceuticals, Inc.