

UV PROTECTION VEIL MINERAL SUNSCREEN, TINTED- titanium dioxide, zinc oxide cream
ATHENA PLASTIC SURGERY

Active Ingredients

Titanium Dioxide 8% Zinc Oxide 3.8%

Purpose

Sunscreen

Uses

Helps Prevent Sunburn. If used as directed with other sun protection measures (see "Directions"), decreases the risk of skin cancer and early skin aging caused by the sun.

Warnings

For external use only. Do not use on broken or damaged skin. Stop use and ask a doctor if rash occurs. When using this product, keep out of eyes. Rinse with water to remove. Keep out of reach of children. If swallowed, get medical help or contact a poison control center right away.

Warning

Keep out of reach of children

Directions

Apply liberally 15 minutes before sun exposure. Use a water-resistant product if swimming or sweating. Reapply at least every 2 hours. Children under 6 months: Ask a doctor. Sun Protection Measures: Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad-Spectrum SPF value of 15 or higher and other sun protection measures including: Limit time in the sun, especially from 10 a.m. - 2 p.m. Wear long-sleeved shirts, pants, hats and sunglasses.

Inactive Ingredients

Alumina, Cyclohexasiloxane, Cyclopentasiloxane, Dimethicone, Dimethicone Crosspolymer, Dimethicone/Vinyl Dimethicone Crosspolymer, Dimethiconol, Iron Oxide (CI 77491), Lauryl PEG/PPG-18/18 Methicone, Hydrogen Dimethicone, PEG-10 Dimethicone, Tetrahexyldecyl Ascorbate, Tocopheryl Acetate

Other Information

- Protect thei product from excessive heat and direct sun
- May stain some fabrics
- Cleanse thoroughly at nightttime
- PA++++

DRUG FACTS

ACTIVE INGREDIENTS:

Titanium Dioxide 8%

Zinc Oxide 3.8%

PURPOSE:

Sunscreen

Sunscreen

USES:

Helps prevent sunburn. If used as directed with other sun protection measures (see directions), decreases the risk of skin cancer and early skin aging caused by the sun.

WARNINGS:

For external use only. Do not use on damaged or broken skin. Stop use and ask a doctor if rash occurs. When using this product, keep out of eyes. Rinse with water to remove. Keep out of the reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

DIRECTIONS:

- Apply liberally 15 minutes before sun exposure.
- Reapply:
 - After 80 minutes of swimming or sweating
 - Immediately after towel drying
 - At least every 2 hours
- Children under 6 months: Ask a doctor
- Sun Protection Measures: Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:
 - Limit time in the sun, especially from 10 am-2 pm
 - Wear long-sleeved shirts, pants, hats, and sunglasses

INACTIVE INGREDIENTS:

Alumina, Cyclohexasiloxane, Cyclopentasiloxane, Dimethicone, Dimethicone Crosspolymer, Dimethicone/Vinyl Dimethicone Crosspolymer, Dimethiconol, Iron Oxide (CI 44791), Lauryl PEG/PPG-18/18 Hydrogen Dimethicone, Hydrogen Dimethicone, PEG-10 Dimethicone, Tetrahexyldecyl Ascorbate, Tocopheryl Acetate

OTHER INFORMATION:

- Protect this product from excessive heat and direct sun
- May stain some fabrics
- Cleanse thoroughly at nighttime
- PA+++

Athena Plastic Surgery

& Med Spa



UV Protection Veil

Mineral Sunscreen

TINTED

SPF 40 Broad Spectrum

Water Resistant (80 mins)

UVA|UVB Protection

Athena Plastic Surgery | 772-266-5003 |

www.athenaplasticsurgery.com

903 SE Monterey Commons Blvd, Stuart, FL 34996



3 85926 20350 7

1.8 oz | 53 g

UV PROTECTION VEIL MINERAL SUNSCREEN, TINTED

titanium dioxide, zinc oxide cream

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:85926-203(NDC:58892-203)
Route of Administration	TOPICAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)		TITANIUM DIOXIDE	80 mg in 1 g
ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)		ZINC OXIDE	38 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
DIMETHICONOL (2000 CST) (UNII: T74O12AN6Y)	
CYCLOHEXASILOXANE (UNII: XHK3U310BA)	
HYDROGEN DIMETHICONE (20 CST) (UNII: 12Z59IF64N)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
ALPHA-TOCOPHEROL ACETATE (UNII: 9E8X80D2L0)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
DIMETHICONE CROSSPOLYMER (UNII: UF7620L1W6)	
TETRAHEXYLDECYL ASCORBATE (UNII: 9LBV3F07AZ)	
CI 77491 (UNII: 1K09F3G675)	
CYCLOPENTASILOXANE (UNII: 0THT5PCI0R)	
ALUMINA (UNII: LMI26O6933)	
CI 77492 (UNII: EX438O2MRT)	
CI 77499 (UNII: XM0M87F357)	
LAURYL PEG/PPG-18/18 METHICONE (UNII: ZJ5S27D9NX)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:85926-203-50	53 g in 1 BOTTLE, PUMP; Type 0: Not a Combination Product	10/24/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M020	10/24/2019	

Labeler - ATHENA PLASTIC SURGERY (108606334)

Establishment

Name	Address	ID/FEI	Business Operations
Custom Analytics LLC		144949372	analysis(85926-203)

Establishment

Name	Address	ID/FEI	Business Operations
Fragrance Manufacturing Inc		793406000	manufacture(85926-203)

Establishment

Name	Address	ID/FEI	Business Operations
Fallien Cosmeceuticals Ltd.		958388357	relabel(85926-203)

