

**SALICYLIC ACID- wart remover liquid**  
**Chain Drug Consortium, LLC**

-----  
**Premier Value Liquid Wart Remover**

***Active ingredient***

Salicylic acid

***Purpose***

Wart Remover

***Use***

- for the removal of common and plantar warts. The common wart is easily recognized by the rough 'cauliflower-like' appearance of the surface. The plantar wart exists only on the bottom of the foot and interrupts footprint pattern.

***Warnings***

**For external use only.**

**Flammable**

- keep away from fire or flame
- **cap bottle tightly** and store at room temperature away heat

**Do not use**

- if you are a diabetic or have poor blood circulation, except under the advice and supervision of a doctor or podiatristirritated, infected or reddened skin
- genital warts and warts on the face
- moles, birthmarks and warts with hair growing from them
- mucous membranes

**When using this product**

- avoid contact with eyes. If product gets into the eyes, flush with water for 15 minutes
- avoid inhaling vapors

**Stop use and ask doctor if**

discomfort persists

**Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center right away.

### **Directions**

- wash affected area, may soak wart in warm water for 5 minutes
- dry area thoroughly
- apply small amount of wart remover with applicator to sufficiently cover each wart
- let dry and repeat this procedure once or twice daily until wart is removed for up to 12 weeks

### **Other information**

store between 20°C to 30°C (68°F to 86°F)

### **Inactive ingredients**

camphor, castor oil, ethanol, ethyl ether, nitrocellulose

### **Questions?**

Call 1-866-964-0939

### **Principal Display Panel**

Premier Value

Maximum Strength

Liquid Wart Remover  
with Salicylic acid

60% More than Compound W

Removes common & plantar warts safely

1/2 FL OZ (15mL)



and plantar warts safely and easily

1/2 FL OZ (15mL)



60% More than Compound W®

MAXIMUM STRENGTH  
LIQUID WART REMOVER  
WITH SALICYLIC ACID

BWRLHRPV-2



Removes common warts and plantar warts safely and easily

MAXIMUM STRENGTH  
LIQUID WART REMOVER  
WITH SALICYLIC ACID

1/2 FL OZ (15mL)



NO VARNISH AREA

E.P.O.  
8-40986 01437-6



If for any reason you are not satisfied with this product, please return it to the store where purchased for a full refund.

DISTRIBUTED BY CHAIN DRUG CONSORTIUM, 3301 NW BOCA RATON BLVD., SUITE 101, BOCA RATON, FL 33431  
Jackson, WY 83001 owner of the registered trademark Compound W®.  
This product is not manufactured or distributed by Medtech Products, Inc.

Questions? Call 1-866-964-0939

**Inactive ingredient** flexible collation containing alcohol and ether

**Other information** store between 20° to 30°C (68° to 85°F)

**Directions** ■ wash or soak the affected area and dry thoroughly  
■ apply wart remover liquid, with enclosed applicator, until wart is covered; then let dry  
■ reapply subsequent applications over existing application  
■ repeat this procedure once or twice daily, until wart is completely removed.  
for up to 12 weeks

**Keep out of reach of children.**  
If swallowed, get medical help or contact a Poison Control Center right away.

**Stop use and ask a doctor if discomfort persists:**  
■ avoid inhaling vapors  
■ avoid contact with eyes; if product gets into the eye, flush with water for 15 minutes

**When using this product**  
■ moles, birthmarks and warts with hair growing from them ■ mucous membranes  
■ irritated, infected or reddened skin ■ genital warts and warts on the face

**Do not use on**  
supervision of a doctor or podiatrist

**Do not use**  
■ if you are diabetic or have poor blood circulation, except under the advice and


■ cap bottle tightly and store at room temperature away from heat  
■ keep away from fire or flame  
Flammable

**Warnings**  
For external use only.

**Use** ■ for the removal of common and plantar warts. The common wart has a rough, cauliflower-like appearance. The plantar wart exists only on the bottom of the foot and interrupts the footprint pattern.

**Drug Facts**  
**Active ingredient** Salicylic acid 17% w/w.  
**Purpose** Wart removal

**DO NOT USE IF NECKBAND IS BROKEN**



**DO NOT USE IF NECKBAND IS BROKEN**

**USES:** For the removal of common and plantar warts. The common wart has a rough 'cauliflower-like' appearance. The plantar wart exists only on the bottom of the foot and interrupts the footprint pattern.

**DIRECTIONS:** Wash or soak the affected area and dry thoroughly. Apply wart remover liquid with enclosed applicator until wart is covered and let dry. Reapply subsequent applications over existing application. Repeat this procedure once or twice daily, until wart is completely removed, for up to 12 weeks.



**MAXIMUM STRENGTH  
LIQUID  
WART  
REMOVER  
WITH SALICYLIC ACID**

Removes common warts and plantar warts safely and easily

1/2 FL OZ (15mL)

**ACTIVE INGREDIENT:** Salicylic acid (17% w/w)

**INACTIVE INGREDIENT:** Flexible colloidion containing alcohol and ether

**WARNINGS:** For external use only. **Flammable:** Keep away from fire or flame. **Cap bottle tightly** and store at room temperature away from heat. Do not use this product if you are diabetic or have poor blood circulation, except under the advice and supervision of a doctor or podiatrist. Do not use on irritated, infected or reddened skin or on moles, birthmarks, warts with hair growing from them, genital warts, warts on the face and mucous membranes. Avoid contact with eyes. If product gets into the eye, flush with water for 15 minutes. Avoid inhaling vapors. Stop use and ask a doctor if discomfort persists. **Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center right away. **Questions?** 1-866-964-0939

**DISTRIBUTED BY: CHAIN DRUG CONSORTIUM, LLC**

**MADE IN USA**

WRLABELPV-1

## SALICYLIC ACID

wart remover liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:68016-611
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SALICYLIC ACID (UNII: 0414PZ4LPZ) (SALICYLIC ACID - UNII:0414PZ4LPZ)	SALICYLIC ACID	0.17 mg in 15 mL

### Inactive Ingredients

Ingredient Name	Strength
PYROXYLIN (UNII: KYR8BR2X6O)	
CAMPHOR (NATURAL) (UNII: N20HL7Q941)	
CASTOR OIL (UNII: D5340Y219G)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68016-611-00	15 mL in 1 BOTTLE, WITH APPLICATOR; Type 0: Not a Combination Product	05/23/2003	

### Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M028	05/23/2003	

**Labeler** - Chain Drug Consortium, LLC (101668460)

Revised: 1/2025

Chain Drug Consortium, LLC