

**ALEVE EASY OPEN ARTHRITIS CAP- naproxen sodium capsule, liquid filled
Bayer HealthCare LLC.**

ALEVE ®

LIQUID GELS

Drug Facts

Active ingredient (in each capsule)

Naproxen sodium 220 mg (naproxen 200 mg) (NSAID) ¹

¹ nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - menstrual cramps
 - headache
 - toothache
 - the common cold
- temporarily reduces fever

Allergy alert

Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days

- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective doses should be used**
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

adults and children 12 years and older:	<ul style="list-style-type: none">• take 1 capsule every 8 to 12 hours while symptoms last• for the first dose you may take 2 capsules within the first hour• do not exceed 2 capsules in any 8- to 12-hour period• do not exceed 3 capsules in a 24-hour period
children under 12 years:	<ul style="list-style-type: none">• ask a doctor

Inactive ingredients

FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Other information

- **each capsule contains:** sodium 20 mg
- store at 20-25°C (68-77°F) avoid high humidity and excessive heat above 40°C (104°F)
- read all directions and warnings before use. Keep carton.

Questions or comments?

1-800-395-0689 (Mon – Fri 9AM – 5PM EST)

Distributed by:

Whippany

Bayer Healthcare LLC

Whippany, NJ 07981

**EZ OPEN ARTHRITIS CAP
ALL DAY STRONG®**

ALEVE®

LIQUID GELS

naproxen sodium capsules, 220 mg (NSAID)

pain reliever/fever reducer

STRENGTH TO LAST 12 HOURS

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

80

CAPSULES

(LIQUID FILLED CAPSULES)



ALEVE EASY OPEN ARTHRITIS CAP

naproxen sodium capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0280-6091
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
LACTIC ACID (UNII: 33X04XA5AT)	
MANNITOL (UNII: 3OWL53L36A)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POVIDONE K30 (UNII: U725QWY32X)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITAN (UNII: 6O92ICV9RU)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color	blue (Blue with white text)	Score	no score
Shape	CAPSULE	Size	21mm
Flavor		Imprint Code	ALEVE
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0280-6091-80	1 in 1 CARTON	07/28/2007	
1		80 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA021920	02/28/2007	

Labeler - Bayer HealthCare LLC. (112117283)

Revised: 12/2025

Bayer HealthCare LLC.