

PAIN RELIEF REGULAR STRENGTH- acetaminophen tablet
Rite Aid Corporation

Rite Aid 44-104

Active ingredient (in each tablet)

Acetaminophen 325 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold
 - headache
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if the user has

liver disease.

Ask a doctor or pharmacist before use if the user is

taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days in adults
- redness or swelling is present
- pain gets worse or lasts more than 5 days in children under 12 years
- new symptoms occur
- fever gets worse or lasts more than 3 days

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**

adults and children 12 years and over	<ul style="list-style-type: none">▪ take 2 tablets every 4 to 6 hours while symptoms last▪ do not take more than 10 tablets in 24 hours▪ do not take for more than 10 days unless directed by a doctor
children 6-11 years	<ul style="list-style-type: none">▪ take 1 tablet every 4 to 6 hours while symptoms last▪ do not take more than 5 tablets in 24 hours▪ do not take for more than 5 days unless directed by a doctor
children under 6 years	ask a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, povidone, sodium starch glycolate, stearic acid

Questions or comments?

1-800-426-9391

Principal Display Panel

NDC 11822-0104-1

Compare to the active ingredient of **Regular Strength Tylenol®***

**REGULAR STRENGTH
PAIN RELIEF
ACETAMINOPHEN**

ACETAMINOPHEN 325 mg

PAIN RELIEVER/FEVER REDUCER

RITE

AID®

PHARMACY

ACTUAL SIZE

100 TABLETS

*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Regular Strength Tylenol®.

50844 REV0417D10412

DISTRIBUTED BY:

RITE AID, 30 HUNTER LANE,
CAMP HILL, PA 17011

www.riteaid.com

SATISFACTION

GUARANTEE

If you're not satisfied, we'll
happily refund your money.

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

NDC 11822-0104-1 Compare to the active ingredient of **Regular Strength Tylenol®**

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING



0 11822 31764 1

REGULAR STRENGTH PAIN RELIEF ACETAMINOPHEN

ACETAMINOPHEN 325 mg

PAIN RELIEVER/FEVER REDUCER



ACTUAL SIZE

100

TABLETS



100

TABLETS

REGULAR STRENGTH PAIN RELIEF ACETAMINOPHEN

ACETAMINOPHEN 325 mg

PAIN RELIEVER/FEVER REDUCER

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Drug Facts	Purpose Pain reliever/fever reducer
Active ingredient (in each tablet) Acetaminophen 325 mg	
Uses	<ul style="list-style-type: none"> ■ temporarily relieves minor aches and pains due to: <ul style="list-style-type: none"> ■ the common cold ■ headache ■ backache ■ minor pain of arthritis ■ toothache ■ muscular aches ■ premenstrual and menstrual cramps ■ temporarily reduces fever
Warnings	<p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if</p> <ul style="list-style-type: none"> ■ adult takes more than 4,000 mg of acetaminophen in 24 hours ■ child takes more than 5 doses in 24 hours ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product <p>Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:</p> <ul style="list-style-type: none"> ■ skin redness ■ hives ■ rash <p>If a skin reaction occurs, stop use and seek medical help right away.</p>
Do not use	<ul style="list-style-type: none"> ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product

B-1702-104-12-R2
REV0417D10412

Drug Facts (continued)	<p>Other information</p> <ul style="list-style-type: none"> ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ see end flap for expiration date and lot number <p>Inactive ingredients corn starch, povidone, sodium starch glycolate, stearic acid</p> <p>Questions or comments? 1-800-426-9391</p> <p><small>* This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Regular Strength Tylenol®.</small></p>
-------------------------------	--

DISTRIBUTED BY:
RITE AID, 30 HUNTER LANE,
CAMP HILL, PA 17011
www.riteaid.com

SATISFACTION GUARANTEE

If you're not satisfied, we'll happily refund your money.

Drug Facts (continued)	<p>Ask a doctor before use if the user has liver disease.</p> <p>Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin.</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ pain gets worse or lasts more than 10 days in adults ■ redness or swelling is present ■ pain gets worse or lasts more than 5 days in children under 12 years ■ new symptoms occur ■ fever gets worse or lasts more than 3 days <p>These could be signs of a serious condition.</p> <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>Directions</p> <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 30%;">adults and children 12 years and over</td> <td> <ul style="list-style-type: none"> ■ do not take more than directed ■ take 2 tablets every 4 to 6 hours while symptoms last ■ do not take more than 10 tablets in 24 hours ■ do not take for more than 10 days unless directed by a doctor </td> </tr> <tr> <td>children 6-11 years</td> <td> <ul style="list-style-type: none"> ■ take 1 tablet every 4 to 6 hours while symptoms last ■ do not take more than 5 tablets in 24 hours ■ do not take for more than 5 days unless directed by a doctor </td> </tr> <tr> <td>children under 6 years</td> <td>ask a doctor</td> </tr> </table>	adults and children 12 years and over	<ul style="list-style-type: none"> ■ do not take more than directed ■ take 2 tablets every 4 to 6 hours while symptoms last ■ do not take more than 10 tablets in 24 hours ■ do not take for more than 10 days unless directed by a doctor 	children 6-11 years	<ul style="list-style-type: none"> ■ take 1 tablet every 4 to 6 hours while symptoms last ■ do not take more than 5 tablets in 24 hours ■ do not take for more than 5 days unless directed by a doctor 	children under 6 years	ask a doctor
adults and children 12 years and over	<ul style="list-style-type: none"> ■ do not take more than directed ■ take 2 tablets every 4 to 6 hours while symptoms last ■ do not take more than 10 tablets in 24 hours ■ do not take for more than 10 days unless directed by a doctor 						
children 6-11 years	<ul style="list-style-type: none"> ■ take 1 tablet every 4 to 6 hours while symptoms last ■ do not take more than 5 tablets in 24 hours ■ do not take for more than 5 days unless directed by a doctor 						
children under 6 years	ask a doctor						

Rite Aid 44-104

PAIN RELIEF REGULAR STRENGTH

acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-0104
Route of Administration	ORAL		

Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	325 mg	
Inactive Ingredients				
Ingredient Name		Strength		
STARCH, CORN (UNII: O8232NY3SJ)				
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)				
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)				
STEARIC ACID (UNII: 4ELV7Z65AP)				
Product Characteristics				
Color	white	Score	2 pieces	
Shape	ROUND	Size	10mm	
Flavor		Imprint Code	44;104	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-0104-1	1 in 1 CARTON	07/13/1990	
1		100 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:11822-0104-7	1 in 1 CARTON	07/13/1990	09/06/2021
2		125 in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M013	07/13/1990		

Labeler - Rite Aid Corporation (014578892)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(11822-0104)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(11822-0104)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(11822-0104)

Revised: 8/2025

Rite Aid Corporation