BASIC CARE TUSSIN CF SEVERE- acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride solution Amazon.com Services LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Amazon Tussin CF Severe Drug Facts

Active ingredients (in each 20 mL)

Acetaminophen, USP 650 mg
Dextromethorphan HBr, USP 20 mg
Guaifenesin, USP 400 mg
Phenylephrine HCl, USP 10 mg

Purposes

Pain reliever/fever reducer
Cough suppressant
Expectorant
Nasal decongestant

Uses

- temporarily relieves these symptoms occurring with a cold or flu:
- cough due to minor throat and bronchial irritation
- nasal congestion
- sinus congestion and pressure
- minor aches and pains
- sore throat
- headache
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen

• 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

When using this product

do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days

- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than 5 doses in any 24-hour period
- do not exceed recommended dosage. Taking more than the recommended dose (overdose) may cause serious liver damage.
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	20 mL every 4 hours
children under 12 years	do not use

Other information

- each 20 mL contains: sodium 15 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

anhydrous citric acid, benzyl alcohol, edetate disodium, FD&C red #40, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose, triacetin, xanthan gum

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

For Ages 12 & Over

Compare to Robitussin® Maximum Strength Severe Multi-Symptom Cough Cold + Flu active ingredients

Maximum Strength

Tussin CF Severe

Multi-Symptom

Cough Cold + Flu

Pain Reliever/Fever Reducer (Acetaminophen)

Cough Suppressant (Dextromethorphan HBr)

Expectorant (Guaifenesin)

Nasal Decongestant (Phenylephrine HCl)

Relieves:

Cough, Sore Throat

Body Aches, Fever

Nasal Congestion

Chest Congestion

Non-Drowsy

For Adults

Dosage Cup Included

8 FL OZ (237 mL)



BASIC CARE TUSSIN CF SEVERE

acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride solution

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72288-723
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	650 mg in 20 mL	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL	
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL	

Inactive Ingredients		
Ingredient Name	Strength	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)		
BENZYL ALCOHOL (UNII: LKG8494WBH)		
EDETATE DISODIUM (UNII: 7FLD91C86K)		
FD&C RED NO. 40 (UNII: WZB9127XOA)		
GLYCERIN (UNII: PDC6A3C0OX)		
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
WATER (UNII: 059QF0KO0R)		
SODIUM BENZOATE (UNII: OJ245FE5EU)		
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)		
SUCRALOSE (UNII: 96K6UQ3ZD4)		
TRIACETIN (UNII: XHX3C3X673)		
XANTHAN GUM (UNII: TTV12P4NEE)		
SORBITOL (UNII: 506T60A25R)		

Product Characteristics			
Color	RED	Score	
Shape		Size	
Flavor	BERRY	Imprint Code	
Contains			

I	Packaging				
4	tem Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:72288-723- 34	1 in 1 CARTON	03/04/2021		
1		237 mL in 1 BOTTLE; Type 0: Not a Combination Product			

Marketing Information				
Marketing	Application Number or Monograph	Marketing Start	Marketing End	
Category	Citation	Date	Date	

OTC monograph final part341 03/04/2021

Labeler - Amazon.com Services LLC (128990418)

Revised: 3/2021 Amazon.com Services LLC