

HEADACHE RELIEF EXTRA STRENGTH- acetaminophen, aspirin, caffeine tablet Walgreens

Drug Facts

Active ingredients (in each caplet)

Acetaminophen 250 mg

Aspirin 250 mg (NSAID)*

Caffeine 65 mg

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever

Pain reliever

Pain reliever aid

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - a cold
 - arthritis
 - muscular aches
 - toothache
 - premenstrual and menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives

- facial swelling
- shock
- asthma (wheezing)

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 8 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

- you have liver disease
- stomach bleeding warning applies to you
- you have asthma
- you are taking a diuretic
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis. or kidney disease

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout, or arthritis
- taking any other drug, or are under a doctor's care for any serious condition

Stop use and ask doctor if

- an allergic reaction occurs, Seek medical help right away
 - any new symptoms appear.
- you experience any of the following signs of stomach bleeding
 - feel faint

- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- ringing in the ears or loss of hearing occurs
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- painful area is red or swollen

these could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets every 6 hours; do not take more than 8 caplets in 24 hours
- children under 12 years: ask a doctor

Other information

- store between 20°-25°C (68°-77°F)

Inactive ingredients

colloidal silicon dioxide, corn starch, hypromellose, microcrystalline cellulose, polyethylene glycol, povidone, stearic acid, talc

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to the active ingredients in Excedrin® Extra Strength††

Headache Relief

ACETAMINOPHEN / ASPIRIN (NSAID)

PAIN RELIEVER / CAFFEINE / PAIN RELIEVER AID

EXTRA STRENGTH

CAPLETS

††This product is not manufactured or distributed by GSK Consumer Healthcare distributor of Excedrin® Extra Strength.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

DISTRIBUTED BY: WALGREENS CO., 200 WILMOT RD., DEERFIELD, IL 60015

Package Label

VALUE SIZE NDC 0363-1150-25
Walgreens
Compare to the active ingredients in Excedrin® Extra Strength!
Headache Relief
ACETAMINOPHEN / ASPIRIN (NSAID) / PAIN RELIEVER / CAFFEINE / PAIN RELIEVER AID
EXTRA STRENGTH
CAPLETS
250 CAPLETS ACTUAL SIZE

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts
Active ingredients (in each caplet)
Acetaminophen 250 mg
Aspirin 250 mg (NSAID)
Caffeine 65 mg
*acetaminoid, anti-inflammatory drug

Uses
Temporarily relieves minor aches and pains due to:
■ headache ■ a cold ■ arthritis ■ muscular aches ■ toothache
■ premenstrual and menstrual cramps

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction, which may include:
■ hives ■ facial swelling ■ shock ■ asthma (wheezing)
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin redness ■ blisters ■ rash
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
■ more than 8 caplets in 24 hours, which is the maximum daily amount
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older

Drug Facts (continued under label)
Walgreens Pharmacist Recommended
Walgreens Pharmacist Survey
††This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Excedrin® Extra Strength.
DISTRIBUTED BY: WALGREENS CO., 200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED walgreens.com ©2020 Walgreen Co.
REVO718-F
REV-0920

how2recycle.info
Recycle A Plastic Cap
PLASTIC BOTTLE
ITEM 905195 W10064-0920-F
3 11917 17718 2
Lot No.:
Exp. Date:
PLD-H6814
LBR000063
FEEL HERE

Drug Facts (continued)
■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed
Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.
Do not use ■ if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
Ask a doctor before use if ■ you have liver disease ■ stomach bleeding warning applies to you ■ you have asthma ■ you are taking a diuretic ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
Ask a doctor or pharmacist before use if you are
■ taking a prescription drug for diabetes, gout, or arthritis
■ taking any other drug or are under a doctor's care for any serious condition
Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away. ■ any new symptoms appear ■ you experience any of the following signs of stomach bleeding ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ ringing in the ears or loss of hearing occurs ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days ■ painful area is red or swollen
These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not use more than directed ■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 2 caplets every 6 hours; do not take more than 8 caplets in 24 hours ■ children under 12 years: ask a doctor

Other information ■ store between 20°-25°C (68°-77°F)
Inactive ingredients colloidal silicon dioxide, corn starch, hypromellose, microcrystalline cellulose, polyethylene glycol, polydioxane, stearic acid, talc

Questions or comments?
Call 1-877-755-9858 Monday-Friday 9AM-5PM EST

WALGREENS Headache Relief Extra Strength

HEADACHE RELIEF EXTRA STRENGTH

acetaminophen, aspirin, caffeine tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-1150
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	

Product Characteristics

Color	white	Score	no score
Shape	CAPSULE	Size	17mm
Flavor		Imprint Code	S53
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-1150-25	250 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/30/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	07/30/2020	

Labeler - Walgreens (008965063)