

# **ACETAMINOPHEN- acetaminophen tablet**

## **Prostat First Aid**

-----

### **PROSTAT Acetaminophen 325mg**

#### **Active Ingredient**

Acetaminophen USP, 325mg (in each tablet)

#### **Purpose**

Pain Reliever / Fever Reducer

#### **Do Not Use**

With any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

If you are allergic to acetaminophen or any of the inactive ingredients in this product.

#### **Uses**

Temporarily relieves minor aches and pains due to

Common cold

headache

minor pain of arthritis

toothache

muscular aches

premenstrual and menstrual cramps

temporarily relieves fever

#### **Warnings**

Liver warning

This product contains acetaminophen.

The maximum daily dose of this product is 10 tablets (3,250 mg) in 24 hours for adults or 5 tablets (1,625 mg) in 24 hours for children. Severe liver damage may occur if:

- adults take more than 4000mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours, which is the maximum daily amount.
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product
-

## Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include

- skin reddening
- blisters
- rash

if a skin reaction occurs stop use and seek medical help right away

## **Ask a doctor before use if you have**

Liver Disease

## **Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin.

## **Keep out of reach of children**

Keep out of reach of children, overdose warning.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222).

Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## **Directions**

Do not take more than directed. (see overdose warning).

Adults and children 12 years and older

- Take 2 tablets every 4 to 6 hours while symptoms last
- Do not take more than 10 tablets in 24 hours unless directed by a doctor
- Do not use for more than 10 days unless directed by a doctor

Children 6 years to under 12 years:

- take 1 tablet every 4 to 6 hours while symptoms last
- do not take more than 5 tablets in 24 hours
- do not use for more than 5 days unless directed by a doctor

Children under 6 years of age:

- Ask a doctor

## **Stop Use and ask a doctor if**

- Pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be a sign of a serious condition

**If pregnant or breast-feeding**

Ask a health professional before use

**Overdose Warning**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as children even if you do not notice symptoms.

**Other Information**

- Store at 20°C - 25°C (68°F - 77°F)

**Inactive Ingredients**

hydroxypropyl methyl cellulose, polyethylene glycol, povidone K-30, Pregelatinized starch, Sodium starch glycoate, Stearic Acid

**Questions or Comments?**

Call 1-888-900-2920

**Distributed By:**

Distributed By:

ProStat First Aid

La Verne, CA. 91750 USA

\*This product is not manufactured or distributed by the Johnson & Johnson Corporation, owner of the registered trademark Tylenol®

Made in the USA

**Label**

PACKET

FRONT

Acetaminophen

2 Tablets, 325mg each

Pain Reliever/Fever Reducer  
for Adults

Active ingredient (in each tablet)

Acetaminophen USP, 325mg

Purpose

Pain reliever/fever reducer

USES:

■ temporarily relieves minor aches and pains due to:  
■ headache ■ muscular aches ■ backache ■ minor pain of  
arthritis ■ the common cold ■ toothache ■ premenstrual and  
menstrual cramps ■ temporarily reduces fever. **WARNINGS:**  
**Liver warning:** This product contains acetaminophen. The  
maximum daily dose of this product is 10 tablets (3,250 mg) in  
24 hours for adults or 5 tablets (1,625 mg) in 24 hours for  
children. Severe liver damage may occur if ■ adult takes more  
than 4,000 mg of acetaminophen in 24 hours ■ child takes  
more than 5 doses in 24 hours, which is the maximum daily  
amount ■ taken with other drugs containing acetaminophen  
■ adult has 3 or more alcoholic drinks every day while using  
this product. **Allergy alert:** acetaminophen may cause  
severe skin reactions. Symptoms may include: ■ skin reddening  
■ itching ■ rash. If a skin reaction occurs, stop use and seek  
medical help right away. If pregnant or breast-feeding, ask  
a health professional before use. **KEEP OUT OF REACH** ▶

BACK

**OF CHILDREN. OVERDOSE WARNING:** In case of  
overdose, get medical help or contact a Poison Control  
Center right away. (1-800-222-1222) Quick medical  
attention is critical for adults as well as for children even if  
you do not notice any signs or symptoms. **DIRECTIONS:**  
■ do not take more than directed (see overdose  
warning), adults and children 12 years and over:  
■ take 2 tablets every 4 to 6 hours while symptoms last  
■ do not take more than 10 tablets in 24 hours, unless  
directed by a doctor ■ do not use for more than 10 days  
unless directed by a doctor **children 6 years to under 12  
years:** ■ take 1 tablet every 4 to 6 hours while symptoms  
last ■ do not take more than 5 tablets in 24 hours ■ do not  
use for more than 5 days unless directed by a doctor  
**children under 6 years:** ■ ask a doctor. **OTHER  
INFORMATION:** store between 20-25°C (68-77°F).  
**QUESTIONS OR COMMENTS?** call 888-900-2920

Do not use if printed packet is torn or punctured.

READ AND KEEP CARTON FOR COMPLETE  
WARNINGS AND INFORMATION

Distributed by:  
ProStat First Aid, La Verne, CA 91750

ACETAMINOPHEN

acetaminophen tablet

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58228-0061
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	325 mg
Inactive Ingredients			
Ingredient Name			Strength

STEARIC ACID (UNII: 4ELV7Z65AP)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
STARCH, CORN (UNII: O8232NY3SJ)				
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)				
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)				
POVIDONE K30 (UNII: U725QWY32X)				
Product Characteristics				
Color	white	Score	no score	
Shape	ROUND	Size	12mm	
Flavor		Imprint Code	G323	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58228-0061-1	10000 in 1 BAG	08/06/2025	
1		2 in 1 PACKET; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013		08/06/2025	

**Labeler** - Prostat First Aid (061263699)

Revised: 8/2025

Prostat First Aid