

# INFANTS IBUPROFEN- ibuprofen suspension Walgreens

-----  
**Ibuprofen Oral Suspension, USP**

## ***Drug Facts***

### ***Active ingredient (in each 1.25 mL)***

Ibuprofen 50 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

### ***Purpose***

Pain reliever/Fever reducer

### ***Uses***

temporarily:

- reduces fever
- relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

### ***Warnings***

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Sore throat warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

**Do not use**

- if the child has ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

**Ask a doctor or pharmacist before use if the child is**

- under a doctor's care for any serious condition
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- side effects occur. You may report side effects to FDA at 1-800-FDA-1088.
- child experiences any of the following signs of stomach bleeding:
  - feels faint
  - vomits blood
  - has bloody or black stools
  - has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

## Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away

### **Directions**

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- shake well before using
- find right dose on chart below. If possible, use weight to dose; otherwise use age.
- mL = milliliter
- measure with the dosing device provided. Do not use with any other device.
- dispense liquid slowly into the child's mouth, toward the inner cheek
- if needed, repeat dose every **6 to 8 hours**
- do not use more than **4 times a day**

### **Dosing Chart**

Weight (lb)	Age (mos)	Dose (mL)
under 6 mos		ask a doctor
12 to 17 lbs	6 to 11 mos	1.25 mL
18 to 23 lbs	12 to 23 mos	1.875 mL

### **Other information**

- store between 20° to 25°C (68° to 77°F)

### **Inactive ingredients**

carboxymethylcellulose sodium, citric acid anhydrous, glycerin, microcrystalline cellulose, natural and artificial berry fruit punch type flavor, polysorbate 80, purified water, sodium benzoate, sorbitol solution, sucrose, and xanthan gum.

### **Questions or Comments?**

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

### **Principal display panel**

Compare to the active ingredient in Concentrated Motrin® Infants' Drops††

**infants'**

**Ibuprofen**

CONCENTRATED IBUPROFEN

ORAL SUSPENSION, USP (NSAID)

50 mg PER 1.25 mL

PAIN RELIEVER / FEVER REDUCER

Alcohol Free

Dye-Free

Non-staining

- Lasts up to 8 hours

AGES 6 -23 MONTHS

BERRY FLAVOR

Use only with enclosed syringe

FL OZ (mL)

IMPORTANT: READ ALL PRODUCT INFORMATION BEFORE USING. KEEP THIS BOX FOR IMPORTANT INFORMATION. THIS PRODUCT IS INTENDED FOR USE IN CHILDREN AGES 6 MONTHS TO 23 MONTHS OF AGE.

††This product is not affiliated with, manufactured by, or produced by the makers or owners of Motrin®.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR PRINTED FOIL UNDER CAP IS BROKEN OR MISSING.

DISTRIBUTED BY: WALGREEN CO.

200 WILMOT RD., DEERFIELD, IL 60015

**Package label**

NDC 0363-5100-01

**IMPORTANT: READ ALL PRODUCT INFORMATION BEFORE USING. KEEP THIS BOX FOR IMPORTANT INFORMATION. THIS PRODUCT IS INTENDED FOR USE IN CHILDREN AGES 6 MONTHS TO 23 MONTHS OF AGE.**

**Drug Facts**  
**Active Ingredient (in each 1.25 mL)**  
 Ibuprofen 50 mg (NSAID)\* — Pain reliever/fever reducer  
 \*nonsteroidal anti-inflammatory drug

**Uses** temporarily:  
 ■ reduces fever  
 ■ relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

**Warnings**  
**⚠️ Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
 ■ hives ■ facial swelling  
 ■ asthma (wheezing) ■ shock  
 ■ skin redness ■ rash ■ blisters  
 If an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:  
 ■ has had stomach ulcers or bleeding problems  
 ■ takes a blood thinning (anticoagulant) or steroid drug  
 ■ takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
 ■ takes more or for a longer time than directed  
**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Drug Facts (continued)**

**Sore throat warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

**Do not use**  
 ■ if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer  
 ■ right before or after heart surgery

**Ask a doctor before use if**  
 ■ stomach bleeding warning applies to your child  
 ■ child has a history of stomach problems, such as heartburn  
 ■ child has problems or serious side effects from taking pain relievers or fever reducers  
 ■ child has not been drinking fluids  
 ■ child has lost a lot of fluid due to vomiting or diarrhea  
 ■ child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke  
 ■ child has asthma ■ child is taking a diuretic

**Ask a doctor or pharmacist before use if the child is**  
 ■ under a doctor's care for any serious condition  
 ■ taking any other drug

**When using this product**  
 ■ take with food or milk if stomach upset occurs

**Stop use and ask a doctor if** ■ side effects occur. You may report side effects to FDA at 1-800-FDA-1088.  
 ■ child experiences any of the following signs of stomach bleeding:  
 ■ feels faint ■ vomits blood ■ has bloody or black stools  
 ■ has stomach pain that does not get better  
 ■ child has symptoms of heart problems or stroke:  
 ■ chest pain ■ trouble breathing  
 ■ weakness in one part or side of body  
 ■ slurred speech ■ leg swelling  
 ■ the child does not get any relief within first day (24 hours) of treatment  
 ■ fever or pain gets worse or lasts more than 3 days  
 ■ redness or swelling is present in the painful area  
 ■ any new symptoms appear

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**  
 ■ this product does not contain directions or complete warnings for adult use  
 ■ do not give more than directed  
 ■ shake well before using

**Drug Facts (continued)**

■ find right dose on chart below. If possible, use weight to dose; otherwise use age.  
 ■ mL = milliliter  
 ■ measure with the dosing device provided. Do not use with any other device.  
 ■ dispense liquid slowly into the child's mouth, toward the inner cheek  
 ■ if needed, repeat dose every 6 to 8 hours  
 ■ do not use more than 4 times a day

**Dosing Chart**

Weight (lb)	Age (mos)	Dose (mL)
under 6 mos		ask a doctor
12 to 17 lb	6 to 11 mos	1.25 mL
18 to 23 lb	12 to 23 mos	1.875 mL

**Other information**  
 ■ store between 20° to 25°C (68° to 77°F)

**Inactive ingredients**  
 carboxymethylcellulose sodium, citric acid anhydrous, glycerin, microcrystalline cellulose, natural and artificial berry fruit punch type flavor, poly sorbate 80, purified water, sodium benzoate, sorbitol solution, sucrose, and xanthan gum

**Questions or comments?**  
 Call 1-877-753-3935  
 Monday-Friday 9AM-5PM EST

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR PUNCHED FOR UNDER CAP OR BROKEN OR MISSING.**

Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands. This product is not affiliated with, manufactured by, or produced by the makers or owners of Motrin®.

DISTRIBUTED BY: WALGREEN CO.  
 200 WILMOT RD., DEERFIELD, IL 60015  
 100% SATISFACTION GUARANTEED  
 walgreens.com ©2021 Walgreen Co.  
 LB8623 Rev. 01/06/22 W3ORG0222-F REV-0722



Compare to the active ingredient in Concentrated Motrin® Infants' Drops\*

# Infants' Ibuprofen

CONCENTRATED IBUPROFEN ORAL SUSPENSION, USP (NSAID)  
 50 mg per 1.25 mL  
 PAIN RELIEVER/FEVER REDUCER

Alcohol Free Dye-Free

Non-staining

• Lasts up to 8 hours

AGES 6-23 MONTHS



1 FL OZ (30 mL)

Berry flavor  
 USE ONLY WITH ENCLOSED SYRINGE

Lot No.:  
 Exp. Date:

VOID AQUEOUS

ITEM 268964 W10041-0822-L  
  
 3 11917 21867 0

PDA-STBA F030815

## WALGREENS Infants ibuprofen Berry Flavor

INFANTS IBUPROFEN			
ibuprofen suspension			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-5100
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	50 mg in 1.25 mL	
Inactive Ingredients			
Ingredient Name	Strength		

<b>CARBOXYMETHYLCELLULOSE SODIUM</b> (UNII: K679OBS311)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCROSE</b> (UNII: C151H8M554)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	BERRY	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-5100-01	1 in 1 CARTON	12/04/2020	03/01/2027
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079058	12/04/2020	03/01/2027

**Labeler** - Walgreens (008965063)