

## **ASPIRIN- aspirin tablet**

### **PROSTAT FIRST AID**

-----

### **PROSTAT Aspirin 325mg (NSAID)**

**\*nonsteroidal anti-inflammatory drug**

#### **Active Ingredient**

Aspirin 325mg (NSAID)

Nonsteroidal anti-inflammatory drug

#### **Purpose**

Pain Reliever / Fever Reducer

#### **Do Not Use**

- If you are allergic to aspirin or any other pain releiver / fever reducer
- If you have ever had an allergic reaction to this product or any of its ingredients

#### **Uses**

Temporarily relieves:

Headache

Pain and fever of colds

toothache

menstrual pain

muscle pain

minor pain of arthritis

#### **Warnings**

##### **Reye's Syndrome**

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behaviour with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's Syndrome, a reare but serious illness.

##### **Allergy Alert**

Aspirin may cause severe allergic reaction, which may include, facial swelling, hives, shock, asthma (wheezing).

##### **Stomach bleeding warning**

This product contains an NSAID which may cause severe stomach bleeding. The chance is higher if you

- Have had stomach ulcers or bleeding problems
- Take blood thinning (anticoagulant) or steroid drug
- Are age 60 or older
- take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen or others)
- Take more or for a longer time than directed
- Have 3 or more alcoholic drinks every day while using this product

**Ask a doctor before use if**

- Stomach bleeding warning applies to you.
- You have a history of stomach problems, such as heartburn
- You have asthma
- You have high blood pressure, heart disease, liver cirrhosis or kidney disease.
- You are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug for gout, diabetes, arthritis

**Directions**

Do not take more than directed. See overdose warning

Drink a full glass of water with each dose

Adults and children 12 years and older

- Take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor.

Children under 12 years

- Do not use unless directed by a doctor

**Stop Use and ask a doctor if**

- An allergic reaction occurs. Seek medical help right away
- You experience any of the following signs of stomach bleeding, feel faint, vomit blood, have bloody stools, have stomach pain that does not get better
- Ringing in the ears or loss of hearing occurs, pain gets worse or lasts more than 10 days, fever gets worse or lasts more than 3 days.
- New symptoms occur,
- Redness or swelling is present
- These could be signs of a serious condition

**If pregnant or breast-feeding**

Ask a health professional before use. It is especially important not to use aspirin at 20

weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Other Information**

- Store at 25°C (77°F)
- Excursions permitted between 15°C-30°C (59°F-86°F)
- See end flap for expiration date and lot number
- Use by expiration date on package
- Tamper Evident: Do not use if packet is open or torn
- Read and Keep carton for complete warnings and information

**Inactive Ingredients**

Corn Starch, Hyperomelose, Polyethylene glycol, propylene glycol

**Keep out of reach of children**

In case of overdose, get medical help or contact a poison control center right away.

**Questions or Comments?**

Call 1-888-900-2920

**Distributed By:**

Distributed By:

ProStat First Aid

La Verne, CA. USA

1-888-900-2920

\*This product is not manufactured or distributed by the Bayer Healthcare LLC. owner of the registered trademark Bayer®

Made in the USA

**Label**

PACKET

FRONT

Aspirin

2 Tablets, 325mg each

Pain Reliever/Fever Reducer

(NSAID)

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)\*

\*Nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

● temporarily relieves ● headache ● pain and fever of colds ● toothache ● menstrual pain ● muscle pain ● minor pain of arthritis ● **Warnings: See carton for complete warnings. Reye's Syndrome:** Children and teenagers who have or are experiencing than chicken pox or the like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. ● **Allergy alert:** Aspirin may cause severe allergic reaction, which may include ● facial swelling ● hives ● difficulty breathing ● **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ● have had stomach ulcers or bleeding problems ● take a blood thinning (anticoagulant) or steroid drug ● are age 60 or older ● use other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ● take more or for a longer time than directed ● use 3 or more alcoholic drinks every day while using this product. ● **Do not use if** ● you are allergic to aspirin or any other pain reliever/fever reducer ● if you have ever had an allergic reaction to this product or any of its ingredients ● **Ask a doctor before use if** ● stomach bleeding warning applies to you ● you have a history of stomach problems, such as heartburn ● you have asthma ● you have high blood pressure, heart disease, liver problems or kidney disease ● you are taking a diuretic ● **Ask a doctor or pharmacist before use if you are taking a prescription drug for** ● gout ● diabetes ● arthritis ● **Stop use and ask a doctor if** ● you

BACK

allergic reaction occurs. Seek medical help right away ● you experience any of the following signs of stomach bleeding ● feel faint ● vomit blood ● have bloody or black stools ● have stomach pain that does not get better ● ringing in the ears or a loss of hearing occurs ● pain gets worse or lasts more than 10 days ● fever gets worse or lasts more than 3 days ● new symptoms occur ● redness or swelling is present. These could be sign of a serious condition. ● **If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. ● **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. ● **DIRECTIONS ● do not take more than directed ●** drink a full glass of water with each dose ● **adults and children 12 years and older:** 1 tablet 1 or 2 tablets every 4 hours or 3 tablets every 6 hours to a maximum of 12 tablets in 24 hours unless directed by a doctor ● **children under 12 years:** do not use unless directed by a doctor ● **OTHER INFORMATION ●** use by expiration date on package ● store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ● **INACTIVE INGREDIENTS** Croscarmellose, Hypromellose, Polyethylene glycol, Pregelatinized starch ● **QUESTIONS OR COMMENTS?** Call 888-888-3333

**TAMPER EVIDENT: DO NOT USE IF PACKET IS OPEN OR TORN**

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

Distributed by:  
ProStat First Aid, La Verne, CA 91750

MADE IN THE USA

ASPIRIN

aspirin tablet

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58228-0071
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)		ASPIRIN	325 mg
Inactive Ingredients			
Ingredient Name			Strength

STARCH, CORN (UNII: O8232NY3SJ)				
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
Product Characteristics				
Color	white	Score	score with uneven pieces	
Shape	ROUND	Size	10mm	
Flavor		Imprint Code	44157	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58228-0071-1	10000 in 1 BAG	08/05/2025	
1		2 in 1 PACKET; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M013	08/04/2025	

**Labeler** - PROSTAT FIRST AID (061263699)

Revised: 8/2025

PROSTAT FIRST AID